

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
13 December 2001 (13.12.2001)

PCT

(10) International Publication Number  
**WO 01/94599 A1**

- (51) International Patent Classification<sup>7</sup>: C12N 15/63, A61K 39/08 (74) Agents: NILSSON, Brita et al.; Stockholms Patentbyrå Zacco AB, P.O. Box 23101, S-104 35 Stockholm (SE).
- (21) International Application Number: PCT/SE01/01280 (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (22) International Filing Date: 7 June 2001 (07.06.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
0002139-4 7 June 2000 (07.06.2000) SE  
0101479-4 26 April 2001 (26.04.2001) SE
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): SMITTSKYDDSIINSTITUTET [SE/SE]; S-171 82 Solna (SE).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): BURMAN, Lars, G. [SE/SE]; Södermannagatan 64A, S-116 65 Stockholm (SE). ÅKERLUND, Thomas [SE/SE]; Salagatan 15A, S-753 30 Uppsala (SE). MUKHERJEE, Kakoli [IN/IN]; 324B, B.B. Chatterjee Road, Kasba, Calcutta 700 042 (IN). KATAGIHALLIMATH, Nainesh [IN/IN]; 57, IInd Main, III stage, IIIrd Block, Basaweshwarnagar, Bangalore 560 079 (IN).
- Published:**  
— with international search report  
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: GENE EXPRESSION CASSETTE AND ITS USE

(57) Abstract: A gene expression cassette comprising a secretory leader sequence encoding a signal peptide from *Clostridium difficile* having an amino acid sequence selected from SEQ ID NO: 1-12 and signal peptides of analogous exported clostridial N-acetylmuramoyl-L-alanine amidase-like proteins, linked to a DNA sequence encoding a heterologous polypeptide, and optionally a part of SEQ ID NO: 22-233, is described. The gene expression cassette is inserted into a vector, and the vector is used to transform a host organism. Compositions, formulations, vaccines and medicaments based on spores of such engineered host organisms are used e.g. for colonization of a mammal.

BEST AVAILABLE COPY

WO 01/94599 A1

### Gene expression cassette and its use

The present invention relates to a gene expression cassette and in particular to the use of the cassette in methods for presenting polypeptides on the surface of bacterial cells and/or secreting them into the surroundings of the latter. The invention further relates to gene expression constructs that are used to transform bacterial host cells. Uses of the invention include immunisation, in particular mucosal immunisation, induction of immunological tolerance and anti-tumour therapy in humans and animals. The intended vaccines and anti-cancer agents will also make use of bacterial spores produced by clostridia, *e.g. Clostridium difficile*, for both industrial production of the vaccine and for local production of the desired polypeptides at the body sites desired.

Vaccines against infection represent the greatest advance in medicine with unparalleled impact on morbidity and mortality at relatively low cost. Despite their cost-effectiveness, the cost associated with modern vaccines is still of concern and limits their use, particularly in developing countries. A number of factors contribute to the cost of injectable vaccines including the requirements for vaccines with defined sub-cellular components, for purity and sterility of the vaccine preparations, for testing of administration routes and combinations with different adjuvants, for maintaining the cold chain in distribution of the vaccines, and for using sterile syringes and needles. The need for repeated vaccinations also contributes to increased costs. Furthermore, for many infectious diseases a vaccine has not yet been made available

Also, in recent years great interest has been shown in mucosal immunisation, i.e. the exposure of mucosal surfaces to an antigen to elicit a general humoral and mucosal immune response, i.e. also at distant sites (Mucosal Immunology, Ed. P.L.Ogra et al., Acad. Press, 1999). Although this field is in its infancy, two such products are so far on the market, namely the oral polio vaccine and an oral (drinkable) vaccine against cholera and diarrhea due to *Escherichia coli*. The latter is an inactivated vaccine containing killed *Vibrio cholerae* organisms plus the cholera toxin subunit B which is non-toxic, immunogenic and shared between the cholera toxin and the toxin of "enterotoxigenic *E. coli*" (ETEC), the main cause of "travellers diarrhea". It is hoped that mucosal immunisation via for example intranasal or peroral administration of the antigen will provide a real alternative to injectable vaccines. As

most microbes invade humans and animals *via* mucous membranes we anticipate that the mucosal route will turn out to be the superior alternative for vaccination in many instances.

The scientific consensus at this point appears to be that live vaccines are potentially superior candidates for single-dose long lasting vaccination, because the carrier organism will continue to produce the antigen and boost immunity *in vivo*. However, experience with the bacterial carriers currently studied, for example the intestinal bacteria *Salmonella* and *E. coli* is that they are not generally classified as safe and are further difficult to distribute. Therefore, probiotic organisms such as *Lactobacilli* have been proposed as suitable carriers of foreign antigens. Problems have been identified, these include limited shelf-life unless freeze-dried which reduces viability, and difficulties in colonising the gut of the human recipient and evoking an immune response.

A classical way of enhancing the immunogenicity of vaccines is co-administration of so-called adjuvants. These may be molecules such as aluminium hydroxide or lipid vesicles that increase the exposure time for the vaccine by slowing its removal from the site of injection or "danger molecules" of microbial origin that increase the immune response in a non-specific way. Thus, in recent years it has been found that adjuvants also act by evoking production of immunomodulatory peptides called cytokines and chemokines (Brewer JM, Alexander J, Cytokines Cell Mol Ther 4:233-246, 1997. Ulanova M, Classical and non-classical antigen-presenting molecules in immune responsiveness, Thesis, Göteborg University, Sweden, 2000, ISBN 91-628-4228-5). Due to possible adverse effects the use of cytokines themselves as adjuvants must await a better understanding of optimal selection and dosing of these molecules together with vaccines. It is likely that applying cytokine adjuvants by the mucosal route is less critical than parenteral administration from a side effect point of view.

Interest has also been shown in the extracellular presentation of foreign protein antigens by bacteria, included as part of bacterial surface layer proteins. A surface layer (S-layer) protein is herein defined as any molecule of proteinaceous nature, including e.g. protein, glyco- or lipoprotein occurring in the outer layer of a bacterium and capable of being exposed on the surface of the bacterium. S-layer proteins are a main constituent of the cell wall of some gram-positive bacterial genera. They may be continuously and spontaneously produced in larger amounts than any other class of protein in the cell. WO-95/19371 describes a fusion protein of at least a part of a S-layer protein and a heterologous peptide, the intention being that the polypeptide is expressed and presented on the surface of the cell. A range of

bacterial hosts is mentioned including *Staphylococcus*, *Streptococcus*, *Bacillus*, *Clostridium* and *Listeria*. A preference for *Bacillus* is stated and the examples use *B. sphaericus*. Elsewhere, WO-97/28263 describes processes for the recombinant preparation of S-layer proteins in gram-negative host cells. It is suggested that these proteins could include antigenic species. FR-A-2778922 describes the use of genes which regulate the synthesis of toxin products in *Clostridium* bacteria, to produce polypeptides.

We first set out to investigate toxin release and to identify other extracellular proteins produced by *Clostridium difficile* (*C. difficile*). *C. difficile* is an anaerobic spore forming pathogen causing *C. difficile* associated diarrhea (CDAD) and pseudomembranous colitis (PMC) by producing two toxins, A and B.

In contrast to studies by Kamiya et al (J.Med.Microbiol.,1992, 37, 206-210) and Ketley et al (J.Med.Microbiol.,1984, 18, 385-391) we found that the accumulation of extracellular toxins is not accompanied by cell lysis suggesting a toxin export mechanism.

Our first detailed analysis (described in Example 9 below) of proteins occurring extracellularly as well as in the cell wall and membrane fraction of strains VPI 10643 and 630 and analysis of the DNA/genes encoding these proteins revealed a genomic segment containing eighteen genes (Figure 1A). Seven proteins (ORF1,3,5-7,9 and 11), when compared with publicly available sequence showed some homology to N-acetyl muramoyl L-alanine amidase (CwlB/LytC) and modifier protein of major autolysin (LytB) from *B. subtilis*, as well as S-layer proteins from *Lactobacillus spp* (Tables 1 and 4). The amidase motif was located either at the C-terminal or the N-terminal end of the S-layer protein ORFs (see examples in Figure 1B). Other ORFs showed similarity to genes involved in polypeptide secretion (ORF2/*secA*), polysaccharide and capsule synthesis, and possibly glucosylation of the S-layer proteins. Further database searches indicated that the amidase motif confers anchorage of the S-layer proteins to the clostridial cell wall peptidoglycan-teichoic acid.

A search in the revised *C. difficile* database revealed five additional genes upstream of ORF1 which had similarities to the previously found ones, i.e. they had a two-domain architecture one showing homology to the CwlB/LytC and LytB proteins. These ORFs thus had the putative cell wall binding amidase motif typical of the other S-layer ORFs and were designated D, E, G, H and I by us (Fig. 2 and Table 1).

Significantly, we found that the N-termini of all S-layer ORFs contained a typical signal peptide for export via *sec*-dependent secretion and that for e.g. ORF1 the predicted signal peptide cleavage site (Figure 3) was identical to that found in the protein sequence.



Furthermore, the secreted ORF1 product was further cleaved into two peptides in strain 630, the C-terminal one containing the N-acetyl muramoyl L-alanine amidase like sequence (Figure 1B and Example 9). Although we identified the cleavage point, the kinetics and precise mechanism of this proteolytic event remains unknown. It is likely that the S-layer protein cleavage product containing the amidase motif provides cell-wall binding, whereas the other peptide showing more sequence variability is more surface exposed and providing antigen variation between strains (serotypes). We found no significant match between sequences within ORF1 and the S-layer homology motif (SLH domain) found in most presently known S-layer proteins although a weak similarity to an S-layer protein from *Lactobacillus helveticus* was found for the N-terminal part of ORF1 (Figure 1B). Also, apart from the shared amidase motif, we found no significant homology between the different S-layer ORFs that could suggest how the variable cleavage product(s) get anchored to the cell-wall binding peptide (inner S-layer) or to each other to form the outermost layer recently distinguished by electron microscopy (Cerquetti M et al. Characterization of surface layer proteins from different *C. difficile* clinical isolates. Microb Pathogenesis 28: 363-372, 2000).

Our subsequent analysis of 21 *C. difficile* serogroup type strains, probably representing all major genetic lineages of the species, indicated that a gene segment corresponding to ORF1 and its upstream region plus ORF.2 is generally present in *C. difficile* (Example 1C).

The N-terminal part of ORF6 showed homology to eukaryotic cysteine proteases (Fig. 1B). ORF5 has been suggested to be involved in adhesion to epithelial cells (Abstract; The Third International Meeting on the Molecular Genetics and Pathogenesis of the Clostridia, June 8-11, 2000, Chiba, Japan).

The present invention is based, at least in part, on the above discoveries. We have identified and developed a polypeptide expression and secretion system that may be used to produce a desired polypeptide on the surface of and/or into the surroundings of bacteria, for introduction into an appropriate mammal. The system may be used for example to initiate mucosal vaccination. A particular advantage of the system is that it may be used with any convenient *Clostridium* species, independently of any normal S-layer protein production. Furthermore, in case of *C. difficile* it is possible to use strains lacking the 5-gene toxicity cassette encoding the two major virulence factors toxins A and B and thus avoid the risk of CDAD when administering the engineered peptide producing strains to humans aged 2-4 years or more (neonates and small children are insensitive to the toxins) or animals.

Therefore, in a first aspect of the invention we provide a gene expression cassette comprising a secretory leader sequence selected from any one of ORF1, ORF3, ORF5-7, ORF9 or ORF11 (SEQ ID NO: 1 - 7) (c.f. Figure 1 and Table 1) of *C. difficile* strain 630 linked to a DNA sequence encoding a heterologous polypeptide. Alternatively, the secretory leader sequence is from any one of ORF D, E, G, H and I (SEQ ID NO: 8 - 12) (cf. Figure 2 and Table 1) or from any analogous S-layer ORF taken from any *C. difficile* strain.

By "heterologous" we mean a nucleic acid sequence or protein not native to the clostridial strain being used.

Use of each of the secretory leader sequences mentioned above represents a separate and independent aspect of the invention. The secretory leader sequence is preferably from ORF1.

A recent publication by Karjalainen et al, Infection and Immunity, May 2001, p3442-3446 provides confirmation and analysis of most of the ORFD - ORFI - ORF1 - ORF11 S-layer gene cluster in *C. difficile* strain 630. The nucleotide and polypeptide sequences disclosed by Karjalainen et al are incorporated herein by reference.

In a further aspect of the invention the gene expression cassette further includes a promoter of prokaryotic origin. The promoter is preferably a strong promoter and in general is placed 5' of the secretory leader sequence in the gene cassette.

In a further aspect of the invention the gene expression cassette further includes a DNA sequence encoding at least a functional portion of an S-layer protein of *C. difficile* fused to a nucleic acid coding sequence coding for a heterologous polypeptide such that the resulting fusion polypeptide will be expressed and presented on the outer surface of the host cell harbouring the cassette. If desired the polypeptide can also be released from the bacteria, e.g. by excluding the S-layer amidase motif from the construct (cf. Figure. 3).

In a further aspect of the invention the engineered gene expression cassette optionally further comprises at least a functional part of the secretory (*secA*) gene represented by ORF2. This may be used to complement or replace the function of the normal *C. difficile sec* gene in order to ensure efficient translocation of the peptide(s) produced by the cassette across the cytoplasmic membrane.

An example of a preferred gene expression cassette is conveniently illustrated in Figure 3.

The promoter in the gene expression cassette is conveniently a strong promoter, this may be the native promoter for ORFs 1 - 12 of *C. difficile* of strain 630 (Table 1).

Alternatively, the promoter sequence is from any one of ORF D - I (cf. Table 1), alternatively from any other analogous S-layer ORF from a *C. difficile* strain or from another gene, preferably from this species (see Specific description B1). The promoter may thus be another prokaryotic promoter that is strong, inducible or constitutive, and functional in the polypeptide producing bacterium. In all potential applications a distinct advantage of this cassette is the very large amounts of protein produced and exported.

The gene expression cassette is conveniently placed in a vector or specifically a plasmid carrying a transposon belonging to for example the Tn916, Tn5387 or the Tn5398 families. After transfection of a *C. difficile* host organism these transposons are able to insert themselves into its chromosome thereby making the engineered cassette a stable trait of the bacterium (cf. Figure 4). For other *Clostridia* other vectors may be preferable, e.g. the engineered shuttle plasmid pJIR750. Unlike the *C. difficile* plasmids currently available, this vector can replicate within both an *E. coli* and a *C. perfringens* host and is not dependent on integration of the plasmid into the host chromosome. Any convenient *Clostridium* species may be used, to date over 70 species have been defined by rRNA sequence analysis. These include *C. difficile* and classical pathogens as *C. perfringens*, *C. tetani* and *C. botulinum*, also *C. acetobutylicum* that is being genetically manipulated and used for industrial production of acetic acid and *C. beijerinckii* that has been transformed with *E. coli* genes.

*C. perfringens* is currently the species most amenable to genetic engineering. It is a normal, moderate level, fecal coloniser of most, if not all, humans. *C. difficile* is found in the fecal flora of most newborns, less often in adults but commonly in hospitalized individuals. As *C. difficile* is an early, normally colonising intestinal organism and even toxigenic strains are unable to cause CDAD in newborns and infants up to 2-4 years of age, we believe that recombinant *C. difficile* producing desired antigens and adjuvants is suitable for oral vaccination at any convenient time after birth.

Whereas *C. perfringens* normally produces many toxins about half of wild *C. difficile* strains are genetically non-toxigenic, which may be an advantage from a safety point of view. *C. difficile* toxin negative strains are preferred as host cells for the gene expression cassettes of this invention, at least for individuals aged 2-4 years or more (see above).

The nucleic acid sequence coding for a heterologous polypeptide is placed in the gene expression cassette before or after insertion into a convenient vector or plasmid. The insertion points for the nucleic acid sequence are at the discretion of the skilled scientist, there may be in the variable or in the constant region of the relevant ORF nucleotide sequence. Routine

experimentation may be used to determine convenient and particular insertion points. In Figure 3 we disclose polypeptide cleavage sites that need to be taken into consideration (See Specific description B3).

Examples of convenient plasmids include those mentioned in Figure 4, for example pCI195 and pSMB47. Convenient transposons include those belonging to the Tn916, Tn 5397 and Tn5398 families for transfection into *C. difficile* and for example pJIR750 for *C. perfringens* or other *Clostridia*. Any convenient heterologous nucleic acid sequence may be placed into the gene expression cassette. In a further aspect of the invention we provide a vector or plasmid comprising a gene cassette of the invention.

The vector or plasmid may then be transfected into a convenient host using techniques known in the art (see for example: Gene 82: 327-333, 1989). For *C. difficile* it is at present preferred to introduce the plasmid into a *Bacillus* species such as *B. subtilis* and then transfer the target DNA by filter mating (conjugation) into a convenient *C. difficile* strain (outlined in Fig. 4). This will generally require the use of a conjugative transposon-bearing plasmid such as pCI195 or pSMB47 (J. Antimicrob. Chemother. 35: 305-315, 1995; FEMS Microbiol. Lett., 168: 259-268, 1998; D. Lyras, J. I. Rood, Clostridial genetics, in Gram-positive pathogens, ed. V. A. Fischetti, Am. Soc. Microbiol, 2000). However, we anticipate that further materials and procedures will become available for the direct introduction of plasmids or other foreign DNAs into *Clostridia* and particularly *C. difficile*. For e.g. *C. perfringens* such vectors and techniques are to some extent already available.

In a further aspect of the invention we provide a Clostridial bacterium transformed with a gene expression cassette of the invention encoding the desired fusion peptide or entirely heterologous polypeptide(s).

The transformed Clostridial bacterium, when administered orally to any convenient mammal such as a human or animal will lead to the intestinal colonization, production and presentation of the desired polypeptide particularly in the large bowel that is the natural site of colonization of *C. difficile*. The bowel wall is surrounded by an immense immune apparatus, the so-called Peyer's patches and thus, specialized in mounting immune responses of various types. Large bowel colonization by a clostridial vaccine or peptide producer strain thus enables a much longer immune stimulus than a traditional injection. In contrast to clostridia, the alternative and much studied S-layer producers for vaccine purposes, *Bacillus spp*, are free-living, obligate aerobic bacteria and unable to replicate in the anaerobic bowel lumen and

thus, unable to colonize a recipient mammal. For clostridial colonization and peptide delivery in hypoxic tissues iv administration is used.

It will be appreciated that Clostridia carrying the gene expression cassette of this invention including DNA encoding different heterologous peptides allows the highly efficient production and export of these polypeptides in hypoxic tissues after iv administration, or into the gut, particularly the colon, of the orally colonized individual for a variety of prophylactic or therapeutic uses. Another advantage of this gene cassette for expression of heterologous peptides is its versatility, i.e. that it is normally used to produce and export peptides of varying size and having completely different amino acid sequences, in their N-terminal or C-terminal end.

In further independent aspects of the invention the recombinant gene expression cassette is used to produce in the gut, for example

- (i) peptides and enzymes for therapy and prophylaxis of various diseases, e.g. peptides having specific antimicrobial activity, cytokines against inflammatory bowel disease, and  $\beta$ -lactamases to prevent diarrhea due to antibiotic therapy
- (ii) single, fusion or multiple polypeptide antigens of microbial, animal or mammalian origin for neonatal immune balancing, vaccination against infections, allergy, metabolic or auto-immune disease, cancer, (in)fertility, and drug addiction.
- (iii) carrier molecules (so-called adjuvants) separate or fused to the antigen in order to amplify or modulate the immune response to the antigen in a desired way according to (ii), e.g. a strong IgA response against a mucosal invader.

In a still further aspect of the invention the gene expression cassette of the invention may be used to provide recombinant clostridia for local production of peptides in tissues after iv administration of their spores (see below), for example for the prophylaxis and/or treatment of fibrinolysis in arterial or venous occlusion and/or for revitalising gangrenous and/or necrotic tissue in various diseases. Furthermore, for anti-tumour therapy by local production of

- (i) immune stimulating human peptides for improving tumour host defence,
- (ii) enzymes that convert a pro-drug to a cytostatic agent inside a tumour (thus avoiding systemic side effects)
- (iii) cytotoxins of e.g. bacterial origin to destroy tumour cells

(iv) angiogenesis inhibitors at local concentrations enough to prevent local blood vessel formation and thus, tumour growth

(v) signal transduction inhibitors.

In a further aspect of the invention we provide a pharmaceutical or veterinary composition which comprises a transformed viable *Clostridial* cell with the ability to present and/or to secrete the desired polypeptide together with a pharmaceutically or veterinary acceptable carrier or diluent.

The composition may be formulated as a vaccine. The composition may be administered orally, or intranasally or alternatively, the polypeptide can be isolated, purified and administered parenterally, e.g. subcutaneously or intramuscularly.

The amount of the desired peptide(s) presented and/or secreted by the transformed strain may be modulated in the body by using

(i) promoters with different strength (power),

(ii) a promoter or regulator responding to external stimuli (inducible, e.g. by a specific carbohydrate) normally present in the gut or administered together with the engineered bacterium,

(iii) different dosage regimens (number of bacteria per dose and doses per time period) or

(iv) methods that influence the ability of the strain to colonise and propagate in the gut for convenient periods of time. Relevant factors include ability to compete with other bacteria, adhere to mucosal cells, and to avoid expulsion by local immune response mechanisms. The latter can be achieved by exploiting induction of tolerance to the natural S-layer antigen of the strain in the neonate (see above and below) or by using its putative normal antigenic variation, suggested to us by the presence of the large number of S-layer ORFs present in *C. difficile*, either by allowing the vaccine strain to change serotype during natural long-term colonization or by repeated applications (colonizations) over time of different strains producing the same antigen but having a different serotype antigen.

The transformed *Clostridia* as anaerobic organisms are conveniently produced by fermentation under for example low oxygen tension and purified and recovered as known in the art for native *Clostridia*, for example by washing and freeze-drying. They may be formulated together with excipients as needed, for example magnesium stearate, lactose, or carboxymethyl cellulose, into solid dosage forms, e.g. in capsules, predominantly for oral administration. The dosage forms may be protected against the acidity of the stomach by a

suitable enteric coating, comprising for example Eudragite "S", Eudragite "L", cellulose acetate, cellulose phthalate or hydroxypropyl cellulose. A preferred dosage form comprises freeze-dried transformed *Clostridia* contained in vials or ampoules, optionally under inert gas. Preferably, the transformed *Clostridia* cells are administered orally or intranasally, as an aqueous, reconstituted suspension of the lyophilized cells e.g. in water or physiological saline, optionally with addition of pharmaceutically acceptable buffers, e.g. sodium bicarbonate, phosphate or citrate to keep the pH of the suspension between 6 and 8, preferably between 6.5 and 7.5.

The dosage forms produced as described above may comprise a mixture of viable and non-viable bacteria depending on the process and/or the storage conditions. The viable, transformed *Clostridia* will, after oral administration, become attached to those parts of the gut, for example the lower intestinal tract, which provide appropriate growing conditions and proliferate, producing the desired polypeptide in increasing amounts. This will provide for an enhanced and sustained physiological effect, for example immunisation, of the polypeptide.

If exposure to defined amounts of the polypeptide is desired, non-viable transformed *Clostridia* presenting the polypeptide can be administered. The non-viable cells can be obtained as known in the art, e.g. by exposing the live cells to agents, e.g. heat, formaldehyde, antibiotics or solvents, which kill them. It is also possible to use cell walls (sacculi) or to use S-layer fragments obtained by mechanical or other disruption of the bacterial cells. These agents can be formulated into pharmaceutical and veterinary compositions as described above for live transformed *Clostridia*.

By "secretion" or "release" we mean that the heterologous polypeptide is exported out from the host cell into the surrounding environment as a soluble antigen. This is conveniently achieved by fusing the DNA coding for the polypeptide to a DNA sequence coding for a signal peptide sequence of any one of ORF1, ORF3, ORF5-7, ORF 9 or ORF11 (SEQ ID NO:1-7, cf. Figure 1 and Table1) preferably to that of ORF1 and expressing it as described above under control of a strong promoter and exporting it with the aid of the *sec* gene (ORF2) product. Alternatively, the DNA codes for a signal peptide sequence of any one of ORF D, E, G, H or I (SEQ ID NO: 8-12, cf. Figure 2 and Table 1), or that of any other suitable secreted bacterial protein.

By "presentation" we mean that the polypeptide is translocated across the cell membrane and presented on the surface of the bacterium in a sufficient manner for it to act as, for example, a particulate antigen. The DNA coding for the heterologous polypeptide may

then for example be fused to a S-layer coding sequence, which codes at least for a functional cell wall binding portion of a S-layer protein of *C. difficile* (Figure 3) and expressed as described above to get exposure of the heterologous polypeptide on the outside of the host cell and thus, hooked to the amidase motif of the S-layer protein. Alternatively, omitting this motif from the construct in order to get increased release of the heterologous peptide (see Specific description B3).

The heterologous polypeptide may be a foreign epitope or immunogen giving rise to antibodies that protect against disease, we note that many antibodies elicited are not protective. It typically comprises an antigenic determinant of a pathogen. The pathogen may be a virus, bacterium, fungus, yeast or parasite. The antigen may also be a "self" molecule for prevention or cure of disease (see below). The heterologous polypeptide may further be an antimicrobial peptide, e.g. for elimination of undesired microorganisms, and an anti-tumour peptide (see below) or a molecule that changes the immune response of the gut from a negative one, such as allergy or auto-immune tissue destruction, to a positive one, such as infection protection (see above). For example, *Lactobacillus* components are believed to prevent allergy development and live lactobacilli are currently given to infants in successful trials for prevention of allergy (Björkstén B, pers comm, and Kalliomäki et al. Lancet 357: 1076-1097, 2001). Cystein proteases such as cathepsin are thought to change the intestinal mucosal response to infection from a Th2 type (disease promoting) to a Th1 response (infection protection). Alternatively, the polypeptide(s) may be enzyme(s) that improve digestion of food, or that together synthesize a polysaccharide antigen of a microorganism, an antibiotic, or a specific vitamin or other nutrient or hormone useful to the host mammal. An enzyme produced by the the engineered *Clostridial* bacterium may also be an antibiotic inactivating enzyme, e.g. a beta-lactamase, to be given together with or after the antibiotic for prevention of CDAD or non-specific antibiotic induced diarrhoea, common problems in hospitals today.

The heterologous polypeptide may also be a part of an antibody molecule. This may comprise the constant part in order for example to obtain an enhanced non-specific immune response or the response to a co-administered antigen (adjuvant effect). Alternatively, it may be the variable part directed against any surface or secreted component of a microorganism (toxin, antigen, adhesin) in order to prevent its ability to colonize and cause intestinal disease. The expression product of the cassette of the invention may also represent the immune



stimulating part of allergy causing antigens lacking their IgE interacting part, thus evoking an antibody response but avoiding an allergic reaction (anti-allergy vaccination).

Whether the heterologous polypeptide is to be provided alone or fused with a carrier peptide, or presented cell-bound, released or both depends on its desired function. For example, for a polypeptide acting as an enzyme, free "secreted" molecules may be most effective, whereas in case of vaccination an antigen fused to a carrier peptide or being a cell-bound ("presented") polypeptide on a bacterium, strongly adhering to or being phagocytosed by the gut mucosa, may give the best mucosal immune response.

The immune response to a heterologous peptide may be increased by fusion to the repeating C-terminal sequences encoding the non-toxic motifs of the *C. difficile* toxins A and B that enable these to enter the colonic mucosal cells by receptor-mediated endocytosis, and/or to a portion of toxin B responsible for intracellular and intercellular spread of the antigen (see Barth et al below). Thus, by using adhering clostridial bacteria producing a desired heterologous peptide antigen fused to non-toxic parts of the *C. difficile* toxins, the mucosal immune response may be boosted (adjuvant effect).

A further improved immune response may be obtained by exploiting the natural S-layer proteins of *C. difficile* that seems to anchor the organism to the mucosa. It is likely that the amidase like fragments are directed inwards to provide cell wall anchorage, whereas the sequence unique fragments represent the the outermost portion of the S-layer protein serving as surface antigen (see above) and probably also as adhesin by which *C. difficile* attaches to the mucosal cell surface as recently suggested by Waligora et al. (Infect Imm 69:2144-2153, 2001). Thus, by switching between expression of its different S-layer ORFs over time each *C. difficile* strain may achieve surface antigen variation and thus, immune evasion and prolonged colonization in the gut. However, after deliberate colonization of a newborn with a certain *C. difficile* strain, particularly if expressing or being administered together with a "danger molecule", tolerance (see above and below) to its S-layer serotype protein may be obtained enabling the use of the same *C. difficile* serotype for efficient deliberate long-term colonization of this individual, e.g. for vaccination purposes, also during later periods of life.

In a further aspect a carrier peptide or adjuvant, e.g. a "danger molecule" is used in addition to the desired heterologous polypeptide, administered or produced *in vivo* either as a separate molecule or fused to the principal (antigenic) polypeptide. This is in order to amplify desired specific immune responses for prevention or therapy of infection, or in the neonate also for shifting its general response towards anti-infection and tolerance of "self" and

therefore away from allergy and auto-immunity (see above and below). The "danger molecule" or adjuvant is a species that may stay in a human or animal body for a long time, such as up to one, three, six months or up to one year. Alternatively, or in addition, this species is capable of eliciting a stronger immune response than the desired heterologous polypeptide acting alone. "Danger molecules" are often of microbial origin, rapidly recognized and strongly reacted upon both by the innate/primitive and the trained/specialized immune system (see above). A convenient reference for this aspect of the invention is the thesis by Carola Rask of the Department of Medical Microbiology and Immunology at Göteborg University, Sweden, ISBN 91-628-4497-0 "Cholera toxin B subunit as a carrier for inducing mucosal immunity and/or peripheral tolerance."

By using a mixture of different genetically engineered *Clostridia* strains, presenting and/or secreting different heterologous polypeptides, which act additively or synergistically, it is possible to achieve an enhanced physiological response. Thus oral immunization with a mixture of genetically engineered *Clostridia* strains, each presenting and/or secreting a different polypeptide derived from the same or another microbial pathogen, will provide both a broader immune response and a so called adjuvant effect, i.e. a more complete immune response and a better vaccination against the pathogen than obtained by using just one strain expressing a single immunogenic epitope of the pathogen.

In a further aspect of the invention we provide a medicament or therapeutic agent which comprises a *Clostridial* bacterium transformed with a gene cassette of the invention and capable of presenting on the surface of the bacterium and/or secreting a polypeptide in a human or animal body.

The medicament or therapeutic agent is conveniently a lyophilised powder for reconstitution as a suspension or for production of a solid pharmaceutical form such as a capsule or a tablet. The therapeutic agent can be administered orally or intranasally. For oral administration capsule or tablet formulations may be used. To protect the compositions against the acidity of the stomach buffer substances, e.g. sodium bicarbonate, may be used and/or the formulations may be covered with enteric coatings, e.g. Eudragite "S" or "L", cellulose acetate, cellulose phthalate or hydroxypropyl cellulose. A convenient way for oral administration of the therapeutic agents is to provide them as lyophilised powders, and shortly before administration to suspend these in for example water, fruit juice or physiological saline, optionally with addition of sodium bicarbonate or neutral citrate, or phosphate buffer to protect against the acidity of the stomach. Any convenient dose may be used, this may be in

the range from 1 to  $10^{11}$  bacteria, more conveniently we anticipate this to be in the range from about  $10^3$  to about  $10^9$  bacteria.

A principal use of the invention is in vaccination. Therefore in a further aspect we provide a vaccine which comprises a *Clostridial* bacterium transformed by a gene cassette of the invention and capable of secreting and/or presenting an antigen on the surface of the bacterium in a human or animal body.

In addition to improving the existing anti-infection vaccines and creating new ones, there is also current interest in several novel uses for vaccines.

Allergy One strategy is engineered anti-allergy vaccines containing the immunostimulatory part of each antigen but lacking the part which interacts with IgE and thus, normally elicits the allergic reaction. Another new approach is to induce an immune response towards human IgE, that normally governs the allergic response, by turning these molecules into "non-self" ones e.g. by coupling to IgE of animal origin. The use of these hybrid IgE molecules as vaccine is expected to elicit production of anti IgE antibodies that thus, inactivate human IgE thereby preventing allergy.

Alternatively, allergy may be prevented by stimulating the immune apparatus of the newborn in such a way that cellular, IgG and IgA antibody responses to microbial antigens, i.e. anti-infection, will be preferred to IgE production against allergens (immune balancing).

Auto-immune diseases. Another new proposed area for vaccines is to boost tolerance to "self" antigens *in utero* and/or in the newborn in order to prevent later development of auto-immune disorders such as type 1 diabetes, rheumatoid arthritis, inflammatory bowel disease and multiple sclerosis. This may be achieved either by non-specific tilting of the newborn immune system towards anti-infection and away from auto-immunity and allergy as mentioned above, or by applying the "self" molecule (e.g. human insulin or other beta cell antigens, connective or CNS tissue antigens) coupled to or together with a "danger molecule" of microbial origin (e.g. part of the tetanus or cholera toxin, see above) here in order to amplify the normal immunotolerance response to e.g. insulin and thus, the natural avoidance of juvenile diabetes.

Pregnancy and metabolic diseases. In contrast to the newborn, exposure to a "self" antigen especially when coupled to a "danger molecule" may in the adult individual lead to an immune response to the antigen rather than reinforced tolerance. Such vaccines boosting specific auto-immunity may be used for prophylaxis and therapy by eliciting antibodies directed against specific "self" target molecules, such as sperm or egg components or human

gonadotropin (hCG) to prevent fertility, enzymes in cholesterol biosynthesis to prevent arteriosclerosis, beta amyloid for prevention and cure of Alzheimer's disease, other brain proteins to counteract prion and Creutzfeld Jacobs disease.

Drug addiction. A further novel application of vaccines includes the use of drugs molecules such as nicotine or heroin as part of the antigen for induction of anti drug antibodies that block its activity and remove the drug and thereby abolishes its CNS effect, in order cure addiction.

Cancer. Novel multicomponent vaccines containing "danger molecules" may be of use also against cancer both by boosting the innate immune defense, by eliciting anti-tumour antibodies and cellular immune responses or by stimulating apoptosis of cancer cells.

Before the optimal anti-infection and other types of mucosal vaccines can be achieved a lot more has to be learned, e.g. about the interaction of *C. difficile* with M-cells and mucosal cells in the gut, the uptake, processing and presentation of each antigen, and optimizing the choice and presentation of adjuvant so that the immune response can be maximized and modulated in the desired way and thus, perfected for each application in order to obtain a desired cellular, IgA, IgG and subclass or combined response. Thus, the choice, size and form (soluble, particulate) of antigen, the vector (live?), the choice and form of carrier molecule (adjuvant, separate, fused to the antigen), recipient (mother, child, both, adult), timing of administration etc need to be tailored in each case.

In the above approaches to vaccination against infection in mammals and for immune modulation in newborns and adults recombinant clostridia producing desired polypeptides may be of particular interest as live vectors for mucosal immunization since certain species, e.g. *C. perfringens* and *C. difficile* belongs to the normal gut flora. *C. difficile* is particularly common in the neonatal period, and toxin negative strains thus can be given orally at all ages without ethical concerns. *C. difficile* appears to be a particularly good candidate also for delivery of antigens for gut mucosal immunization as exposure to microscopic numbers of the organism during hospital stay resulting in asymptomatic carriage is enough to yield an immune response to its toxins (NEJM 2000). Furthermore, we have observed in animals that asymptomatic gut colonization by *C. difficile* results in an immune response also to its S-layer protein (see below). These responses are probably enhanced by the the non-toxic part of the *C. difficile* toxins that are used for their receptor mediated pinocytosis into the mucosal cells. Toxin B then can form membrane pores in the pinocytic vacuoles containing toxin and presumably also in phagocytic vacuoles containing whole bacteria (Barth H et al, Low pH

induced formation of ion channels by *C. difficile* toxin B in target cells. J Biol Chem 276(14): 10670-10676, 2001). Thereby the toxin and other bacterial components may be released into the cytosol of the mucosal cells and may spread also to neighbouring cells including to antigen presenting cells and thus, enhancing an immune response. Such an unusual adjuvant effect of *C. difficile* toxin B obtained by breakage of phagocytic vacuoles and intercellular spread of internalized antigens and bacteria can alternatively also be obtained e.g. by including the membrane attacking peptide listeriolysin O from *Listeria monocytogenes* in a recombinant *C. difficile* strain in order to boost immunity as has been shown in experiments using other gut mucosal delivery systems (Dietrich G et al, From evil to good: a cytolysin in vaccine development, Trends in Microbiology 9:23-28, 2001).

Clostridia furthermore represent a unique torpedo able to deliver a desired heterologous polypeptide to hypoxic tissues such as tumours. This is because spores of these obligate anaerobic organisms given intravenously are known to settle and be able to germinate into growing clostridial cells in the hypoxic parts of tumours but not in healthy tissues. This phenomenon was described already in 1955 (reference 7 in Theys J et al, FEMS Immunol Med Microbiol 30:37-41, 2001) and is currently being studied for therapeutic applications by several groups (see also Abstracts, The 3<sup>rd</sup> Int Meeting on Molecular Genetics and Pathogenesis of the Clostridia, Chiba, Japan, June 8-11, 2000). Thus, anti-tumour peptides including apoptosis inducing peptides, cytokines, toxins and other proteins, such as enzymes locally converting pro-drugs to active anti-cancer chemotherapeutic agents, thus minimizing systemic side effects, all produced by recombinant clostridia inside tumours may become novel approaches to cure cancer. We propose that another approach to anti-tumour therapy is iv injection of spores of recombinant clostridia producing various angiogenesis inhibitors, mostly of peptide nature and currently used in many clinical trials (phase I: angiostatin, SU6688, combrestatin A-4 prodrug, PTK787/ZK2284; phase II: endostatin, anti-VEGF Ab, TNF-470, IL-12, 2-methoxyestradiol, squalamine, vitaxin, EMD 121974, COL-3, CGS-27023A, CAI; phase III: thalidomide, marimastat, INF-alfa, neovastat, BMS-275291, SU5416, AG3340, IM862 as summarized in Larsson H, Regulation of angiogenesis, Thesis, 2001, Uppsala University, Sweden, ISBN-91-554-4954-9).

A problem in these studies and possible to overcome by local production of enzymes, toxins and other peptides is limited effect *in vivo* due to short half-life in serum and thus, insufficient local concentrations of e.g. the cytostatic agent or the anti-angiogenesis peptide. In the latter application recombinant clostridia may become particularly powerful (self-

accelerating), because the effect of the peptide will further lower the oxygen tension and thus, enhance bacterial growth and further production of the peptide(s)

We further believe, that that iv administration of spores from recombinant clostridia may be used also against other diseases involving local tissue hypoxia such as fibrinolytic and other agents for venous or arterial occlusion, and oxygen releasing or other tissue vitalizing molecules for tissue necrosis and gangrene.

We now found and disclose that *Clostridial* spores may be used to deliver heterologous polypeptide(s) to a human or animal body. This is an important step forward. A spore is a dormant or resting state of a bacterial cell. Unlike bacterial spores from species belonging to the obligate aerobic genus *Bacillus* (see above), ingested *Clostridial* spores naturally germinate into vegetative bacteria that can grow anaerobically and naturally colonise a human or animal gut. Intake of the spores of the genetically engineered *Clostridia* is preferably through the oral route. Spores are able to resist stomach HCl and digestive enzymes. Upon contact with bile they will germinate and establish themselves in the colonic flora as vegetative bacteria presenting and/or secreting for example the desired heterologous peptide *in vivo*.

Therefore, in a further aspect of the invention we provide a therapeutic agent which comprises spores of *Clostridia* transformed with a construct capable of expressing, secreting or presenting a heterologous polypeptide in the mammalian body after conversion (germination) to live (vegetative) bacteria.

The construct is preferably a recombinant gene cassette of the invention as outlined before. The mammalian body is preferably a human or animal.

The use of *Clostridial* spores has a number of advantages including low production cost, relative ease of production, very long shelf life independently of the mode of storage, ease of administration, production of antigen at the site of action, and an oral route of immunisation which may be superior to a parenteral one. As mentioned above for bacteria, spores are suitable for administration of mixtures (cocktails) of recombinant *Clostridia* having different desired properties.

The use of live vaccines administered via the oral route may lead to further fecal-oral transmission and enhanced immunization of a population. On the other hand, this may also be considered as to be unwanted spread of genetically modified organisms in the environment. Spores of *Clostridia* survive readily in the environment whereas the vegetative forms have a very limited capability to survive in an oxygen-containing milieu. The invention may be

further developed to create *Clostridia* that are unable to reconvert to spores, once they have germinated in the colon. One way is to modify a genetic element present in *C. difficile* that is similar to the so-called *skin* (Sigma K intervening) element of *B. subtilis*. This element truncates the sigma K factor necessary for sporulation, and becomes removed by a specific excision system during sporulation (Krogh, S. et al.(1996) and Takemaru, K. et al.(1995)). By genetic modification of the excision system, and insertion of wild-type copies with inducible promoters, there is a possibility to create a host strain that is able to sporulate only during special conditions, e.g. in the presence of a special chemical (IPTG) or at low temperature (20 °C). Such construction would allow the production of spores *in vitro*, whereas no new spores are created in the vaccinated host. The spread of genetically modified *Clostridic* microorganisms to the environment would still occur, but the probability of survival of these organisms would in practice be very low or nil.

Spores of the transformed *Clostridia* are produced, purified and isolated in the same way as for native *Clostridial* strains. They may thus be readily obtained from a stationary phase culture for example by treatment with ethanol, acid or heat or by combinations of such measures followed by purification and isolation in a conventional way. As outgrown spores will have the same properties as the parental bacteria, purification of the spores may not even be necessary.

Pharmaceutical and veterinary compositions for oral administration, comprising spores of transformed *Clostridia* and pharmaceutically acceptable carriers, diluents and excipients are further provided by the invention. They have the ability to colonise the intestinal tract of humans and animals with live *Clostridial* bacteria producing and presenting or secreting the heterologous polypeptide coded for by the modified gene cassette provided by the invention or by any other construct. The pharmaceutical and veterinary compositions may comprise tablets, capsules, powder for reconstitution or any other form suitable for oral administration to humans or animals. Examples of pharmaceutically acceptable carriers and diluents are lactose and carboxymethyl cellulose. A convenient way of oral administration of these therapeutic agents is to provide them as lyophilized, or just dried, powders; shortly before administration they are suspended in for example water, physiological saline or fruit juice. The dose is as indicated above for *Clostridial* bacteria.

In a further aspect of the invention we provide a method of treatment for the human or animal body, which comprises of administering a therapeutic agent comprising *Clostridial* spores capable of expressing a heterologous polypeptide in a human or animal body.

As in the treatment with *Clostridial* bacteria described above it will be possible to treat with mixtures of transformed *Clostridial* spores to obtain in the body a mixture of different heterologous polypeptides, which may act synergistically or additively.

It will be appreciated that the therapy may be either prophylactic or therapeutic.

5       The method may be applied to any convenient mammal such as a human or animal. Convenient animals include domestic animals such as dogs and cats, also cattle, pigs, chicken and horses.

10       In a further aspect of the present invention we provide a method for immunisation which method comprises administering to a mammalian body *Clostridial* spores capable of expressing a heterologous antigen after germination.

15       Examples of convenient *Clostridium* spores include the spores of *C. difficile* and *C. perfringens*, which normally colonise the large intestine of man. For animals other *Clostridia* such as *C. tetani* may also be useful. Thus, the intensity and duration of antigen exposure in the gut (clostridial colonization) in a particular host can be varied by not only exploiting and manipulating e.g. adherence of *C. difficile* (see above), but also by selecting the appropriate *Clostridium* species with regard to the intended host mammal.

In a further aspect of the invention we provide *Clostridial* spores transformed with a gene expression cassette of the invention.

20       It will be appreciated that the methods and materials of the invention may also be used for other applications such as the display of antibodies and peptide libraries. They may also be used for screening proteins and antigens and also to provide a support for immobilising an enzyme, peptide and/or antigen.

**Sequence numbers of proteins encoded by the ORFs of the invention**

<u>Protein encoded by ORF</u>	<u>SEQ ID NO:</u>
ORF 1	SEQ ID NO: 22
ORF 3	SEQ ID NO: 23
ORF 5	SEQ ID NO: 24
ORF 6	SEQ ID NO: 25
ORF 7	SEQ ID NO: 26
ORF 9	SEQ ID NO: 27
ORF 11	SEQ ID NO: 28



**Protein encoded by ORF    SEQ ID NO:**

ORF D	SEQ ID NO: 29
ORF E	SEQ ID NO: 30
ORF G	SEQ ID NO: 31
ORF H	SEQ ID NO: 32
ORF I	SEQ ID NO: 33

The present invention is particularly directed to a gene expression cassette comprising a secretory leader sequence encoding a signal peptide from *Clostridium difficile* having an amino acid sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7 and signal peptides of analogous exported clostridial N-acetylmuramoyl-L-alanine amidase-like proteins, linked to a DNA sequence encoding a heterologous polypeptide. The signal peptides of the analogous clostridial N-acetylmuramoyl-L-alanine amidase-like proteins may also be selected from *Clostridium difficile* signal peptides having an amino acid sequence of any one of SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, and SEQ ID NO: 12.

The gene expression cassette may further include a promoter of prokaryotic origin, e.g. selected from clostridial promoters comprising a nucleotide sequence selected from the group consisting of SEQ ID NO: 13 – 21, or from the promoters of ORFs 1-11 or D-I mentioned above.

The gene expression cassette according to the invention may further include a DNA sequence encoding at least a cell wall binding portion of a protein of prokaryotic origin functioning in clostridia such that a fusion polypeptide may be presented on the outer surface of a host cell harbouring the cassette.

The gene expression cassette according to the invention may in particular include a DNA sequence encoding at least a functional cell wall binding portion of an S-layer protein of *C. difficile* selected from any one of the polypeptides having an amino acid sequence selected from SEQ ID NO: 22 - 33 such that a fusion polypeptide may also be presented on the outer surface of a host cell harbouring the cassette. The DNA encoding the cell wall binding portions of SEQ ID NO: 22-33 may be omitted such that the fusion peptide is secreted into the surrounding milieu by the host cell harbouring the cassette.

Further, the gene expression cassette according to the invention may be such that the DNA sequence encoding the heterologous peptide is inserted at a point downstream the first (signal) proteolytic cleavage sites in the gene encoding a polypeptide having an amino acid sequence selected from SEQ ID NO: 22 – 33, optionally including or excluding its second  
5 cleavage site.

In addition, the gene expression cassette according to the invention may further comprise at least a functional part of a secretory (*secA*) gene recognizing the signal peptide, to allow translocation of a heterologous polypeptide and/or fusion polypeptide across the cytoplasmic membrane of a host cell harbouring the expression cassette. For example, the  
10 secretory gene may be from *C. difficile* and encode a polypeptide having the amino acid sequence SEQ ID NO: 34.

In a preferred embodiment of the invention the gene expression cassette is the one that is shown in Figure 3.

The invention is also directed to a vector comprising a gene expression cassette  
15 according to the invention, such as a plasmid.

The invention is further directed to host organism transformed with a vector according to the invention for expression of the heterologous polypeptide and/or fusion polypeptide.

In an embodiment of the invention the host organism is a *Clostridium* host organism transformed with a vector according to the invention for expression of the heterologous  
20 polypeptide and/or fusion polypeptide.

In a preferred embodiment, the host organism is *C. difficile* or *C. perfringens*.

Further, the invention is directed to a pharmaceutical or veterinary composition or formulation which comprises *Clostridial* cells transformed with a vector according to the invention, with the ability to present on the cell surface and/or to secrete an expressed  
25 heterologous polypeptide or fusion polypeptide, together with a pharmaceutically or veterinary acceptable carrier or diluent. Preferably, the composition or formulation is suitable for oral or intranasal administration. The composition or formulation according to the invention may further comprise, as adjuvants, non-toxic motifs of the *C. difficile* toxins A and/or B that enable the heterologous polypeptide and/or fusion polypeptide to enter the  
30 colonic mucosal cells of a mammal by receptor-mediated endocytosis, and/or a portion of toxin B responsible for its intracellular and intercellular spread. The composition or formulation according to the invention may alternatively additionally comprise a further fused or separate carrier peptide or adjuvant, in addition to the expressed heterologous polypeptide

and/or fusion polypeptide, to elicit a stronger or differently directed immune response than that against the expressed heterologous polypeptide acting alone.

The invention is, in another aspect, directed to a vaccine which comprises a *Clostridial* bacterium transformed with a vector according to the invention and capable of presenting on the surface of the bacterium and/or secreting an antigen in a human or animal body, and optionally an adjuvant described in conjunction with a composition or formulation of the invention. The vaccine may comprise a mixture of at least two differently engineered *Clostridia* strains, each capable of presenting on the surface of the bacteria and/or secreting a different heterologous polypeptide and/or fusion polypeptide. Further, the vaccine may comprise spores of *Clostridia* cells or bacteria transformed with a vector according to the invention and capable of germinating into cells which are able to grow, express, and present or secrete a heterologous polypeptide and/or fusion polypeptide, and optionally also an adjuvant described in conjunction with a composition or formulation of the invention, in a mammalian body. The vaccine may comprise a mixture of spores from at least two differently engineered *Clostridia* strains. Each of these strains is capable of presenting on the surface of the bacterium and/or secreting a different heterologous polypeptide and/or fusion polypeptide. The spores are preferably from *C. difficile* or *C. perfringens*.

The invention is in yet another aspect directed to a medicament which comprises a *Clostridial* bacterium transformed with a vector according to the invention and capable of presenting on the surface of the bacterium and/or secreting a heterologous polypeptide and/or fusion polypeptide in a human or animal body, and optionally an adjuvant described in conjunction with a composition or formulation of the invention. The medicament may comprise a mixture of at least two differently engineered *Clostridia* strains, each capable of presenting on the surface of the bacteria and/or secreting a different heterologous polypeptide and/or fusion polypeptide. Further, the medicament may comprise spores of *Clostridia* cells or bacteria transformed with a vector according to the invention and capable of germinating into cells which are able to grow, express, and present or secrete a heterologous polypeptide and/or fusion polypeptide, and optionally an adjuvant described in conjunction with a composition or formulation of the invention, in a mammalian body. The medicament may comprise a mixture of spores from at least two differently engineered *Clostridia* strains. Each of these strains is capable of presenting on the surface of the bacterium and/or secreting a different heterologous polypeptide and/or fusion polypeptide. The spores are preferably from *C. difficile* or *C. perfringens*.

The invention is in still another aspect directed to a method for vaccination of a mammal, which comprises administering a therapeutically or prophylactically effective dose of a vaccine according to the invention to the mammal. Spores used in the vaccine are preferably from *C. difficile* or *C. perfringens*.

5 The invention is also directed to a method for prophylactic or therapeutic treatment of a mammal, which comprises administering a therapeutically or prophylactically effective dose of a medicament according to the invention to the mammal. Spores used in the medicament are preferably from *C. difficile* or *C. perfringens*.

The invention is additionally directed to a *C. difficile*-associated diarrhea (CDAD) vaccine comprising spores according to the invention and capable of expressing, after  
10 germination,

- (i) relevant parts of the *C. difficile* toxins, alone or together with a
- (ii) suitable adjuvant to provide an IgA response to the toxin antigenic epitopes and
- (iii) S-layer protein antigenic variants (serotype antigens) or fimbrial antigens to obtain,  
15 after administration to a mammal, a polyvalent anti-S-layer (or anti-fimbrial) IgA response preventing *C. difficile* colonization of the mammal.

The invention will now be illustrated but not limited by reference to the following Figures, Specific Descriptions, Tables, and Examples wherein:

**Figure 1A** shows our first simple layout of the *C. difficile* strain 630 genomic segment encoding the S-layer genes. ORF2 represents *secA* and ORFs 1,3, 7-9 and 11 S-layer protein genes. For explanations of these and other ORFs, see Table 1 and Example 9.

**Figure 1B** represents the result of comparisons between three of the S-layer ORFs with published sequences of other genes. The "amidase enhanced precursor" sequence is equivalent to the N-acetyl muramoyl L-alanine amidase motif mentioned in the text.

**Figure 2.** Defining the upstream region of ORF 1-12. The figure illustrates additional information and genetic organisation of the *C. difficile* S-layer genes (cf. Figure 1), found after searches in the revised *C. difficile* database at the Sanger Centre. The genes upstream of ORF 1 to 12 are denoted A to I (see also Table 1). The numbers +1, +2 and +3 indicate the reading frame of the ORFs relative to the start point of the contig. ORFs D, E, G, H and I had  
25 the amidase motif typical of genes encoding the *C. difficile* S-layer proteins.

**Figure 3** shows an example of a preferred gene expression cassette here taken from *C. difficile* strain 630 and containing a strong promoter, the secretory leader peptide from ORF1, the signal peptide cleavage site the area of insertion of foreign DNA encoding the  
30

heterologous peptide, the second (optional) peptide cleavage site in the N-acetyl muramoyl L-alanine amidase motif, and the *secA* gene (ORF2).

Figure 4 shows a preferred strategy for introducing a recombinant gene cassette of the invention back into *C. difficile* via *B. subtilis*.

- 5 Figure 5. Further details of a particular *C. difficile* S - layer gene cassette. This is a 4960 bp cassette taken from strain 630 encoding an S-layer protein of 2160 bp in its original form (ORF1). The 210 bp region (pr, promoter) upstream of ORF1 includes gene control elements for the S-layer protein included in the cassette. Also shown are an intervening 244 bp region and the 2346 bp *secA* sequence.
- 10 Figure 6. Strategy for the engineering of ORF1 (Figure 5) to express a recombinant protein (for example as outlined in Example 2 and 2A). The 613 bp variable region (vr) is replaced by a foreign DNA. For example, three fragments encoding the Hepatitis B virus surface antigen (HBsAg) were selected: (i) the full length HBsAg that includes the pre S1, pre S2 and the S gene (1207 bp); (ii) the S gene (740 bp); and (iii) the subtype from the S gene (minimum
- 15 antigenic epitope, 421 bp).
- Figure 7. Cloning strategy for the construction of ORF1 - *secA* (with and without the native promoter) together with the different lengths of the HBsAg antigenic loop (full length, S gene and Sub type - see legend to Fig. 6 above) using a PCR based method and cloning into the TA vector in *E. coli*. The primers indicated were used also for PCRs to help checking the
- 20 correctness of the constructs. The expected and obtained constructs were 5564, 5097 and 4778 bp respectively.

### Specific descriptions

- 25 A. Characterisation of the genomic segment responsible for S-layer protein expression in *C. difficile* (outlined in Fig. 1 and Fig. 2, see also Table 1 and Example 9):
1. The main surface layer proteins expressed by *C. difficile* strain 630 has been found to encoded by a single open reading frame (ORF1) encoding a 72 kDa protein. The gene product of ORF1 is postrationally cleaved at two sites yielding three different
- 30 peptides; the leader peptide and the final S-layer proteins of apparent molecular weights of 36 kDa and 45 kDa.

2. The C-terminus of ORF1 shows similarity to N-acetyl muramoyl L-alanine amidase, and the N-terminus shows weak similarity to surface layer proteins from *L. helveticus* (Fig. 1B).
  3. The gene immediately downstream of ORF1 (ORF2) encodes the SecA protein responsible for secretion of proteins with signal peptides.
  4. The genes downstream of *secA* encodes proteins with similarities to ORF1 (ORF3, 5-7, 9 and 11, see Fig. 1).
  5. ORF1 is efficiently expressed and its product is efficiently exported in strain 630, whereas e.g. ORF3 is expressed more than 100-fold less in strain VPI 10643, indicating a strong termination between ORF1 and 3 (as judged by identification of exported proteins by two-dimensional gel electrophoresis).
  6. The upstream region including the putative promoter has not been characterized functionally, but the very high expression of ORF1 in various growth conditions indicates the action of a strong, constitutive promoter. It is also active in *E. coli* (see Example 2).
  7. In the revised *C. difficile* sequence database from Sanger Centre, the upstream region of ORF 1 was included and revealed 9 new ORFs (A-I) of which 5 (D, E, G, H and I) had the N-acetylmuramoyl L-alanine amidase motif typical of the *C. difficile* S-layer protein ORFs (Fig. 2 and Table 1). The putative promoter region for ORF1 is thus situated between ORFI and ORF (See Example 1A and Table 2).
- The S-layer proteins from strain VPI 10463 have similar molecular weights but different pI as compared to those of strain 630, and the N-terminal sequences of the two S-layer proteins from VPI 10463 showed no similarity with those of strain 630. Studies of strains from different serogroups showed that the S-layer proteins vary in pI and molecular weight. The downstream region of the gene segment may in part be more conserved, since the N-terminal sequence from another extracellular protein from strain VPI 10643 was identical to ORF3 of strain 630. Our results indicate that ORF1 is located at part of the chromosome that is capable of expressing and exporting various S-layer proteins depending on the strain.

**B. Designing a preferred engineered gene cassette based on the *C. difficile* S-layer gene segment data (outlined in Fig. 3).**

1. Any strong prokaryotic promoter functional in *Clostridia* can be used to express the heterologous peptide, e.g. the promoter of ORF1 or any of the promoters of genes encoding

other highly expressed proteins in *C. difficile* such as certain electron transfer proteins (our unpublished data and Fix A and Fix B in Example 9) or ribosomal proteins.

2. A secretory leader peptide, preferably the leader peptide from ORF1, is fused with the heterologous peptide, to ensure its translocation across the cell membrane

3. Depending on the desired fate of e.g. the antigen (secreted, surface presented or both), the heterologous peptide is optionally fused to the amidase part of the S-layer protein optionally including the part involved in the proteolytic cleavage event (Figure 3). Thus, for maximum release of the peptide the secretory leader of e.g. ORF1 may be sufficient. On the other hand, maximum cell-wall binding may require fusion to the amidase portion but omitting the proteolytic cleavage sequence in the middle of the gene (Figure 3). If both free and bound heterologous peptide is desired one recombinant cassette of each type present in the same *Clostridium* strain or a mix of two different strains, each harbouring one the recombinant cassettes, can be used. The peptide cleavage site may be exploited if for instance the antigen and an adjuvant are produced in a fused form, to obtain equal amounts of the two, but are desired as separate peptides on the outside of the producer bacterium. To what extent parts of the N-terminal (variable) portion of e.g. ORF1 can be used to optimize the localization of the heterologous peptide requires further experimentation:

4. The *secA* gene is usually included in the construct to ensure efficient translocation of the polypeptide across the cell membrane.

5. The gene construct is made in plasmids suitable for transformation of both *E.coli* and *C. perfringens* (e.g. pJIR750 or 751) or in plasmids suitable for conjugation into *C. difficile* via *B. subtilis* (e.g. pCI195 or pSMB47, Figure 4).

## Examples

### **Example 1**

Defining a minimal gene segment responsible for production and export of the heterologous protein (an engineered gene cassette, see also Figure 3).

#### **A. Defining the promoter start (the 5' end).**

The promoter region may be further characterised in different *C.difficile* strains, for example by the following steps:

1. Defining the DNA upstream of ORF1 by circular PCR. One or several restriction enzymes are used to cut the DNA outside ORF1. Useful enzymes are NdeI, TaqI, PstI, and BsmAI.

A Southern blot is performed using a probe against ORF1 to confirm size of the fragment (optimal size is around 1-3 kb). The cleaved genomic DNA is then ligated into circles. PCR is then performed using primers (directed outwards of each other) directed against ORF1. The PCR product is then cloned and sequenced. Optionally, PCR is performed with one primer directed against ORF1 (lower primer) and different arbitrarily designed primers. The products are then cloned and sequenced.

2. Identification of the transcription start. Primer extension with primer located at the 5 prime end of the genomic segment is used.
3. By computerised search for homologies to known promoters. The new sequence data for strain 630 enabled a search for putative promoters upstream of ORF1 and the result is shown in Table 2. Due to the AT rich genome, several putative promoters were found. The actual promoter start has to be experimentally determined as described in point 2 above.

#### **B. Defining the termination point (the 3' end).**

Primers directed at different parts of ORF1, 2 and 3 to determine transcript abundance using Northern blots. Termination loops in the RNA may be identified by computer analysis.

**C. Conservation of the S-layer locus in other *C. difficile* strains: characterization of part of the S-layer genomic segment (ORF1 and proximal parts of its upstream region and of part of ORF2) in different *C. difficile* serogroups in order to check for conservation of this region of the genome.**

Primer pairs that are directed against the identified upstream-transcription start region and the proximal part of ORF2 were designed. PCR was performed on different strains belonging to all serogroups to confirm the generality of the expression center of the S-layer locus.

(i) PCR performed with chromosomal DNA as template from the different serogroups of *C. difficile*.

The primers used:

CONS1: 5 prime- TAT AAT GTT GGG AGG AAT TTA AGA – 3 prime, total length 24 nt

(5 prime end starts at 8th nt upstream of ORF1, ends at 32nd nt)

CONS2: 5 prime- CAA ATC CAA ATT CAC TAT TTG TAC – 3 prime, total length 24 nt

(5 prime end starts at 2983rd nt downstream of ORF1, ends at 2959th nt)



Total size of expected PCR pdt (from strain 630 sequence): 2975 bp (includes ORF1 and the proximal part of ORF2).

The enzyme/system used: Expand™ Long Template PCR System from Boehringer Mannheim.

- 5 Reaction conditions (as specified by the manufacturer): In a total reaction volume of 50ml, 350mM dNTPs, 300nM primers, 50ng chromosomal DNA template, 1x supplied PCR buffer with 1.75mM MgCl<sub>2</sub>, and 2.5U of a mix of Taq and Pwo DNA polymerase. 10ml of the reaction mix was run on a 0.8% Agarose-TBE (Tris-Borate- EDTA buffer) to check for product.

- 10 PCR cycle conditions: Initial denaturation - 92°C for 2 mins

For 30 cycles - denaturation - 92°C for 10 secs

annealing - 40°C for 30 secs

elongation - 68°C for 2 mins

Final elongation - 68°C for 2 mins

15

PCR results:

	<u>Serogroup tested</u>	<u>Product size</u>	<u>No. of Expts</u>
	A	none	3
	B	around 2900bp	2
20	C	around 2900bp	2
	D	around 2900bp(lesser amt)	2
	F	around 2900bp	2
	G	around 2900bp	2
	H	pdt > 2900bp (lesser amt)	3
25	I	around 2900bp	2
	K	around 2900bp	2
	X	around 2900bp	2
	A2	around 2900bp (lesser amt)	3

	<u>Serogroup tested</u>	<u>Product size</u>	<u>No. of Expts</u>
	A3	around 2900bp (lesser amt)	2
	A4	around 2900bp (lesser amt)	2
5	A5	none	2
	A6	around 2900bp (lesser amt)	2
	A8	around 2900bp	2
	A9	around 2900bp (lesser amt)	2
	A10	around 2900bp	2
10	S1	around 2900bp	2
	S3	around 2900bp	2
	S4	none	2
	VPI 10643 (G)	around 2900bp	4, last PCR was weak
	630 (X)	around 2900bp (expected size)	5, last PCR was -ve
15	Serogroups A, A5 and S4 did not give any PCR product with several attempts. The PCR reactions were very sensitive to template condition, which had to be prepared fresh.		

(ii) Restriction Enzyme Analysis of the PCR products.

12ul PCR reaction mix (reactions done above) from most of the serogroups (except A, A5-6, A9, and S4) were further subjected to RsaI (Boehringer Mannheim) and Sau3A1 (Amersham Pharmacia Biotech) digestions. Total digestion reaction mixes were run on a 0.8% Agarose-TBE (Tris-Borate- EDTA buffer) to obtain respective digestion patterns.

Each serogroup tested appears to give rise to unique restriction pattern though the apparent size of the PCR product appears to be similar except for gp H.

25 Conclusion:

It appears that the promoter-ORF1-ORF2 organization and the size of the genetic segment between the putative promoter and ORF2 (*secA* gene), i.e. the ORF 1 equivalent, is generally conserved in *C. difficile*. There may be sequence variation between ORF 1 equivalents from strains of different serotype and also between two strains of the same serotype, presumably reflecting the variable (non-amidase) portion of these genes.

### Example 2

Cloning of the *C. difficile* promoter-ORF1-ORF2 “cassette” from strain 630 and construction of a recombinant cassette in *E. coli* encoding heterologous proteins to be transferred to *C. perfringens* and *C. difficile* and used for immunisation.

1. Primer pairs are designed that include the promoter region and part of ORF1 including the leader peptide sequence (Figure 5). PCR is performed followed by cloning of the product into *E. coli*-*C. perfringens* shuttle vectors pJIR750 or pJIR751 (Plasmid 229: 233-235, 1993) in frame of a reporter gene such as  $\beta$ -lactamase or at least a part of the hepatitis B virus (HBV) antigen. A convenient source of HBV antigen is the SMI strain no. 8423/87 having the genotype A and subtype adw2 (cf. Magnus et al, J.Gen. Virology, 1993, 74, 1341-1348). The plasmid is isolated from *E. coli*, purified and used to transform *C. perfringens*, and the engineered strain is isolated for further use. The *secA* gene is optionally included in the construction to optimise secretion.
2. Check for expression and secretion of the reporter gene using an HBV antigen based assay or a  $\beta$ -lactamase assay of the transformed *E.coli* and *C. perfringens* strain. Gnotobiotic mice and rats are fed with spores of the engineered *Clostridium* strain to obtain colonisation. Expression of antigen is checked by the HBV antigen based assay or  $\beta$ -lactamase assay in feces and immune response by antibody response in feces and in serum.

**A. Cloning of ORF1 - ORF2 (*secA*) in order to construct a fusion with a foreign antigen**  
 PCR was performed with chromosomal DNA from Strain 630 as template. The primers used:

AMP1: 5' - GGA ATT CCA TGA ATA AGA AAA ATA TAG CA - 3',

total length 29 nt (5' end starts at the first codon of ORF1, ends at 7th codon )

AMP2: 5' - CGG GAT CCC GTT TTT AGT TAA ATT TAT ATA AG - 3',

total length 32 nt (5' end starts at the stop codon for *secA*)

Analogous PCRs but with the first primer in the upstream region in order to include a putative native promoter were also performed.

Total size of expected PCR product (from strain 630 sequence) : 4770 bp (4960 bp including the promoter) (Figure 5).

The enzyme/system used: Expand™ Long Template PCR System from Boehringer Mannheim. Reaction conditions (as specified by the manufacturer): In a total reaction volume of 50ul, 350mM dNTPs, 300nM primers, 150ng chromosomal DNA template from strain 630, 1 x supplied PCR buffer with 1.75mM MgCl<sub>2</sub>, and 2.5U of a mix of Taq and Pwo DNA polymerase. 10ul of the reaction mix was run on a 0.8% Agarose-TBE (Tris-Borate- EDTA buffer) to check for product.

PCR cycle conditions: Initial denaturation - 92°C for 2 mins

For 30 cycles - denaturation - 92°C for 10 secs

annealing - 40°C for 30 secs

elongation - 68°C for 4 mins

final elongation - 68°C for 5 mins

#### Results:

The expected PCR products were obtained and cloned into pGEMT vector (Promega). The plasmid containing the insert will be subjected to partial digestion with PvuII enzyme (sites at position 282 and 895 of the insert) to eliminate the 613 bp internal fragment from ORF1, where the foreign antigen is planned to be inserted (Figure 6). The digestion time had to be standardised. The foreign antigen used was the hepatitis B virus (HBV) surface antigen (HbsAg).

Three HBsAg regions were used (Figure 6): (a) Total (413 aa, 3-1243 bp; 1240 bp)

(b) Central (259 aa, 465-1243 bp; 778 bp)

(c) C-terminal (140 aa, 822-1243 bp; 421 bp)

References for HBV antigen sequences include Prange et al, 1995, 14 (2), 247-256 and Chen et al, 1996, 93, 1997-2001.

Alternative antigens that may be used include relevant epitopes of the rota virus and hepatitis A virus.

#### **B. Cloning of the *C. difficile* ORF1-secA cassette containing the three HBsAg DNAs in *E. coli*.**

##### Result:

The cloning strategy (Figure 7) using PCR and the TA vector was successful according to DNA analyses including agarose gel electrophoresis and PCR.

**C. ELISA to check for expression in *E. coli* of the three ORF1-HBsAg-*secA* DNA constructs in the TA vector.**

A commercially available ELISA (Abbott) was performed on sonicated samples of three overnight cultures of *E. coli* each containing one of the three different HBsAg DNAs inserted into ORF1-*secA* (with the native *C.difficile* ORF1 promoter) and cloned into the TA vector.

**Results:**

Cut off value = 1.0, higher values are regarded as positive.

<i>Sample (plasmid)</i>	<i>Culture OD*</i>	<i>Sonicate-supernatant (periplasm +cytoplasm)</i>	<i>Pellet following sonication (cell membrane)</i>	<i>Spent medium</i>
1.3	1.1 <sup>#</sup>	0.92	Test Failed	0.61
1.3 <sup>a</sup>	1.9	4.76	3.70	0.69
2.8	1.6 <sup>#</sup>	0.99	1.33	0.74
2.8 <sup>a</sup>	1.8	1.14	4.33	0.69
3.7	1.4 <sup>#</sup>	18.56	34.58	0.70
3.7 <sup>a</sup>	2.0	5.10	5.11	0.69
TA vector	2.0 <sup>#</sup>	0.70	1.13	0.69
TA vector <sup>a</sup>	2.0	0.74	0.91	0.70
PBS(buffer)	NT	0.77	NT	NT

10 1.3 : Full length HBsAg(pre S1, pre S2 and S gene),

2.8 : S gene,

3.7 : Partial S gene(the minimum HBsAg epitope)

a : Duplicates

# : Samples stored overnight in cold before analysis.

15 NT : Not tested

\* : Culture OD at the time of harvest. Cells from 1ml culture were pelleted and resuspended in 0.5ml of PBS before sonication.

The above experiment was repeated with cells from 5ml cultures resuspended into 1ml PBS to see if increased protein concentrations would give higher titer values. Here a control *E. coli*

culture with the TA vector carrying the ORF1-*secA* cassette but without any HBsAg insert was also included.

<i>Sample</i>	<i>Culture OD</i>	<i>Sonicate-supernatant (cytoplasm)</i>	<i>Pellet following sonication (cell membrane)</i>
1.3	2.4	8.17	3.08(1/10 dil)
2.8	1.8	5.07	2.02(1/10 dil)
3.7	2.4	42.0	1.15(1/100 dil)
orf1secA (no HBsAg insert)	2.4	Test Failed	0.92

##### 5 Conclusion:

All three constructs containing HBV DNA expressed both soluble and and particulate HBsAg. Thus, the cloning proved to be successful and the native *C. difficile* ORF1 promoter was to some extent functional also in *E. coli*.

##### 10 Example 3

Cloning of the ORF1-ORF2 (*secA*) "cassette" into *C. difficile* and construction of recombinant protein useful for immunisation studies.

1. Cloning of a gene expression cassette according to examples 1 and 2 in plasmids pCI195 or pSMB47 followed by transfer to a non-toxigenic strain of *C. difficile* by mating via *B. subtilis* is performed according to standard methods reported (J. Gen. Microbiol 136: 1343-1349, 1990; Plasmid 31: 320-323, 1994, see also Figure 4). The engineered strain is isolated for further use.
2. Check for expression, secretion and antibody response *in vitro* and *in vivo* (see Example 2).

##### 20 A. Cloning of the ORF1-HBsAg-*secA* constructs into the shuttle plasmid vector pJIR750

Results:

Ligation mixtures containing the desired recombinant plasmids were obtained and judged by agarose gel electrophoresis and PCR. However, upon transformation into *E. coli* the plasmid constructs were fragmented. This indicated that a plasmid replication machinery better at handling large plasmids in *E. coli* than that of pJIR750 (colE based) needs to be used. Also,

attempts to transform *C. perfringens* with our recombinant pJIR750 plasmids are being performed.

#### Example 4

5 Production of transformed *C. perfringens* expressing and presenting foreign antigen for vaccination.

##### A. Live bacteria.

10 *C. perfringens*, transformed with a gene cassette coding for a foreign antigen fused to ORF1 and obtained as in Example 2 is cultivated under anaerobic conditions in a fermenter until a cell density of at least  $10^7$  bacteria per ml is obtained. The broth is cooled to 11°C, the bacteria recovered by centrifugation and the supernatant discarded. The pellet is twice washed with cold 0.1 M phosphate buffer, pH 7 and centrifuged. The final pellet is resuspended in the phosphate buffer to a concentration of about  $10^9$  organisms per ml. One ml portions of the  
15 suspension are dispensed into glass ampoules and freeze-dried to remove the water. The final product is obtained by sealing of the ampoules *in vacuo*.

##### B. Bacterial envelopes.

20 Transformed *C. perfringens* bacteria are produced as in A above. The final pellet is suspended in 50 mM Tris-HCl, pH 7.2, and sonicated for 1-10 min (40 watt, Bransic Sonic Power co. Sonicator). Triton X-100 is added to a final concentration of 2% and the mixture incubated under stirring at 11°C for 30 min.. The cells are collected by centrifugation and washed three times with cold distilled water. The pellet is resuspended in 5 mM MgCl, containing DNase (11mg/ml) and RNase (11mg/ml) and incubated for 15 min at 11°C. The resulting envelopes  
25 are recovered by centrifugation, washed three times with cold distilled water, resuspended in cold distilled water and freeze-dried to give the envelopes as a powder suitable for formulation in capsules or tablets, for suspension in e.g. physiological saline for oral administration.

30 **Example 5**

Production of Clostridial spores, germination of spores *in vitro* and *in vivo*, and colonization and immune response to *C. difficile* in animals given spores orally

Clostridial strain producing the heterologous peptide is allowed to grow anaerobically in Peptone-Yeast extract-glucose or another medium optimal for sporulation for 48-72h to ensure maximum conversion of the vegetative bacteria into spores during the stationary phase. The remaining vegetative bacteria are killed by heat or ethanol treatment, eliminated by the bacteriolytic enzymes lysozyme or lysostaphin and the remaining spores are purified by centrifugation.

#### Results:

Spores from *C. difficile* strain 630 were readily observed on 3-day old blood agar plates as well as in PYG medium, whereas spores from *C. perfringens* NCTC 8798 were found only during growth in Duncan Strong (DS) medium, after preincubation in Fluid Thioglycolate (FTG) medium.

Prepared spores (washed in ethanol and resuspended in sterile PBS) were checked for germination ability on plates (TCCFA) as well as in gnotobiotic rats. Feces from 1 week old rats were positive for bacterial growth in feces 1-2 days after receiving *C. difficile* or *C. perfringens* spores orally, confirming the ability of the spores to germinate also *in vivo* and lead to colonization of the animal gut.

Antisera from 5 rats colonised for one week by *C. difficile* were pooled and used for Western blotting of *C. difficile* protein extracts. Western blotting revealed immunological reactions to bands corresponding to the *C. difficile* S-layer proteins confirming that antibodies were produced against these *C. difficile* antigens upon feeding with spores, spore germination and colonization of the animals.

### Example 6

#### Production of transformed *C. difficile* spores for use in oral immunisation.

##### A. Capsules.

The *C. difficile* spores obtained according to Example 5 are mixed together with Mg stearate (1%) and lactose (30%), granulated in ethanol and compressed to tablets, containing  $10^6$  spores, which are filled into gelatine capsules.

##### B. Powder for suspension.

The *C. difficile* spores obtained according to Example 5 are mixed together with Mg stearate (1%) and carboxymethyl cellulose (25%), granulated in ethanol. The granulate is dried and dispensed into vials to give an amount of about  $10^6$  spores in each vial. For oral



administration the content of the vial is suspended in water or for example orange juice immediately before intake.

### Example 7

#### 5 Use of S-layer genes for epidemiological typing.

Present methods to detect and follow the spread of certain *C. difficile* strains in the environment and between infected patients (i.e. "fingerprinting") include e.g. serotyping and PCR ribotyping. PCR ribotyping is a PCR based approach to amplify the region between the 16S and 23S genes of *C. difficile*, and which has been shown to resolve and detect over 100  
10 different patterns or strains. Different serotypes are likely to represent differences of the surface-exposed proteins, i.e. variations of S-layer proteins among strains. Our results with PCR amplification and following cleavage with restriction enzymes indicate that this region is present in almost all serogroups and that the cleavage pattern also varies among these groups (see Example 1C, (ii)). Thus, a molecular method including PCR combined with restriction  
15 enzyme cleavage or direct sequencing of the variable part of the ORF1 or another part of this segment may be a method which is faster and more reliable than serotyping and in particular also more reliable than PCR ribotyping for fingerprinting.

### Example 8

#### 20 Vaccination against CDAD

Immunity to CDAD after an episode of the infection is regarded to be short (months). This may be due to that anti-toxin antibodies are mainly of the serum IgG classes and not the secreted IgA class made to protect the gut mucosal surface, because the toxins are internalized by the gut mucosal cells (see above) and not by the M-cells specialized in furthering an IgA  
25 response. A further problem may be that immunity to the 20 *C. difficile* S-layer serotypes is required for prevention of colonization and thus the best protection against infection. For these reasons, it is likely that injectable vaccines against CDAD based on the toxins and under development may turn out to offer poor protection.

As an alternative, we provide a polyvalent live oral vaccine containing (i) the most  
30 prevalent toxin-producing serotypes (S-layer variants), here attenuated by knock-out of their toxin genes, and (ii) carrying a recombinant ORF1-*secA* cassette encoding relevant parts of the toxin genes and (iii) an adjuvant peptide ensuring uptake of the immunogenic toxin epitopes by e.g. M-cells in order to obtain an IgA anti-toxin response.

**Example 9**

Analysis of extracellular and membrane fraction proteins in *Clostridium difficile* by two-dimensional PAGE, N-terminal sequencing and data base searches – identification of genes encoding the S-layer proteins

Identification of extracellular proteins

Analysis of its extracellular protein pattern by 2-D PAGE showed that two proteins of 50 kDa and 36 kDa were very abundant. Subsequent analysis of membrane preparations from *C. difficile* VPI 10643 corroborated the almost exclusive fractionation of the 50 kDa and 36 kDa proteins into the membrane fraction, when compared with the soluble fraction. Also in strain 630 two proteins with similar molecular weights but with different pI were abundant in the extracellular as well as in the membrane fraction. Thus, the 50 kDa and 36 kDa proteins were likely to constitute the *C. difficile* surface(S)-layer proteins, which are known to be partially shed from the bacterial surface into the culture supernatant (Luckevich and Beveridge, 1989; Tsukagoshi *et al.*, 1984). The N-terminal sequence of spot no. 1 from VPI 10463 did not show any homology to other proteins in the *C. difficile* strain 630 genome database (Table 3). The N-terminal part of spot no. 2 showed similarity to an open reading frame encoding a 72 kDa protein in the *C. difficile* genome database (Table 3; ORF1, see also Figure 1). However, only nine out of 15 amino acids matched close to the N-terminus of ORF1. Strikingly, the N-terminal sequences of the corresponding proteins from strain 630 were different from those of VPI 10463 and both matched to ORF1 but at two different positions (spot No.10 and 11 in Table 1).

Several proteins were specifically found in PY cultures, i.e. during high toxin production (Table 3, spot no. 3, 4, 5, and 6). The N-terminal sequence of spot no. 3 matched with an ORF of 47.5 kDa in the *C. difficile* genome database. This ORF showed weak homology to a hypothetical protein in the *Plasmodium falciparum* genome database. The N-terminal sequence of spot no. 4 matched with an ORF of 39 kDa that showed homology to a phage-like element PBSX protein (XkdK) from *Bacillus subtilis*. The N-terminal sequences of spot no. 5 and 6 matched with two ORFs of 38 and 22 kDa, respectively, and these ORFs had the highest similarity to the FixB and FixA proteins from *Escherichia coli*. Spot no. 7, 8 and 9 were absent in PY but abundant in culture medium from PYG cultures. Spot no. 7 and 8 matched to an ORF located on the same contig as ORF1 (Table 3; ORF3). The N-terminus of

spot no. 9 matched to a central part of ORF1, and is likely to be a proteolytic fragment of a protein encoded by ORF1.

#### Analysis of the surface layer genes

5           The identification of the S-layer genes revealed a genomic segment including seven genes (ORF1, 3, 5-7, 9 11) with significant homology to N-acetyl muramoyl L-alanine amidase (CwlB/LytC) and modifier protein of major autolysin (LytB) from *Bacillus subtilis* (Fig.1, Table 4). In addition to the LytB/LytC similarity, the N-terminal part of ORF6 showed similarity to eukaryotic cysteine proteases, and the highly expressed ORF1 (above) showed  
10 weak similarity to S-layer proteins from *Lactobacillus* and *Streptococcus* spp. (Fig. 1). The N-terminus of ORF1 contained a typical signal peptide for export via the Sec-dependent secretion and the predicted cleavage site was identical to that found in the protein sequence (not shown). However, no typical protein cleavage site was identified within ORF1 that would allow processing of the 72 kDa protein further to give the finally sized S-layer proteins found  
15 (50 and 36 kDa). Strikingly, no significant match between the *C. difficile* S-layer ORFs and the S-layer homology motif (SLH domain) found in all presently known S-layer proteins was obtained (not shown). Most of the remaining genes in this genomic segment showed similarity to genes involved in secretion, polysaccharide and capsule synthesis (Fig. 1; Table 4). At least 2 other genomic sequence segments were found that contained genes similar to CwlB/LytC,  
20 indicating a complex variability (not shown).

#### **Summary of results**

          The most dominant surface-exposed protein in many bacterial species is the S-protein. This protein crystallizes into a regular monolayer on the outside surface of the bacteria: the S-  
25 layer. The S-layers satisfy multiple roles for the cell and function as protective coats, as structures involved in cell adhesion and surface recognition, as molecular sieves, as molecular and ion traps, as scaffolding for enzymes and as virulence factors (Sleytr and Beveridge, 1999; Sara and Sleytr, 2000). Though all S-layers share general features (all are made of relatively large proteins, self-assemble and are paracrystalline), comparative studies indicate  
30 that S-layers are non-conserved structures and are of limited taxonomical value (Kuen and Lubitz, 1996; Sleytr et al. 1999). Chemical analysis and genetic studies of a variety of S-layers have shown that they are composed of a single, homogenous protein or glycoprotein species with molecular weights ranging from 40 to 170 kDa. The S-layers of *Clostridium*

*difficile* (Takeoka et al., 1991) and *Bacillus anthracis* (Mesnage et al., 1998; Etienne-Toumelin et al., 1995) consist of two types of S-layer subunits which together form a defined lattice type but do not cross-react with polyclonal antibodies. Typically, S-layer proteins are often weakly acidic proteins (pIs between 4 to 6), containing 40-60% hydrophobic amino acids, and possess few or no sulfur-containing amino acids (Messner, 1996). S-protein production is directed by single or multiple promoters in front of the S-protein gene, yielding stable mRNAs. Most bacteria secrete S-proteins via the general secretory pathway (*sec*-pathway). Silent S-protein genes have been found in *Campylobacter fetus* and *Lactobacillus acidophilus*. These silent genes are placed in the expression site in a fraction of the bacterial population via inversion of a chromosomal segment (Boot and Pouwels, 1996).

The S-layer has been detected in some *C. difficile* strains and preliminary characterization has been done from *C. difficile* C253 (Mauri et al., 1999). Another independent study purified and identified the S-layer subunits from *C. difficile* GAI 0714 (Takeoka et al., 1991). In both cases, the S-layer has been shown to be composed of two different protein subunits with apparent molecular weights of 36 kDa and 47 kDa (*C. difficile* C253) and 32 kDa and 45 kDa (*C. difficile* GAI 0714). The S-layer proteins from VPI 10463 and strain 630 was here found to be similar in size but with significant pI differences. The N-terminal sequences varied significantly especially for the larger protein. The N-terminal sequences as determined for these proteins also indicate that they are not identical. Those from strain 630 appear to be processed products from the same gene (ORF1, Table 3). The N-terminal sequences of spot 1 from VPI 10643 did not find any homologue in the strain 630 database corroborating earlier results that the higher molecular weight S-layer protein was sero-group specific (VPI 10643, group G and strain 630, group X) (Takeoka et al., 1991). Our current work indicated that this arrangement of two S-layer proteins was also true for all the different serotypes tested (data not shown). The spot 2 found a partial match with another ORF (ORF3) in the same contig as ORF1.

Other ORFs located in the same contig also had similarities with ORF1 and ORF3, whose C-terminal parts showed similarities to N-acetyl muramoyl L-alanine amidase (CwlB/LytC) and modifier protein of major autolysin (LytB) from *Bacillus subtilis* (Lazarevic et al., 1992), whereas the N-terminal part showed weak similarities to surface-layer proteins from *Lactobacillus helveticus* (Callegari et al., 1998) and *Streptococcus* spp. It is interesting to note that N-acetyl muramoyl L-alanine peptidoglycan amidase is the major autolysin of *B. subtilis* and has high affinity for cell walls, which is enhanced by the modifier protein, but

small amounts of cell free autolysin can be detected in cultures of *B. subtilis*. Thus, the amidase-like motif that appears to be typical of *C. difficile* S-layer proteins probably confers their anchorage to the cell wall peptidoglycan-teichoic acid.

Considering the highly competitive situation of closely related organisms in their natural habitats, it is obvious that the S-layer surface has to contribute to diversification rather than to conservation. With respect to this, the importance of S-layer variation leading to the expression of alternative S-layer genes under different stress factors such as those imposed by the immune system of a host in response to an S-layered pathogen or drastic changes in the growth and environmental conditions for nonpathogens is conceivable (Dworkin and Blaser, 1997; Sara et al., 1996). This could probably explain the variation in S-layer proteins even amongst the same species as in the case of *C. difficile*.

Identification of spot 4 as having similarity with XkdK, a protein encoded by the phage-like element PBSX from *Bacillus subtilis* (Krogh et al., 1996) and being located in a contig with other ORFs having similarity with other PBSX encoded proteins is very interesting. PBSX is a bacteriophage-like bacteriocin, or phibacin, of *B. subtilis* 168 (Okamoto et al., 1968). When *B. subtilis* 168 cells are exposed to treatments that induce the SOS response (such as UV light, mitomycin C), the cells lyse after incubation of 1h and release particles of PBSX (Seaman et al., 1962). The spot 4 is completely absent in PYG supernatants. Taken together this could indicate that toxin production (high in PY) and expression of this phage-like protein in *C. difficile* is a response to certain stress, environmental or otherwise, that decides whether it will resort to toxin expression, sporulation or both.

The N-terminal sequences of spots 7 (41 kDa) and 8 (38 kDa) are identical (Table 3) and correspond to the same ORF (ORF3, Fig 1), whose N-terminal part is similar to N-acetyl muramoyl L-alanine amidase (CwlB/LytC) from *B. subtilis*. However, the size of the proteins in the gel do not match the size expected from the ORF3 (encodes a 67.5 kDa protein). Both ORF1 and ORF3 have clear signal sequences at the beginning which is missing in the protein spots sequenced, thus indicating that these are indeed secreted and processed following translation. This could also possibly explain the difference in size and pI for the two different spots, and the discrepancy between expected and observed molecular weights on SDS-gels.

The spot 9 (24 kDa) has a N-terminal sequence (Table 1) which corresponds to an internal fragment of ORF1. The expected size of this fragment is around 21 kDa which corresponds closely with what is observed experimentally. Clearly there are post-translational processing events which could be enacted in the cell envelope or in the supernatant. It is

important however to note that spots 7-9 are present in PYG supernatants only, when the cells start sporulating.

The spots 9 and 10 are also processed products from ORF1 and are present in both PY supernatant and membrane fractions. However, these samples are obtained from strain 630.

5 The results obtained thus far indicate that this operon (contig) (Fig 1) is present in both VPI 10643 and strain 630, but different ORFs are expressed by the two strains.

## Experimental procedures

### Strains and growth medium

10 The toxin-producing *C. difficile* strain VPI 10463 (CCUG 19126, Culture Collection, University of Göteborg, Sweden) was grown in either PY, PYG (purchased from the Karolinska Hospital, Stockholm, Sweden) or SDM medium. SDM is identical to MADM (Karasawa *et al.*, 1995; Yamakawa *et al.*, 1994; Yamakawa *et al.*, 1996), except that the concentrations of glycine and threonine were 100 mg/L and 200 mg/L, respectively, and that  
15 Ca-D-pantothenate, pyridoxine and biotin were used as the sole vitamin sources. PY(G) was prepared by adding cysteine (500 mg/L), boiling for 20 min while purging with an anaerobic gas mixture (10% CO<sub>2</sub>, 10% H<sub>2</sub>, 80% N<sub>2</sub>) for 20 min, sterilised by filtration (Acrodisc, Gelman sciences) and aliquoted into tubes with serum vial-style necks (Bellco Glass) while flushing with anaerobic gas. The tubes were closed with butyl stoppers secured with  
20 aluminium crimp seals. SDM was prepared accordingly.

### Growth conditions, sampling and optical density measurements

For each experiment, a tube containing 20 ml SDM was inoculated with 0.2 ml thawed bacterial suspension (stored at -70°C) using a syringe and a needle that was passed through  
25 the rubber septum of the tube. To avoid entry of O<sub>2</sub>, the syringe was equilibrated with anaerobic gas before inoculation. The tube was put horizontally on a rotary shaker (50 rpm, 37 °C), and on the next day, the culture was serially diluted into PY or PYG. On day three, samples were collected from the diluted cultures and OD was measured at 600 nm using a Hitachi U-1100 spectrophotometer.

30

### Sample preparation and membrane fractionation

Culture samples were centrifuged at 16000 x g for 3 min, whereafter the supernatants were removed, filtered, and stored at -20°C for later analysis. The pellet was frozen at -20°C

for 30 min or longer, thawed, dissolved in 1 ml sterile water and sonicated on ice for 3 x 30 s at 100 W (Labsonic 1510, B. Braun). Larger cell pellets, obtained from >1 ml culture, was sonicated for longer times. The cell extracts were centrifuged at 5000 x g for 5 min. The pellet was separated as the low speed pellet (LSP), and the supernatant was further centrifuged at 50000 x g for 20 min. The pellet was separated as the high speed pellet (HSP), and the supernatant (soluble fraction) was stored at -20°C. The LSP and the HSP were resuspended in 1x PBS (Phosphate buffered saline). Protein amount was measured using a kit (Biorad) and a BSA standard curve according to the manufacturer's instructions.

The culture supernatants were precipitated using trichloroacetic acid (TCA) to a final concentration of 10%. The pellets were washed with ice-cold Acetone, air dried and finally resuspended in 1x PBS to obtain the extracellular protein fraction. Protein estimation and analysis was carried out as described earlier.

#### Immunoprecipitation

Immunoprecipitation was performed in microtiter wells coated with antibodies against toxin A (PCG-4, r-Biopharm) or toxin B (~~xxx~~, r-Biopharm), Ten µg/ml antibody in 0.04 M Na<sub>2</sub>CO<sub>3</sub>, 0.06 M NaHCO<sub>3</sub>, pH 9.6 was added to microtiter wells and incubated for 1 h at 37°C and washed four times with PBS containing 0.05% (v/v) Tween-20, pH 7.4. The wells were loaded with cell extract, culture supernatant medium or PBS (negative control), incubated 90 min at 25°C, and washed four times with PBS. After addition of 50 µl SDS sample buffer solution (below) and heating for 5 min at 95°C, the precipitated proteins were analysed by SDS-PAGE.

#### SDS-polyacrylamide gel electrophoresis (SDS-PAGE)

SDS-PAGE was performed using pre-cast polyacrylamide gels (ExcelGel 8-18% gradient gels, Pharmacia Biotech) and a Multiphor II horizontal slab gel apparatus (Pharmacia Biotech) according to the manuals provided by the manufacturer. The samples were mixed 1:1 with SDS sample buffer solution (0.05 M Tris, 1% (w/v) SDS, 10 mM DTT, 0.01% (w/v) bromophenol blue, pH 7), incubated 5 min at 95°C, loaded onto the gels and run at 15°C. Chemicals were obtained from Sigma, and molecular weight markers from Pharmacia Biotech. The gels were stained with silver (PlusOne, Pharmacia Biotech) using a Hoefer automatic gel stainer (Pharmacia Biotech), digitised by scanning (Scanjet 3c/T, Hewlett-Packard), and transferred to ClarisDraw (Claris Software) on a Macintosh computer.

### Immunoblotting

Proteins were separated by SDS-PAGE transferred to polyvinylidene fluoride membranes (Immobilon P<sup>SEQ</sup>, Millipore) using the Pharmacia Novablot transfer equipment and a continuous buffer system (39 mM glycine, 48 mM Tris, 0.0375% (w/v) SDS, 20% (v/v) methanol) according to the Multiphor II manual. The membranes were dried at 25°C for 1.5 h, blocked with 0.5% Tween-20 for 20 min, and then incubated with toxin A or toxin B antibodies (r-Biopharm, 0.2 µg/ml in TST buffer containing 0.05 M Tris, 0.5 M NaCl, 0.1% Tween-20, pH 9) for 1 h. After three washes in TST, the membranes were incubated with horse-radish peroxidase conjugated anti-mouse antibodies (DAKOPATTS, diluted 1:10000 in TST) for 1 h and washed three times in TST. A chemiluminiscent signal (ECL Plus, Amersham) was used to detect the bands. The relative amount of toxin B was measured on scanned x-ray films using the Molecular Analyst software (Biorad).

### Two-dimensional polyacrylamide gel electrophoresis (2-D PAGE)

Protein samples were obtained as described under sample preparation. For 2-D PAGE, 40 ml aliquots of each sample was mixed with 160 ml of buffer III [9.9M Urea, 4% (v/v) Igepal CA630, 2.2% (v/v) ampholytes 3-10, 100mM DTT, 2% (w/v) CHAPS]. Protein mixtures were focused at 20°C on 180mm IPG Drystrip pH 4-7 (Pharmacia Biotech) using the Multiphor II 2-D gel Kit according to the manufacturer's instructions. The second dimension was run on 12% (linear) SDS acrylamide gels. The gels were stained with silver (PlusOne, Pharmacia Biotech) using a Hoeffler automatic gel stainer (Pharmacia Biotech). The chemicals were obtained from Sigma except for Pharmalytes (Pharmacia Biotech).

### N-terminal sequencing

For N-terminal Sequence determination, gels were transferred to Immobilon-P polyvinylidene difluoride membrane (Millipore) as described under immunoblotting. The membrane was stained with Coomassie blue; protein spots were excised for sequence determination. The protein spots cut from the transfer membrane were washed four times in 10% methanol and then dried and frozen. N-terminal sequence analysis was performed at the Protein Analysis Center, Karolinska Institute. Peptide sequences were matched against the *C.difficile* genome database (Sanger Center, UK).



## References

- Bardwell, J.C.A., Lee, J.-O., Jander, G., Martin, N., Belin, D., and Beckwith, J. (1993) A pathway for disulfide bond formation *in vivo*. *Proc Natl Acad Sci USA* 90: 1038-1042.
- Bardwell, J.C.A., McGovern, K., and Beckwith, J. (1991) Identification of a protein required  
5 for disulfide bond formation *in vivo*. *Cell* 65: 581-589.
- Boot, H.J., and Pouwels, P.H. (1996) Expression, secretion and antigenic variation of bacterial S-layer proteins. *Mol Microbiol* 21: 1117-1123.
- Callegari, L., Riboli, B., Sanders, P., Coconelli, S., Kok, J., Venema, G., and Morelli L. (1998) The S-layer gene of *Lactobacillus helveticus* CNRZ 892 : cloning, sequencing and  
10 heterologous expression. *Microbiology* 144: 719-726.
- Cerquetti, M., Pantosti, A., Stefanelli, P., and Mastrantonio, P. (1992) Purification and characterization of an immunodominant 36 kD antigen present on the cell surface of *Clostridium difficile*. *Microb Pathogen* 13: 271-279.
- Dauter, Z., Wilson, K.S., Sieker, L.C., Moulis, J.M., and Meyer, J. (1996) Zinc- and iron-rubredoxins from *Clostridium pasteurianum* at atomic resolution: a high-precision model  
15 for ZnS<sub>4</sub> coordination unit in a protein. *Proc Natl Acad Sci USA* 93: 8836-8840.
- Dworkin, J., and Blaser, M.J. (1997) Molecular mechanisms of *Campylobacter fetus* surface layer protein expression. *Mol Microbiol* 26: 433-440.
- Economou, A. (1999) Following the leader : bacterial protein export through the Sec pathway.  
20 *Trends Microbiol* 7: 315-319.
- Etienne-Toumelin, I., Sirard, J.-C., Duflot, E., Mock, M., and Fouet, A. (1995) Characterization of the *Bacillus anthracis* S-layer : cloning and sequencing of the structural gene. *J. Bacteriol.* 177 : 614-620.
- Garnier, T., and Cole, S.T. (1988) Studies of UV-inducible promoters from *Clostridium perfringens* *in vivo* and *in vitro*. *Mol Microbiol* 2: 607-614.  
25
- Izard, J.W., and Kendall, D.A. (1994) Signal peptides: exquisitely designed transport promoters. *Mol Microbiol* 13: 765-73.
- Kamiya, S., Ogura, H., Meng, X.Q., and Nakamura, S. (1992) Correlation between cytotoxin production and sporulation in *Clostridium difficile*. *J Med Microbiol* 37: 206-210.
- 30 Karasawa, T., Ikoma S., Yamakawa, K., and Nakamura, S. (1995) A defined medium for *Clostridium difficile*. *Microbiology* 141: 371-375.

- Kawata, T., Takeoka, A., Takumi, K., and Masuda, K. (1984) Demonstration and preliminary characterization of a regular array in the cell wall of *Clostridium difficile*. *FEMS Microbiol Lett* **24**: 232-328.
- 5 Klauser, T., Pohlner, J., and Meyer, T.F. (1993) The secretion pathway of IgA protease-type proteins in gram-negative bacteria. *Bioessays* **15**: 799-805.
- Krogh, S., M. O'Reilly, N. Nolan, and K.M. Devine. (1996) The phage-like element PBSX and part of the skin element, which are resident at different locations on the *Bacillus subtilis* chromosome, are highly homologous. *Microbiology* **142** : 2031-2040.
- 10 Lazarevic, V., Margot, P., Suldo, B., and Karamata, D. (1992) Sequencing and analysis of the *Bacillus subtilis* *lytRABC* divergon : a regulatory unit encompassing the structural genes of the N-acetylmuramoyl-L-alanine amidase and its modifier. *J Gen Microbiol* **138**: 1949-1961.
- Luckevich, M.D., and Beveridge, T.J. (1989) Characterization of a dynamic S-layer on *Bacillus thuringiensis*. *J. Bacteriol.* **171** : 6656-6667.
- 15 Masuda, K., Itoh, M., and Kawata, T. (1989) Characterization and reassembly of a regular array in the cell wall of *Clostridium difficile* GAI 4131. *Microbiol Immunol* **33**: 287-298.
- Mauri, P.L., Pietta, P.G., Maggioni, A., Cerquetti, M., Sebastianelli, A., Mastrantonio, P. (1999) Characterization of surface layer proteins from *Clostridium difficile* by liquid chromatography/ electrospray ionization mass spectrometry. *Rapid Commun. Mass Spectrom.* **13**: 695-703.
- 20 McFarland, L.V., Elmer, G.W., Stamm, W.E., and Mulligan, M.E. 1991. Correlation of immunoblot type, enterotoxin production, and cytotoxin production with clinical manifestations of *Clostridium difficile* infection in a cohort of hospitalized patients. *Infect Immun* **59**: 2456-2462.
- 25 Mesnage, S., Tosi-Couture, E., Gounon, P., Mock, M., and Fouet, A. (1998) The capsule and the S-layer: two independent and yet compatible macromolecular structures in *Bacillus anthracis*. *J. Bacteriol.* **180** : 52-58.
- Messner, P. (1996) Chemical composition and biosynthesis of S-layers, p.35-76. In U.B.Sleytr, P.Messner. D.Pum, and M.Sara (ed.), *Crystalline bacterial cell surface layer proteins (S-layers)*. Academic Press, Landes Company, Austin, Tex.
- 30 Nicholls, L., Grant, T.H., and Robins-Browne, R.M. (2000) Identification of a novel genetic locus that is required for *in vitro* adhesion of a clinical isolate of enterohaemorrhagic *Escherichia coli* to epithelial cells. *Mol Microbiol* **35**: 275-288.

- Orme-Johnson, W.H., and Orme-Johnson, N.R. (1978) Overview of iron-sulfur proteins. *Methods Enzymol* 53: 259-268.
- Pugsley, A.P. (1993) The complete general secretory pathway in Gram-negative bacteria. *Microbiol Rev* 57: 50-108.
- 5 Saïra, M., and Sleytr, U.B. (2000) S-layer proteins. *J. Bacteriol.* 182 : 859-868.
- Saïra, M., Kuen, B., Mayer, H.F., Mandl, F., Schuster, K.C., and Sleytr, U.B. (1996) Dynamics in oxygen induced changes in S-layer protein synthesis from *Bacillus stearothermophilus* PV72 and the S-layer deficient variant T5 in continuous culture and studies on the cell wall composition. *J. Bacteriol.* 178 : 2108-2117.
- 10 Sellman, B.T., and Tweten, R.K. (1997) The propeptide of *Clostridium septicum* alpha toxin functions as an intramolecular chaperone and is a potent inhibitor of alpha toxin-dependent cytolysis. *Mol Microbiol* 25: 429-440.
- Simonen, M., and Palva, I. (1993) Protein secretion in *Bacillus* species. *Microbiol Rev* 57: 109-137.
- 15 Takemaru, K., Mizuno, T. Sato, M. Takeuchi, and Y. Kobayashi. (1995) Complete nucleotide sequence of a skin element excised by DNA rearrangement during sporulation in *Bacillus subtilis*. *Microbiology* 141 : 323-327.
- Takeoka, A., Takumi, K., Koga, T., and Kawata, T. (1991) Purification and characterization of S layer proteins from *Clostridium difficile* GAI 0714. *J Gen Microbiol* 137: 261-267.
- 20 Takumi, K., Takeoka, A., and Kawata, T. (1987) Purification and immunochemical properties of a wall protein antigen from *Clostridium difficile* ATCC 110011. *Microbiol Immunol* 31: 837-849.
- Tsukagoshi, N., Tabata R., Takemura, T., Yamagata, H., and Udaka, S. (1984) Molecular cloning of a major cell wall protein gene from protein producing *Bacillus brevis* 47 and its
- 25 expression in *Escherichia coli* and *Bacillus subtilis*. *J. Bacteriol.* 158 : 1054-1060.
- Welch, R.A. (1991) Pore-forming cytolysins of gram-negative bacteria. *Mol Microbiol* 5: 21-528.

Table 1. Identification of ORF 1-12 and sequences upstream of ORF1 in SEQ ID NO: 35 and predicted signal peptide sequences of the S-layer ORFs. See Figures 1 and 2.

Gene	Frame	from	to	Total bp (aa)	Homology/homologous motif (aa domain)	Predicted signal peptide**	SEQ ID NO:
A	+2	227	1366	1140	tRNA- Guanine transglycolase		
B	+3	1500	1793	294	conserved protein (membrane bound)		
C	+3	1851	2213	363	Hypothetical protein YrzE		
D	+1	3046	4941	1896 (631)	N-acetylmutamoyl-L-alanine amidase (aa 320-610)	MLSNKKRSMALVMA GATVMSAAAPIFA↓DNTVTEN	8
E	+3	5373	6806	1434 (477)	N-acetylmutamoyl-L-alanine amidase (aa 20-290)	MKSTLGVENNMKNKSKILAIGLITLFLVMVNTPMV SA↓LTSVE	9
F	+2	7187	13150	5964	Unknown		
G	+1	13360	15240	1881 (626)	N-acetylmutamoyl-L-alanine amidase (aa 320-620)	MNKRKSFRTIAVSTMVA VVTGSATCAYA↓APVLQ	10
H	+1	15556	17208	1653 (550)	N-acetylmutamoyl-L-alanine amidase (aa 5-370)	MENNHNINIKYKNHQGDGMKNMKNKILSLGLAVSLIL VNFKSVNA↓SSVV	11
I	+3	17427	19013	1587 (528)	N-acetylmutamoyl-L-alanine amidase (aa 20-250)	MKVNKRVL SIGLAISLIMAGAPNINA↓LSSIEK	12
ORF1	+3	19224	21383	2160 (719)	N-acetylmutamoyl-L-alanine amidase (aa 380-719)	MNKKNIAMSGLT V LASAAPVFA↓ATTGT	1
ORF2	+1	21628	23973	2346	SecA		34
ORF3	+1	24388	26259	1872 (622)	N-acetylmutamoyl-L-alanine amidase (aa 310-540)	MNKKNL SVMAAAMISTSVAPVFA↓AETTQ	2
ORF4	+2	26420	27073	654	Unknown		
ORF5	+1	27106	28938	1833 (610)	N-acetylmutamoyl-L-alanine amidase (aa 5-300)	MKISKKIVSLLTMTFLT V TLYGNTSNA↓STKDT	3
ORF6	+3	29613	32024	2412 (803)	N-acetylmutamoyl-L-alanine amidase (aa 500-800)	MRKYKSKKL SKLLALSTVCF LIVSTIPVSA↓ENHK	4
ORF7	+2	32321	33898	578 (525)	N-acetylmutamoyl-L-alanine amidase (aa 30-340)	MKAPKTLTILTIALT LSSISIPSYA↓LTEEK	5
ORF8	+3	34032	35351	1320	GDT1 protein		
ORF9	+1	35590	37629	2040 (528)	N-acetylmutamoyl-L-alanine amidase (aa 5-285)	MRGDMMKKTTKLLATGMLSVAMVAPNVVALA↓AENTT	6
ORF10	+1	37873	38535	699	Hypothetical protein		
ORF11	+2	38630	39685	1056 (351)	N-acetylmutamoyl-L-alanine amidase (aa 40-340)	MIKKISTLSVL LISISSTIGVFA↓DANPKR	7
ORF12	+2	39800	41356	1557	Virulence Factor MvIN		

aa = no. of amino acids or their position in translated ORF

↓ = predicted cleavage point

Table 2. Possible promoter starts between ORF I and ORF 1. (1.00 is highest score). ORF 1 start codon is at position 19224.

Position	Score	Sequence	SEQ ID NO:
19026-19071	0.91	TAGTTTATTACATTTTAAAAATTTAGGGTATAAAAACTTGTAACTTGGAG	13
19036-19081	1.00	CATTTTAAAAATTTAGGGTATAAAAACTTGTAACTTGGAGAAAATAATAA	14
19059-19104	0.80	AACTTGTAAACTTGGAGAAAATAATAATTTAAAAAATAGCTTGCAAAA	15
19067-19112	0.99	AACTTGGAGAAAATAATAATTTAAAAAATAGCTTGCAAAAAGAAATAAAA	16
19083-19128	1.00	TAATTTAAAAAATAGCTTGCAAAAAGAAATAAATGGATTATTATAGAG	17
19094-19139	0.99	AATAGCTTGCAAAAAGAAATAAATGGATTATTATAGAGATGTGAGAAAT	18
19123-19168	0.96	TATTATAGAGATGTGAGAAATAATTagGaATATATATGGATGATTATTCTATG	19
19132-19177	0.85	GATGTGAGAAATAATTagGaATATATGGATGATTATTCTATGtAcATAATA	20
19161-19206	0.84	GATTATCTATGtAcATAATAAGAGATGTAAATTTTAATATAATGTTGGG	21

Table 3. Extracellular *C. difficile* proteins identified by N-terminal sequence analysis and genome database searches.

Spot	MW	Source	N-terminal Sequence <sup>a</sup>	Highest match <sup>b</sup>	Comment
1	50	PY	AAKASIADENSPVKLT LKSDXKKDL	No match	
2	36	PY	DDTKVETGDQGYTVV	N-acetyl---(ORF1)	Partly matched
3	47	PY	SEKEILTARLAV	Hypothetical protein	Fragment
4	40	PY	AIGLPSINISSK	PBSX phage protein, XkdK	
5	39	PY	MXDIKLDXFXKX	Fix B	
6	30	PY	MKILVXVKQVXX	Fix A	
7	41	PYG	AETTQVKKETIT	N-acetyl muramoyl-L-alanine amidase (ORF3)	Fragment
8	38	PYG	AETTQVKKETIT	N-acetyl---(ORF3)	Fragment
9	24	PYG	TSLKIADEVGLD	N-acetyl---(ORF1)	Fragment
10	48	PY	ANDTIASQDTPAKVV	N-acetyl---(ORF1)/S-layer protein	Fragment (central part)
11	35	PY	ATTGTQGYTVVKNDD	N-acetyl---(ORF1)/S-layer protein	Fragment (N-terminal)

<sup>a</sup>Underlined sequence indicates that only this part was found to be identical with the ORF in the *C. difficile* strain 630 genome database

<sup>b</sup>Search was made in the *C. difficile* strain 630 genome database by tBLASTn algorithm at www.. Contigs were exported to ORF finder at NCBI, and the entire ORF was subjected to further BLAST searches against the redundant database.

Table 4. Blast search summary of the ORFs in the *C. difficile* strain 630 S- layer contig. ORF numbers refer to those in Fig 1.

ORF	Length (amino acids)	Match (highest E-value) <sup>1</sup>	Source	E value
1	719	N-acetyl/muramoyl-L-alanine amidase	<i>B. subtilis</i>	$1 \times 10^{-23}$
2	781	Preprotein translocase SecA subunit	<i>B. subtilis</i>	0.0
3	623	N-acetyl/muramoyl-L-alanine amidase	<i>B. subtilis</i>	$1 \times 10^{-24}$
4	217	No match		
5	610	N-acetyl/muramoyl-L-alanine amidase	<i>B. subtilis</i>	$2 \times 10^{-42}$
6	803	N-acetyl/muramoyl-L-alanine amidase	<i>B. subtilis</i>	$2 \times 10^{-36}$
7	525	N-acetyl/muramoyl-L-alanine amidase	<i>B. subtilis</i>	$2 \times 10^{-39}$
8	439	GDT1 protein	<i>D. discoideum</i>	$2 \times 10^{-11}$
9	679	N-acetyl/muramoyl-L-alanine amidase	<i>B. subtilis</i>	$4 \times 10^{-56}$
10	232	Hypothetical protein	<i>B. subtilis</i>	$7 \times 10^{-45}$
11	351	N-acetyl/muramoyl-L-alanine amidase	<i>B. subtilis</i>	$1 \times 10^{-11}$
12	518	Virulence factor MviN	<i>S. typhimurium</i>	$2 \times 10^{-53}$
13	568	Phosphomannose murase	<i>B. subtilis</i>	$10^{-159}$
14	352	Mannose-1-phosphate guanylttransferase	<i>E. coli</i>	$5 \times 10^{-68}$
15	157	Cap8J	<i>S. pneumoniae</i>	$5 \times 10^{-6}$
16	385	No match		
17	279	Glucosyl transferase	<i>S. pneumoniae</i>	$5 \times 10^{-11}$

<sup>1</sup>Database search was made using the BLAST algorithm at <http://www.ncbi.nlm.nih.gov/>.

Matches with a higher E value than 0.001 was regarded as no match.

## CLAIMS

1. A gene expression cassette comprising a secretory leader sequence encoding a signal peptide from *Clostridium difficile* having an amino acid sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7 and signal peptides of analogous exported clostridial N-acetylmuramoyl-L-alanine amidase-like proteins, linked to a DNA sequence encoding a heterologous polypeptide.
2. A gene expression cassette according to claim 1, wherein the signal peptides of the analogous clostridial N-acetylmuramoyl-L-alanine amidase-like proteins are selected from *Clostridium difficile* signal peptides having an amino acid sequence of any one of SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, and SEQ ID NO: 12.
3. A gene expression cassette according to claim 1 or 2, which further includes a promoter of prokaryotic origin.
4. A gene expression cassette according to claim 3, wherein the promoter is selected from clostridial promoters comprising a nucleotide sequence selected from the group consisting of SEQ ID NO: 13 – 21.
5. A gene expression cassette according to any one of claims 1-4 which further includes a DNA sequence encoding at least a cell wall binding portion of a protein of prokaryotic origin functioning in clostridia such that a fusion polypeptide may be presented on the outer surface of a host cell harbouring the cassette.
6. A gene expression cassette according to any one of claims 1-4, which further includes a DNA sequence encoding at least a functional cell wall binding portion of an S-layer protein of *C. difficile* selected from any one of the polypeptides having an amino acid sequence selected from SEQ ID NO: 22 - 33 such that a fusion polypeptide may be presented on the outer surface of a host cell harbouring the cassette.



7. A gene expression cassette according to claim 5 or 6, wherein DNA encoding the cell wall binding portions of SEQ ID NO: 22-33 has been omitted such that the fusion peptide is secreted into the surrounding milieu by the host cell harbouring the cassette.

5 8. A gene expression cassette according to claim 6 or 7, wherein the DNA sequence encoding the heterologous peptide is inserted at a point downstream the first (signal) proteolytic cleavage site in the gene encoding a polypeptide having an amino acid sequence selected from SEQ ID NO: 22 – 33, optionally including or excluding its second cleavage site.

10 9. A gene expression cassette according to any one of claims 1-8, which further comprises at least a functional part of a secretory (*secA*) gene recognizing the signal peptide, to allow translocation of a heterologous polypeptide and/or fusion polypeptide across the cytoplasmic membrane of a host cell harbouring the expression cassette.

15 10. A gene expression cassette according to claim 9, wherein the secretory gene is from *C. difficile* and encodes a polypeptide having the amino acid sequence SEQ ID NO: 34.

11. A gene expression cassette as shown in Figure 3.

20 12. A vector comprising a gene expression cassette as claimed in any one of claims 1-11.

13. A vector according to claim 12, wherein the vector is a plasmid.

14. A host organism transformed with a vector according to claim 12 or 13 for  
25 expression of the heterologous polypeptide and/or fusion polypeptide.

15. A *Clostridium* host organism transformed with a vector according to claim 12 or 13 for expression of the heterologous polypeptide and/or fusion polypeptide.

30 16. A host organism as claimed in claim 15 which is *C. difficile*.

17. A host organism as claimed in claim 15 which is *C. perfringens*.

18. A pharmaceutical or veterinary composition or formulation which comprises *Clostridial* cells transformed with a vector according to claim 12 or 13, with the ability to present on the cell surface and/or to secrete an expressed heterologous polypeptide or fusion polypeptide, together with a pharmaceutically or veterinary acceptable carrier or diluent.

5

19. A composition or formulation according to claim 18, which is suitable for oral or intranasal administration.

10

20. A composition or formulation according to claim 18 or 19, which further comprises, as an adjuvant, non-toxic motifs of the *C. difficile* toxins A and/or B that enable the expressed heterologous polypeptide and/or fusion polypeptide to enter the colonic mucosal cells of a mammal by receptor-mediated endocytosis, and/or a portion of toxin B responsible for its intracellular and intercellular spread.

15

21. A composition or formulation according to claim 18 or 19, which further comprises a further fused or separate carrier peptide or adjuvant, in addition to the expressed heterologous polypeptide and/or fusion polypeptide, to elicit a stronger or differently directed immune response than that against the expressed heterologous polypeptide acting alone.

20

22. A vaccine which comprises a *Clostridial* bacterium transformed with a vector according to claim 12 or 13 and capable of presenting on the surface of the bacterium and/or secreting an antigen in a human or animal body, and optionally also an adjuvant described in claim 20 or 21.

25

23. A medicament which comprises a *Clostridial* bacterium transformed with a vector according to claim 12 or 13 and capable of presenting on the surface of the bacterium and/or secreting a heterologous polypeptide and/or fusion polypeptide in a human or animal body, and optionally also an adjuvant described in claim 20 or 21.

30

24. A vaccine according to claim 22, which comprises a mixture of at least two differently engineered *Clostridia* strains, each capable of presenting on the surface of the bacterium and/or secreting a different heterologous polypeptide and/or fusion polypeptide.

25. A medicament according to claim 23, which comprises a mixture of at least two differently engineered *Clostridia* strains, each capable of presenting on the surface of the bacterium and/or secreting a different heterologous polypeptide and/or fusion polypeptide.

5 26. A vaccine which comprises spores of *Clostridia* cells or bacteria transformed with a vector according to claim 12 or 13, and capable of germinating into cells which are able to grow, express, and present or secrete a heterologous polypeptide and/or fusion polypeptide, and optionally an adjuvant described in claim 20 or 21, in a mammalian body.

10 27. A medicament which comprises spores of *Clostridia* cells or bacteria transformed with a vector according to claim 12 or 13, and capable of germinating into cells which are able to grow, express, and present or secrete a heterologous polypeptide and/or fusion polypeptide and optionally an adjuvant described in claim 20 or 21, in a mammalian body.

15 28. A vaccine according to claim 26, which comprises a mixture of spores from at least two differently engineered *Clostridia* strains.

29. A medicament according to claim 27, which comprises a mixture of spores from at least two differently engineered *Clostridia* strains.

20

30. A method for vaccination of a mammal, which comprises administering a therapeutically or prophylactically effective dose of a vaccine according to any one of claims 22, 24, 26 and 28 to the mammal.

25 31. A method for prophylactic or therapeutic treatment of a mammal, which comprises administering a therapeutically or prophylactically effective dose of a medicament according to any one of claims 23, 25, 27 and 29 to the mammal.

30 32. A vaccine according to claim 26 or 28, a medicament according to claim 27 or 29, or a method according to claim 30 or 31, wherein the spores are from *C. difficile* or *C. perfringens*.

33. A *C. difficile*-associated diarrhea (CDAD) vaccine comprising spores according to claim 26 or 32 and capable of expressing, after germination,

- (i) relevant parts of the *C. difficile* toxins, alone or together with a
- (ii) suitable adjuvant to provide primarily an IgA response to the toxin antigenic epitopes

5

and/or

- (iii) S-layer protein antigenic variants (serotype antigens) or fimbrial antigens from *C. difficile* to obtain, after administration to a mammal, a polyvalent anti-S-layer (or anti-fimbrial) IgA response preventing *C. difficile* colonization of the mammal.

10




Figure 1

2/7




Figure 2

3/7




Figure 3




Figure 4



5/7




Figure 5





Figure 6



→ Primers cof2 and cor3 designed to amplify orf1 & secA with native promoter

→ Primers hbf1, 2,3 and hbr4 designed to amplify Hbv



Figure 7

## SEQUENCE LISTING

<110> Smittskyddsinstitutet

<120> Gene expression cassette and its use

<130> 110060300

<140>

<141>

<150> SE 0002139-4

<151> 2000-06-07

<150> 0101479-4

<151> 2001-04-26

<160> 35

<170> PatentIn Ver. 2.1

<210> 1

<211> 29

<212> PRT

<213> Clostridium difficile

<400> 1

Met Asn Lys Lys Asn Ile Ala Ile Ala Met Ser Gly Leu Thr Val Leu  
1 5 10 15

Ala Ser Ala Ala Pro Val Phe Ala Ala Thr Thr Gly Thr  
20 25

<210> 2

<211> 29

<212> PRT

<213> Clostridium difficile

<400> 2

Met Asn Lys Lys Asn Leu Ser Val Ile Met Ala Ala Ala Met Ile Ser  
1 5 10 15

Thr Ser Val Ala Pro Val Phe Ala Ala Glu Thr Thr Gln  
20 25

<210> 3

<211> 32

<212> PRT

<213> Clostridium difficile

<400> 3

Met Lys Ile Ser Lys Lys Ile Val Ser Leu Leu Thr Met Thr Phe Leu  
1 5 10 15

Thr Val Thr Leu Tyr Gly Asn Thr Ser Asn Ala Ser Thr Lys Asp Thr  
20 25 30

<210> 4  
 <211> 34  
 <212> PRT  
 <213> Clostridium difficile

<400> 4  
 Met Arg Lys Tyr Lys Ser Lys Lys Leu Ser Lys Leu Leu Ala Leu Ser  
     1                    5                    10                    15  
 Thr Val Cys Phe Leu Ile Val Ser Thr Ile Pro Val Ser Ala Glu Asn  
             20                    25                    30

His Lys

<210> 5  
 <211> 32  
 <212> PRT  
 <213> Clostridium difficile

<400> 5  
 Met Lys Ala Pro Lys Thr Ile Leu Thr Ile Leu Thr Ile Ala Leu Thr  
     1                    5                    10                    15  
 Leu Ser Ser Ile Ser Ile Ile Pro Ser Tyr Ala Leu Thr Glu Glu Lys  
             20                    25                    30

<210> 6  
 <211> 35  
 <212> PRT  
 <213> Clostridium difficile

<400> 6  
 Met Arg Gly Asp Met Met Lys Lys Thr Thr Lys Leu Leu Ala Thr Gly  
     1                    5                    10                    15  
 Met Leu Ser Val Ala Met Val Ala Pro Asn Val Ala Leu Ala Ala Glu  
             20                    25                    30

Asn Thr Thr  
             35

<210> 7  
 <211> 31  
 <212> PRT  
 <213> Clostridium difficile

<400> 7  
 Met Ile Lys Lys Ile Ser Thr Ile Leu Ser Leu Val Leu Leu Ile Ser  
     1                    5                    10                    15  
 Ile Ser Ser Thr Ile Gly Val Phe Ala Asp Ala Asn Pro Lys Arg  
             20                    25                    30

<210> 8  
 <211> 34  
 <212> PRT  
 <213> Clostridium difficile

<400> 8  
 Met Leu Ser Asn Lys Lys Arg Ser Met Ala Ile Val Met Ala Gly Ala  
     1                    5                    10                    15  
 Thr Val Met Ser Ala Ala Ala Pro Ile Phe Ala Asp Asn Thr Val Thr  
                     20                    25                    30

Glu Asn

<210> 9  
 <211> 41  
 <212> PRT  
 <213> Clostridium difficile

<400> 9  
 Met Lys Ser Thr Leu Gly Val Glu Asn Asn Met Lys Asn Ser Lys Lys  
     1                    5                    10                    15  
 Ile Leu Ala Ile Gly Leu Thr Leu Phe Leu Val Met Val Asn Thr Pro  
                     20                    25                    30

Met Val Ser Ala Leu Thr Ser Val Glu  
                     35                    40

<210> 10  
 <211> 34  
 <212> PRT  
 <213> Clostridium difficile

<400> 10  
 Met Asn Lys Arg Lys Ser Phe Ile Arg Thr Ile Ala Val Ser Thr Met  
     1                    5                    10                    15  
 Ala Val Ala Val Thr Gly Ser Ala Thr Cys Ala Tyr Ala Ala Pro Val  
                     20                    25                    30

Leu Gln

<210> 11  
 <211> 47  
 <212> PRT  
 <213> Clostridium difficile

<400> 11  
 Met Glu Asn Asn His Asn Ile Asn Ile Lys Tyr Lys Asn His Gln Gly  
     1                    5                    10                    15  
 Asp Met Lys Met Asn Lys Lys Ile Leu Ser Leu Gly Leu Ala Val Ser  
                     20                    25                    30

Leu Ile Leu Val Asn Phe Lys Ser Val Asn Ala Ser Ser Val Val  
           35                          40                          45

<210> 12  
 <211> 32  
 <212> PRT  
 <213> Clostridium difficile

<400> 12  
 Met Lys Val Asn Lys Arg Val Leu Ser Ile Gly Leu Ala Ile Ser Leu  
       1                          5                          10                          15

Ile Met Ala Gly Ala Pro Asn Ile Asn Ala Leu Ser Ser Ile Glu Lys  
                   20                          25                          30

<210> 13  
 <211> 50  
 <212> DNA  
 <213> Clostridium difficile

<400> 13  
 tagtttatta cattttaaaa tttagggtat aaaaacttgt aaacttggag 50

<210> 14  
 <211> 50  
 <212> DNA  
 <213> Clostridium difficile

<400> 14  
 cattttaaaa tttagggtat aaaaacttgt aaacttggag aaaataataa 50

<210> 15  
 <211> 50  
 <212> DNA  
 <213> Clostridium difficile

<400> 15  
 aacttgtaaa cttggagaaa ataataattt aaaaaaatag cttgcaaaaa 50

<210> 16  
 <211> 50  
 <212> DNA  
 <213> Clostridium difficile

<400> 16  
 aacttggaga aaataataat ttaaaaaaat agcttgcaaa aagaataaaa 50

<210> 17  
 <211> 50  
 <212> DNA  
 <213> Clostridium difficile

<400> 17  
 taatttaaaa aaatagcttg caaaaagaat aaaaatggat tattatagag 50

<210> 18  
 <211> 50  
 <212> DNA  
 <213> Clostridium difficile

<400> 18  
 aatagcttgc aaaaagaata aaaatggatt attatagaga tgtgagaaat 50

<210> 19  
 <211> 50  
 <212> DNA  
 <213> Clostridium difficile

<400> 19  
 tattatagag atgtgagaaa tattaggaat atatggatga ttattctatg 50

<210> 20  
 <211> 50  
 <212> DNA  
 <213> Clostridium difficile

<400> 20  
 gatgtgagaa atattaggaa tatatggatg attattctat gtacataata 50

<210> 21  
 <211> 50  
 <212> DNA  
 <213> Clostridium difficile

<400> 21  
 gattattcta tgtacataat aaagagatgt aattttaata taatgttggg 50

<210> 22  
 <211> 719  
 <212> PRT  
 <213> Clostridium difficile

<400> 22  
 Met Asn Lys Lys Asn Ile Ala Ile Ala Met Ser Gly Leu Thr Val Leu  
 1 5 10 15  
 Ala Ser Ala Ala Pro Val Phe Ala Ala Thr Thr Gly Thr Gln Gly Tyr  
 20 25 30  
 Thr Val Val Lys Asn Asp Trp Lys Lys Ala Val Lys Gln Leu Gln Asp  
 35 40 45  
 Gly Leu Lys Asp Asn Ser Ile Gly Lys Ile Thr Val Ser Phe Asn Asp  
 50 55 60  
 Gly Val Val Gly Glu Val Ala Pro Lys Ser Ala Asn Lys Lys Ala Asp  
 65 70 75 80  
 Arg Asp Ala Ala Ala Glu Lys Leu Tyr Asn Leu Val Asn Thr Gln Leu  
 85 90 95



Asp Lys Leu Gly Asp Gly Asp Tyr Val Asp Phe Ser Val Asp Tyr Asn  
 100 105 110  
 Leu Glu Asn Lys Ile Ile Thr Asn Gln Ala Asp Ala Glu Ala Ile Val  
 115 120 125  
 Thr Lys Leu Asn Ser Leu Asn Glu Lys Thr Leu Ile Asp Ile Ala Thr  
 130 135 140  
 Lys Asp Thr Phe Gly Met Val Ser Lys Thr Gln Asp Ser Glu Gly Lys  
 145 150 155 160  
 Asn Val Ala Ala Thr Lys Ala Leu Lys Val Lys Asp Val Ala Thr Phe  
 165 170 175  
 Gly Leu Lys Ser Gly Gly Ser Glu Asp Thr Gly Tyr Val Val Glu Met  
 180 185 190  
 Lys Ala Gly Ala Val Glu Asp Lys Tyr Gly Lys Val Gly Asp Ser Thr  
 195 200 205  
 Ala Gly Ile Ala Ile Asn Leu Pro Ser Thr Gly Leu Glu Tyr Ala Gly  
 210 215 220  
 Lys Gly Thr Thr Ile Asp Phe Asn Lys Thr Leu Lys Val Asp Val Thr  
 225 230 235 240  
 Gly Gly Ser Thr Pro Ser Ala Val Ala Val Ser Gly Phe Val Thr Lys  
 245 250 255  
 Asp Asp Thr Asp Leu Ala Lys Ser Gly Thr Ile Asn Val Arg Val Ile  
 260 265 270  
 Asn Ala Lys Glu Glu Ser Ile Asp Ile Asp Ala Ser Ser Tyr Thr Ser  
 275 280 285  
 Ala Glu Asn Leu Ala Lys Arg Tyr Val Phe Asp Pro Asp Glu Ile Ser  
 290 295 300  
 Glu Ala Tyr Lys Ala Ile Val Ala Leu Gln Asn Asp Gly Ile Glu Ser  
 305 310 315 320  
 Asn Leu Val Gln Leu Val Asn Gly Lys Tyr Gln Val Ile Phe Tyr Pro  
 325 330 335  
 Glu Gly Lys Arg Leu Glu Thr Lys Ser Ala Asn Asp Thr Ile Ala Ser  
 340 345 350  
 Gln Asp Thr Pro Ala Lys Val Val Ile Lys Ala Asn Lys Leu Lys Asp  
 355 360 365  
 Leu Lys Asp Tyr Val Asp Asp Leu Lys Thr Tyr Asn Asn Thr Tyr Ser  
 370 375 380  
 Asn Val Val Thr Val Ala Gly Glu Asp Arg Ile Glu Thr Ala Ile Glu  
 385 390 395 400  
 Leu Ser Ser Lys Tyr Tyr Asn Ser Asp Asp Lys Asn Ala Ile Thr Asp  
 405 410 415

Lys Ala Val Asn Asp Ile Val Leu Val Gly Ser Thr Ser Ile Val Asp  
 420 425 430  
 Gly Leu Val Ala Ser Pro Leu Ala Ser Glu Lys Thr Ala Pro Leu Leu  
 435 440 445  
 Leu Thr Ser Lys Asp Lys Leu Asp Ser Ser Val Lys Ser Glu Ile Lys  
 450 455 460  
 Arg Val Met Asn Leu Lys Ser Asp Thr Gly Ile Asn Thr Ser Lys Lys  
 465 470 475 480  
 Val Tyr Leu Ala Gly Gly Val Asn Ser Ile Ser Lys Asp Val Glu Asn  
 485 490 495  
 Glu Leu Lys Asn Met Gly Leu Lys Val Thr Arg Leu Ser Gly Glu Asp  
 500 505 510  
 Arg Tyr Glu Thr Ser Leu Ala Ile Ala Asp Glu Ile Gly Leu Asp Asn  
 515 520 525  
 Asp Lys Ala Phe Val Val Gly Gly Thr Gly Leu Ala Asp Ala Met Ser  
 530 535 540  
 Ile Ala Pro Val Ala Ser Gln Leu Lys Asp Gly Asp Ala Thr Pro Ile  
 545 550 555 560  
 Val Val Val Asp Gly Lys Ala Lys Glu Ile Ser Asp Asp Ala Lys Ser  
 565 570 575  
 Phe Leu Gly Thr Ser Asp Val Asp Ile Ile Gly Gly Lys Asn Ser Val  
 580 585 590  
 Ser Lys Glu Ile Glu Glu Ser Ile Asp Ser Ala Thr Gly Lys Thr Pro  
 595 600 605  
 Asp Arg Ile Ser Gly Asp Asp Arg Gln Ala Thr Asn Ala Glu Val Leu  
 610 615 620  
 Lys Glu Asp Asp Tyr Phe Thr Asp Gly Glu Val Val Asn Tyr Phe Val  
 625 630 635 640  
 Ala Lys Asp Gly Ser Thr Lys Glu Asp Gln Leu Val Asp Ala Leu Ala  
 645 650 655  
 Ala Ala Pro Ile Ala Gly Arg Phe Lys Glu Ser Pro Ala Pro Ile Ile  
 660 665 670  
 Leu Ala Thr Asp Thr Leu Ser Ser Asp Gln Asn Val Ala Val Ser Lys  
 675 680 685  
 Ala Val Pro Lys Asp Gly Gly Thr Asn Leu Val Gln Val Gly Lys Gly  
 690 695 700  
 Ile Ala Ser Ser Val Ile Asn Lys Met Lys Asp Leu Leu Asp Met  
 705 710 715

<210> 23  
 <211> 623  
 <212> PRT

&lt;213&gt; Clostridium difficile

&lt;400&gt; 23

Met Asn Lys Lys Asn Leu Ser Val Ile Met Ala Ala Ala Met Ile Ser  
 1 5 10 15  
 Thr Ser Val Ala Pro Val Phe Ala Ala Glu Thr Thr Gln Val Lys Lys  
 20 25 30  
 Glu Thr Ile Thr Lys Lys Glu Ala Thr Glu Leu Val Ser Lys Val Arg  
 35 40 45  
 Asp Leu Met Ser Gln Lys Tyr Thr Gly Gly Ser Gln Val Gly Gln Pro  
 50 55 60  
 Ile Tyr Glu Ile Lys Val Gly Glu Thr Leu Ser Lys Leu Lys Ile Ile  
 65 70 75 80  
 Thr Asn Ile Asp Glu Leu Glu Lys Leu Val Asn Ala Leu Gly Glu Asn  
 85 90 95  
 Lys Glu Leu Ile Val Thr Ile Thr Asp Lys Gly His Ile Thr Asn Ser  
 100 105 110  
 Ala Asn Glu Val Val Ala Glu Ala Thr Glu Lys Tyr Glu Asn Ser Ala  
 115 120 125  
 Asp Leu Ser Ala Glu Ala Asn Ser Ile Thr Glu Lys Ala Lys Thr Glu  
 130 135 140  
 Thr Asn Gly Ile Tyr Lys Val Ala Asp Val Lys Ala Ser Tyr Asp Ser  
 145 150 155 160  
 Ala Lys Asp Lys Leu Val Ile Thr Leu Arg Asp Lys Thr Glu Thr Val  
 165 170 175  
 Thr Ser Lys Thr Ile Tyr Val Gly Ile Gly Asp Glu Lys Val Asp Leu  
 180 185 190  
 Thr Ala Asn Pro Val Asp Ser Thr Gly Thr Asn Leu Asp Pro Ser Ala  
 195 200 205  
 Glu Gly Phe Arg Val Asn Lys Ile Asp Lys Leu Gly Val Ala Gly Ala  
 210 215 220  
 Lys Asn Ile Asp Asp Val Gln Leu Ala Glu Ile Thr Ile Lys Asn Ser  
 225 230 235 240  
 Asp Leu Asn Thr Val Ser Pro Gln Asp Leu Tyr Asp Gly Tyr Arg Leu  
 245 250 255  
 Thr Val Lys Gly Asn Met Val Ala Asn Gly Thr Ser Lys Ser Ile Ser  
 260 265 270  
 Asp Ile Ser Ala Lys Asp Ser Glu Thr Gly Lys Tyr Lys Phe Thr Ile  
 275 280 285  
 Lys Tyr Thr Asp Ala Ser Gly Lys Ala Thr Glu Leu Thr Val Glu Ser  
 290 295 300

Thr Asn Glu Lys Asp Leu Lys Asp Ala Lys Ala Ala Leu Glu Gly Asn  
 305 310 315 320  
 Ser Lys Val Lys Leu Ile Ala Gly Asp Asp Arg Tyr Ala Thr Ala Val  
 325 330 335  
 Ala Ile Ala Lys Gln Thr Lys Tyr Thr Asp Asn Ile Val Ile Val Asn  
 340 345 350  
 Ser Asn Lys Leu Val Asp Gly Leu Ala Ala Thr Pro Leu Ala Gln Ser  
 355 360 365  
 Lys Lys Ala Pro Ile Leu Leu Ala Ser Asp Asn Glu Ile Pro Lys Val  
 370 375 380  
 Thr Leu Asp Tyr Ile Lys Asp Ile Ile Lys Lys Ser Pro Ser Ala Lys  
 385 390 395 400  
 Ile Tyr Ile Val Gly Gly Glu Ser Ala Val Ser Asn Thr Ala Lys Lys  
 405 410 415  
 Gln Leu Glu Ser Val Thr Lys Asn Ile Glu Arg Leu Ala Gly Asp Asp  
 420 425 430  
 Arg His Thr Thr Ser Val Ala Val Ala Lys Ala Met Gly Ser Phe Lys  
 435 440 445  
 Asp Ala Phe Val Val Gly Ala Lys Gly Glu Ala Asp Ala Met Ser Ile  
 450 455 460  
 Ala Ala Lys Ala Ala Glu Leu Lys Ala Pro Ile Ile Val Asn Gly Trp  
 465 470 475 480  
 Asn Asp Leu Ser Ala Asp Ala Ile Lys Leu Met Asp Gly Lys Glu Ile  
 485 490 495  
 Gly Ile Val Gly Gly Ser Asn Asn Val Ser Ser Gln Ile Glu Asn Gln  
 500 505 510  
 Leu Ala Asp Ile Asp Lys Asp Arg Lys Val Gln Arg Val Glu Gly Glu  
 515 520 525  
 Thr Arg His Asp Thr Asn Ala Lys Val Ile Glu Thr Tyr Tyr Gly Lys  
 530 535 540  
 Leu Asp Lys Leu Tyr Ile Ala Lys Asp Gly Tyr Gly Asn Asn Gly Met  
 545 550 555 560  
 Leu Val Asp Ala Leu Ala Ala Gly Pro Leu Ala Ala Gly Lys Gly Pro  
 565 570 575  
 Ile Leu Leu Ala Lys Thr Asp Ile Thr Asp Ser Gln Lys Asn Ala Leu  
 580 585 590  
 Ser Lys Lys Leu Asn Leu Gly Ala Glu Val Thr Gln Ile Gly Asn Gly  
 595 600 605  
 Val Glu Leu Thr Val Ile Gln Lys Ile Ala Lys Ile Leu Gly Trp  
 610 615 620

&lt;211&gt; 610

&lt;212&gt; PRT

&lt;213&gt; Clostridium difficile

&lt;400&gt; 24

```

Met Lys Ile Ser Lys Lys Ile Val Ser Leu Leu Thr Met Thr Phe Leu
 1           5           10           15

Thr Val Thr Leu Tyr Gly Asn Thr Ser Asn Ala Ser Thr Lys Asp Thr
          20           25           30

Leu Thr Gly Ser Gly Arg Trp Glu Thr Ala Ile Lys Ile Ser Gln Ala
          35           40           45

Gly Trp Thr Lys Ser Glu Ser Ala Val Leu Val Asn Asp Asn Ser Ile
          50           55           60

Ala Asp Ala Leu Ser Ala Thr Pro Phe Ala Lys Ala Lys Asp Ala Pro
          65           70           75           80

Ile Leu Leu Thr Gln Ser Asn Lys Leu Asp Ser Arg Thr Lys Ala Glu
          85           90           95

Leu Lys Arg Leu Gly Val Lys Asn Val Tyr Leu Ile Gly Gly Ser Ile
          100          105          110

Ala Leu Ser Ser Glu Ile Glu Lys Gln Leu Asn Ala Glu Asn Ile Asn
          115          120          125

Phe Glu Arg Ile Ser Gly Asn Ser Arg Tyr Asp Thr Ser Leu Lys Leu
          130          135          140

Ala Glu Lys Leu Asp Arg Glu Lys Ser Ile Ser Lys Ile Val Val Val
          145          150          155          160

Asn Gly Glu Lys Gly Leu Ala Asp Ala Val Ser Val Gly Ala Ile Ala
          165          170          175

Ala Gln Glu Asn Met Pro Ile Ile Leu Ser Asp Ser Glu Asn Gly Thr
          180          185          190

Glu Val Ala Asp Asn Phe Ile Asp Ser Lys Asp Ile Ala Lys Ser Tyr
          195          200          205

Val Ile Gly Gly Thr Tyr Ser Ile Ser Asn Ser Val Glu Arg Ser Leu
          210          215          220

Pro Asn Ala Thr Arg Ile Ala Gly Ser Ser Arg Ser Glu Thr Asn Ala
          225          230          235          240

Lys Ile Ile Glu Glu Phe Tyr Lys Asp Thr Asp Ile Lys Asn Ile Tyr
          245          250          255

Val Thr Lys Asp Gly Thr Lys Asn Lys Asn Asp Leu Ile Asp Ser Leu
          260          265          270

Ala Val Gly Val Leu Ala Ala Lys Asn Ser Ser Pro Ile Val Leu Ala
          275          280          285

Gly Asn Lys Leu Asp Thr Thr Gln Lys Asp Val Leu Asn Thr Lys Ile
          290          295          300

```

Ile Asp Lys Val Thr Gln Ile Gly Gly Leu Gly Asn Glu Asn Val Val  
 305 310 315 320  
 Glu Asp Ile Leu Asp Ile Gln Glu Glu Thr Lys Tyr Thr Val Glu Thr  
 325 330 335  
 Ile Asp Glu Leu Asn Ala Ala Ile Lys Arg Ala Asp Ala Asn Asp Ile  
 340 345 350  
 Ile Lys Phe Lys Pro Glu Lys Glu Lys Thr Ile Asn Asn Ser Phe Ser  
 355 360 365  
 Ile Glu Thr Lys Lys Thr Val Thr Ile Glu Leu Asp Gly Arg Tyr Arg  
 370 375 380  
 Gln Thr Ile Thr Leu Asp Ile Pro Asn Gly Lys Phe Asn Asn Tyr Ala  
 385 390 395 400  
 Glu Ile Glu Gly Gly Val Lys Leu Lys Asn Ile Lys Asn Glu Ser Leu  
 405 410 415  
 Val Asn Lys Gly Ser Ile Gln Asp Leu Asp Ile Tyr Asp Glu Asn Gly  
 420 425 430  
 Cys Lys Ile Glu Asn Glu Ser Ser Gly Glu Ile Trp Phe Val Thr Ile  
 435 440 445  
 Val Glu Glu Ala Asn Asp Val Tyr Ile Val Asn Ser Gly Asp Ile Thr  
 450 455 460  
 Lys Ile Ser Asn Asn Ser Ser Ser Thr Ile Ile Arg Asn Ser Gly Asn  
 465 470 475 480  
 Ile Asp Thr Val Thr Gly Lys Lys Glu Pro Ala Ile Ser Gly Asn Lys  
 485 490 495  
 Pro Lys Val Asn Asp Thr Glu Lys Glu Thr Lys Ala Ala Arg Gly Leu  
 500 505 510  
 Asn Pro Arg Val Glu Ala Cys Ser Val Pro Lys Lys Asp Tyr Val Met  
 515 520 525  
 Ile Thr Ile Pro Asn Ser Pro Lys Asp Ser Arg Tyr Lys Ile Tyr Tyr  
 530 535 540  
 Arg Val Val Tyr Asn Lys Pro Tyr Ala Met Asp Val Gly Asp Lys Ile  
 545 550 555 560  
 Asn Ile Gly Glu Trp Thr Val Ala Pro Thr Asp Glu Glu Pro Phe Leu  
 565 570 575  
 Glu Lys Ala Lys Asn Gly Cys Tyr Val Glu Ala Val Glu Val Asn Thr  
 580 585 590  
 Ser Thr Lys Glu Val Ser Arg Trp Gly Arg Thr Asn Ala Thr Asp Asp  
 595 600 605  
 Gly Phe  
 610

&lt;210&gt; 25

&lt;211&gt; 803

&lt;212&gt; PRT

&lt;213&gt; Clostridium difficile

&lt;400&gt; 25

Met Arg Lys Tyr Lys Ser Lys Lys Leu Ser Lys Leu Leu Ala Leu Ser  
 1 5 10 15

Thr Val Cys Phe Leu Ile Val Ser Thr Ile Pro Val Ser Ala Glu Asn  
 20 25 30

His Lys Thr Leu Asp Gly Val Glu Thr Ala Glu Tyr Ser Glu Ser Tyr  
 35 40 45

Leu Gln Tyr Leu Glu Asp Val Lys Asn Gly Asp Thr Ala Lys Tyr Asn  
 50 55 60

Gly Val Ile Pro Phe Pro His Glu Met Glu Gly Thr Thr Leu Arg Asn  
 65 70 75 80

Lys Gly Arg Ser Ser Leu Pro Ser Ala Tyr Lys Ser Ser Val Ala Tyr  
 85 90 95

Asn Pro Met Asp Leu Gly Leu Thr Thr Pro Ala Lys Asn Gln Gly Ser  
 100 105 110

Leu Asn Thr Cys Trp Ser Phe Ser Gly Met Ser Thr Leu Glu Ala Tyr  
 115 120 125

Leu Lys Leu Lys Gly Tyr Gly Thr Tyr Asp Leu Ser Glu Glu His Leu  
 130 135 140

Arg Trp Trp Ala Thr Gly Gly Lys Tyr Gly Trp Asn Leu Asp Asp Met  
 145 150 155 160

Ser Gly Ser Ser Asn Val Thr Ala Ile Gly Tyr Leu Thr Ala Trp Ala  
 165 170 175

Gly Pro Lys Leu Glu Lys Asp Ile Pro Tyr Asn Leu Lys Ser Glu Ala  
 180 185 190

Gln Gly Ala Thr Lys Pro Ser Asn Met Asp Thr Ala Pro Thr Gln Phe  
 195 200 205

Asn Val Thr Asp Val Val Arg Leu Asn Lys Asp Lys Glu Thr Val Lys  
 210 215 220

Asn Ala Ile Met Gln Tyr Gly Ser Val Thr Ser Gly Tyr Ala His Tyr  
 225 230 235 240

Ser Thr Tyr Phe Asn Lys Asp Glu Thr Ala Tyr Asn Cys Thr Asn Lys  
 245 250 255

Arg Ala Pro Leu Asn His Ala Val Ala Ile Val Gly Trp Asp Asp Asn  
 260 265 270

Tyr Ser Lys Asp Asn Phe Ala Ser Asp Val Lys Pro Glu Ser Asn Gly  
 275 280 285

Ala Trp Leu Val Lys Ser Ser Trp Gly Glu Phe Asn Ser Met Lys Gly  
 290 295 300  
 Phe Phe Trp Ile Ser Tyr Glu Asp Lys Thr Leu Leu Thr Asp Thr Asp  
 305 310 315 320  
 Asn Tyr Ala Met Lys Ser Val Ser Lys Pro Asp Ser Asp Lys Lys Met  
 325 330 335  
 Tyr Gln Leu Glu Tyr Ala Gly Leu Ser Lys Ile Met Ser Asn Lys Val  
 340 345 350  
 Thr Ala Ala Asn Val Phe Asp Phe Ser Arg Asp Ser Glu Lys Leu Asp  
 355 360 365  
 Ser Val Met Phe Glu Thr Asp Ser Val Gly Ala Lys Tyr Glu Val Tyr  
 370 375 380  
 Tyr Ala Pro Val Val Asn Gly Val Pro Gln Asn Asn Ser Met Thr Lys  
 385 390 395 400  
 Leu Ala Ser Gly Thr Val Ser Tyr Ser Gly Tyr Ile Asn Val Pro Thr  
 405 410 415  
 Asn Ser Tyr Ser Leu Pro Lys Gly Lys Gly Ala Ile Val Val Val Ile  
 420 425 430  
 Asp Asn Thr Ala Asn Pro Asn Arg Glu Lys Ser Thr Leu Ala Tyr Glu  
 435 440 445  
 Thr Asn Ile Asp Ala Tyr Tyr Leu Tyr Glu Ala Lys Ala Asn Leu Gly  
 450 455 460  
 Glu Ser Tyr Ile Leu Gln Asn Asn Lys Phe Glu Asp Ile Asn Thr Tyr  
 465 470 475 480  
 Ser Glu Phe Ser Pro Cys Asn Phe Val Ile Lys Ala Ile Thr Lys Thr  
 485 490 495  
 Ser Ser Gly Gln Ala Thr Ser Gly Glu Ser Leu Thr Gly Ala Asp Arg  
 500 505 510  
 Tyr Glu Thr Ala Val Lys Val Ser Gln Lys Gly Trp Thr Ser Ser Gln  
 515 520 525  
 Asn Ala Val Leu Val Asn Gly Asp Ala Ile Val Asp Ala Leu Thr Ala  
 530 535 540  
 Thr Pro Phe Thr Ala Ala Ile Asp Ser Pro Ile Leu Leu Thr Gly Lys  
 545 550 555 560  
 Asp Asn Leu Asp Ser Lys Thr Lys Ala Glu Leu Gln Arg Leu Gly Thr  
 565 570 575  
 Lys Lys Val Tyr Leu Ile Gly Gly Glu Asn Ser Leu Ser Lys Asn Val  
 580 585 590  
 Gln Thr Gln Leu Ser Asn Met Gly Ile Ser Val Glu Arg Ile Ser Gly  
 595 600 605



Ser Asp Arg Tyr Lys Thr Ser Ile Ser Leu Ala Gln Lys Leu Asn Ser  
 610 615 620  
 Ile Lys Ser Val Ser Gln Val Ala Val Ala Asn Gly Val Asn Gly Leu  
 625 630 635 640  
 Ala Asp Ala Ile Ser Val Gly Ala Ala Ala Asp Asn Asn Met Pro  
 645 650 655  
 Ile Ile Leu Thr Asn Glu Lys Ser Glu Leu Gln Gly Ala Asp Glu Phe  
 660 665 670  
 Leu Asn Ser Ser Lys Ile Thr Lys Ser Tyr Ile Ile Gly Gly Thr Ala  
 675 680 685  
 Thr Leu Ser Ser Asn Leu Glu Ser Lys Leu Ser Asn Pro Thr Arg Leu  
 690 695 700  
 Ala Gly Ser Asn Arg Asn Glu Thr Asn Ala Lys Ile Ile Asp Lys Phe  
 705 710 715 720  
 Tyr Pro Ser Ser Asp Leu Lys Tyr Ala Phe Val Val Lys Asp Gly Ser  
 725 730 735  
 Lys Ser Gln Gly Asp Leu Ile Asp Gly Leu Ala Val Gly Ala Leu Gly  
 740 745 750  
 Ala Lys Thr Asp Ser Pro Val Val Leu Val Gly Asn Lys Leu Asp Glu  
 755 760 765  
 Ser Gln Lys Asn Val Leu Lys Ser Lys Lys Ile Glu Thr Pro Ile Arg  
 770 775 780  
 Val Gly Gly Asn Gly Asn Glu Ser Ala Phe Asn Glu Leu Asn Thr Leu  
 785 790 795 800  
 Leu Gly Lys

<210> 26  
 <211> 525  
 <212> PRT  
 <213> Clostridium difficile

<400> 26  
 Met Lys Ala Pro Lys Thr Ile Leu Thr Ile Leu Thr Ile Ala Leu Thr  
 1 5 10 15  
 Leu Ser Ser Ile Ser Ile Ile Pro Ser Tyr Ala Leu Thr Glu Glu Lys  
 20 25 30  
 Leu Ile Gly Asn Gly Arg Tyr Glu Thr Ala Val Lys Ile Ser Gln Lys  
 35 40 45  
 Ala Tyr Ser Ser Ser Pro Asn Val Val Leu Val Asn Asp Asn Ser Leu  
 50 55 60  
 Ala Asp Ala Leu Ser Ala Thr Pro Phe Ala Lys Ala Lys Gly Ala Pro  
 65 70 75 80

Ile Leu Leu Thr Glu Ser Asp Lys Leu Asp Asp Arg Thr Glu Lys Glu  
 85 90 95  
 Ile Lys Arg Leu Gly Ala Lys Asp Ile Tyr Leu Ile Gly Gly Thr Ala  
 100 105 110  
 Val Leu Asn Lys Asp Ile Glu Asn Lys Leu Lys Gly Asn Gly Leu Asn  
 115 120 125  
 Val Glu Arg Ile Asn Gly Lys Asn Arg Tyr Glu Thr Ser Leu Ile Leu  
 130 135 140  
 Ala Asn Lys Leu Lys Asp Ile Lys Asp Ile Lys Glu Val Ala Val Val  
 145 150 155 160  
 Asn Gly Glu Lys Gly Leu Ser Asp Ala Val Ser Val Gly Ala Pro Ala  
 165 170 175  
 Ala Gln Asn Lys Met Pro Ile Ile Leu Ser Asn Pro Lys Asp Gly Val  
 180 185 190  
 Glu Ala Phe Asp Lys Phe Ile Arg Asp Glu Lys Val Ile Lys Ala Tyr  
 195 200 205  
 Val Ile Gly Gly Thr Asn Ser Val Ser Arg Ala Val Glu Lys Ser Leu  
 210 215 220  
 Pro Asn Ala Glu Arg Val Ser Gly Lys Asp Arg Asn Glu Thr Asn Ala  
 225 230 235 240  
 Lys Val Ile Glu Lys Phe Tyr Thr Asp Thr Asn Leu Ser Asn Leu Tyr  
 245 250 255  
 Val Thr Lys Asp Gly Ser Lys Asn Glu Asn Gln Leu Ile Asp Ser Leu  
 260 265 270  
 Ala Val Gly Val Leu Ala Ala Lys Asn Glu Ser Pro Ile Val Leu Val  
 275 280 285  
 Gly Asn Lys Leu Asn Thr Lys Gln Arg Asp Ile Leu Ser Thr Lys Lys  
 290 295 300  
 Leu Asn Thr Ile Thr Gln Val Gly Gly Asn Gly Asn Glu Glu Ala Phe  
 305 310 315 320  
 Asp Glu Ile Lys Ser Leu Gln Glu Lys Thr Val Phe Glu Ala Lys Thr  
 325 330 335  
 Val Glu Glu Leu Thr Asp Met Ile Asn Ile Ala Ser Pro Asn Asp Ile  
 340 345 350  
 Ile Asn Phe Lys Pro Lys Glu Asn Thr Val Asn Glu Ala Phe Arg Met  
 355 360 365  
 Val Thr Asn Lys Pro Ile Thr Val Asn Ile Lys Gly Asp Cys Ser Lys  
 370 375 380  
 Thr Leu Thr Val Asp Met Pro Asn Gly Glu Val Asn Asn Tyr Ala Thr  
 385 390 395 400

<400> 27																
Met	Arg	Gly	Asp	Met	Met	Lys	Lys	Thr	Thr	Lys	Leu	Leu	Ala	Thr	Gly	
1				5					10					15		
Met	Leu	Ser	Val	Ala	Met	Val	Ala	Pro	Asn	Val	Ala	Leu	Ala	Ala	Glu	
			20					25					30			
Asn	Thr	Thr	Ala	Asn	Thr	Glu	Ser	Asn	Ser	Asp	Ile	Asn	Ile	Asn	Leu	
		35					40					45				
Gln	Arg	Lys	Ser	Val	Val	Leu	Gly	Ser	Lys	Ser	Asn	Ala	Ser	Val	Lys	
	50					55					60					
Phe	Lys	Glu	Lys	Leu	Asn	Ala	Asp	Ser	Ile	Thr	Leu	Asn	Phe	Met	Cys	
65					70					75					80	
Tyr	Asp	Met	Pro	Leu	Glu	Ala	Thr	Leu	Asn	Tyr	Asn	Glu	Lys	Thr	Asp	
				85					90					95		
Ser	Tyr	Glu	Gly	Val	Ile	Asn	Tyr	Asn	Lys	Asp	Pro	Glu	Tyr	Leu	Asn	
		100						105					110			
Val	Trp	Glu	Leu	Gln	Ser	Ile	Lys	Ile	Asn	Gly	Lys	Asp	Glu	Gln	Lys	
		115					120					125				
Val	Leu	Asn	Lys	Glu	Asp	Leu	Glu	Ser	Met	Gly	Leu	Asn	Leu	Lys	Asp	
	130					135					140					
Tyr	Asp	Val	Thr	Gln	Glu	Phe	Ile	Ile	Ser	Asp	Ala	Asn	Ser	Thr	Lys	
145					150					155					160	

Ala Val Asn Glu Tyr Met Arg Lys Thr Ser Ala Pro Val Lys Lys Leu  
 165 170 175  
 Ala Gly Ala Thr Arg Phe Glu Thr Ala Val Glu Ile Ser Lys Gln Gly  
 180 185 190  
 Trp Lys Asp Gly Ser Ser Lys Val Val Ile Val Asn Gly Glu Leu Ala  
 195 200 205  
 Ala Asp Gly Ile Thr Ala Thr Pro Leu Ala Ser Thr Tyr Asp Ala Pro  
 210 215 220  
 Ile Leu Leu Ala Asn Lys Asp Asp Ile Pro Glu Ser Thr Lys Ala Glu  
 225 230 235 240  
 Leu Lys Arg Leu Asn Pro Ser Asp Val Ile Ile Ile Gly Asp Asp Gly  
 245 250 255  
 Ser Val Ser Gln Lys Ala Val Ser Gln Ile Lys Ser Ala Val Asn Val  
 260 265 270  
 Asn Val Thr Arg Ile Gly Gly Val Asp Arg His Glu Thr Ser Leu Leu  
 275 280 285  
 Ile Ala Lys Glu Ile Asp Lys Tyr His Asp Val Asn Lys Ile Tyr Ile  
 290 295 300  
 Ala Asn Gly Tyr Ala Gly Glu Tyr Asp Ala Leu Asn Ile Ser Ser Lys  
 305 310 315 320  
 Ala Gly Glu Asp Gln Gln Pro Ile Ile Leu Ala Asn Lys Asp Ser Val  
 325 330 335  
 Pro Gln Gly Thr Tyr Asn Trp Leu Ser Ser Gln Gly Leu Glu Glu Ala  
 340 345 350  
 Tyr Tyr Ile Gly Gly Ser Gln Ser Leu Ser Ser Lys Ile Ile Asp Gln  
 355 360 365  
 Ile Ser Lys Ile Ala Lys Asn Gly Thr Ser Lys Asn Arg Val Ser Gly  
 370 375 380  
 Ala Asp Arg His Glu Thr Asn Ala Asn Val Ile Lys Thr Phe Tyr Pro  
 385 390 395 400  
 Asp Lys Glu Leu Ser Ala Met Leu Val Ala Lys Ser Asp Ile Ile Val  
 405 410 415  
 Asp Ser Ile Thr Ala Gly Pro Leu Ala Ala Lys Leu Lys Ala Pro Ile  
 420 425 430  
 Leu Ile Thr Pro Lys Thr Tyr Val Ser Ala Tyr His Ser Thr Asn Leu  
 435 440 445  
 Ser Glu Lys Thr Ala Glu Thr Val Tyr Gln Ile Gly Asp Gly Met Lys  
 450 455 460  
 Asp Ser Val Ile Asn Ser Ile Ala Ser Ser Leu Ser Lys His Asn Ala  
 465 470 475 480

Pro Thr Glu Pro Asp Asn Ser Gly Ser Ala Ala Gly Lys Thr Val Val  
 485 490 495  
 Ile Asp Pro Gly His Gly Gly Ser Asp Ser Gly Ala Thr Ser Gly Leu  
 500 505 510  
 Asn Gly Gly Ala Gln Glu Lys Lys Tyr Thr Leu Asn Thr Ala Leu Ala  
 515 520 525  
 Thr Thr Glu Tyr Leu Arg Ser Lys Gly Ile Asn Val Val Met Thr Arg  
 530 535 540  
 Asp Thr Asp Lys Thr Met Ala Leu Gly Glu Arg Thr Ala Leu Ser Asn  
 545 550 555 560  
 Thr Ile Lys Pro Asp Leu Phe Thr Ser Ile His Tyr Asn Ala Ser Asn  
 565 570 575  
 Gly Ser Gly Asn Gly Val Glu Ile Tyr Tyr Lys Val Lys Asp Lys Asn  
 580 585 590  
 Gly Gly Thr Thr Lys Thr Ala Ala Ser Asn Ile Leu Lys Arg Ile Leu  
 595 600 605  
 Glu Lys Phe Asn Met Lys Asn Arg Gly Ile Lys Thr Arg Thr Leu Asp  
 610 615 620  
 Asn Gly Lys Asp Tyr Leu Tyr Val Leu Arg Asn Asn Asn Tyr Pro Ala  
 625 630 635 640  
 Ile Leu Val Glu Cys Ala Phe Ile Asp Asn Lys Ser Asp Met Asp Lys  
 645 650 655  
 Leu Asn Thr Ala Glu Lys Val Lys Thr Met Gly Thr Gln Ile Gly Ile  
 660 665 670  
 Gly Ile Glu Asp Thr Val Lys  
 675

<210> 28  
 <211> 351  
 <212> PRT  
 <213> Clostridium difficile

<400> 28  
 Met Ile Lys Lys Ile Ser Thr Ile Leu Ser Leu Val Leu Leu Ile Ser  
 1 5 10 15  
 Ile Ser Ser Thr Ile Gly Val Phe Ala Asp Ala Asn Pro Lys Arg Glu  
 20 25 30  
 Leu Ile Glu Gly Ser Ile Pro Glu Ile Ser Thr Glu Leu Asn Lys Arg  
 35 40 45  
 Ala Phe Lys Asp Ser Lys Glu Val Ile Leu Val Asn Glu Glu Ser Ile  
 50 55 60  
 Val Asp Ser Ile Ser Ala Thr Pro Leu Ala Tyr Ser Lys Asn Ala Pro  
 65 70 75 80

Ile Val Val Thr Lys Ser Lys Asn Leu Gly Arg Val Thr Arg Asn Tyr  
                     85                                    90                                    95  
 Leu Lys Glu Leu Gly Pro Glu Lys Val Thr Ile Val Gly Gly Leu Lys  
                     100                                    105                                    110  
 Ala Val Ser Lys Asp Ala Glu Arg Asn Ile Glu Lys Met Gly Met Lys  
                     115                                    120                                    125  
 Val Glu Arg Ile Arg Gly Lys Asp Arg Tyr Asp Thr Ser Leu Lys Ile  
                     130                                    135                                    140  
 Ala Arg Glu Met Tyr Arg Thr Val Gly Phe Asp Glu Ala Phe Leu Leu  
                     145                                    150                                    155                                    160  
 Ser Ser Thr Thr Gly Leu Glu Asn Ala Ile Ser Val Tyr Ser Tyr Ala  
                     165                                    170                                    175  
 Ala Lys Ser Gly Met Pro Ile Ile Trp Ala Lys Asp Glu Gly Phe Glu  
                     180                                    185                                    190  
 Glu Gln Ile Asp Phe Leu Lys Gly Lys Asn Leu Lys Lys Ile Tyr Ala  
                     195                                    200                                    205  
 Leu Gly Asp Ser Lys Glu Phe Ile Ala Glu Ile Asp Ser Asn Leu Lys  
                     210                                    215                                    220  
 Asn Ile Glu Gly Ile Lys Gln Ile Asn Lys Ser Ser Thr Asn Val Asp  
                     225                                    230                                    235                                    240  
 Leu Ile Lys Lys Phe Tyr Asp Glu Lys Asp Ile Lys Lys Ile Tyr Thr  
                     245                                    250                                    255  
 Ala Arg Leu Asp Phe Gly Ser Arg Ser Asp Val Asn Glu Tyr Ile Ser  
                     260                                    265                                    270  
 Leu Gly Val Val Ser Ala Lys Glu Asn Met Pro Ile Leu Ile Cys Ser  
                     275                                    280                                    285  
 Asp Asn Leu Ser Arg Ala Gln Asp Lys Phe Leu Lys Asp Ser Asn Ile  
                     290                                    295                                    300  
 Asn Asp Val Val Glu Val Gly Tyr Thr Val Gly Asp Tyr Ser Leu Phe  
                     305                                    310                                    315                                    320  
 Lys Ser Ile Phe Asn Leu Thr Phe Leu Ser Cys Ile Val Leu Ile Leu  
                     325                                    330                                    335  
 Leu Leu Leu Leu Ile Thr Phe Arg Ala Leu Arg Tyr Glu Ser Lys  
                     340                                    345                                    350

&lt;210&gt; 29

&lt;211&gt; 631

&lt;212&gt; PRT

&lt;213&gt; Clostridium difficile

&lt;400&gt; 29

Met Leu Ser Asn Lys Lys Arg Ser Met Ala Ile Val Met Ala Gly Ala  
   1                                    5                                    10                                    15

Thr Val Met Ser Ala Ala Ala Pro Ile Phe Ala Asp Asn Thr Val Thr  
 20 25 30  
 Glu Asn Val Asp Lys Asn Tyr Thr Val Ser Ala Lys Asp Ser Ala Lys  
 35 40 45  
 Leu Ile Glu Glu Val Arg Lys Ala Leu Glu Val Lys Phe Glu Asp Thr  
 50 55 60  
 Lys Ala Gly Ala Asn Val Asn Asp Arg Val Tyr Asp Ile Lys Val Asp  
 65 70 75 80  
 Asn Val Asn Leu Thr Asn Ala Thr Gln Leu Gln Asn Lys Ile Asn Ser  
 85 90 95  
 Leu Thr Glu Gly Gln Ser Leu Lys Val Thr Ile Gln Asp Lys Gly His  
 100 105 110  
 Gln Val Leu Gly Gly Lys Val Val Asp Tyr Lys Ile Glu Asn Tyr Lys  
 115 120 125  
 Thr Ala Gln Glu Ile Val Asp Ala Val Asn Ala Tyr Asn Ala Thr Leu  
 130 135 140  
 Ala Glu Asp Ser Asp Asn Lys Leu Thr Ala Thr Ile Lys Ser Thr Asn  
 145 150 155 160  
 Thr Val Glu Val Lys Arg Ala Lys Asp Ser Ala Asn Val Ile Thr Leu  
 165 170 175  
 Asn Val Gly Asp Gln His Leu Asp Phe Ser Lys Val Ile Thr Ser Glu  
 180 185 190  
 Glu Gly Thr Phe Glu Gly Tyr Glu Lys Arg Tyr Ser Asp Ile Asp Ser  
 195 200 205  
 Lys Glu Leu His Thr Val Thr Val Lys Asn Ala Asp Leu Gln Asp Ile  
 210 215 220  
 Ser Ala Glu Glu Leu Phe Asp Gly Ile Arg Leu Thr Thr Leu Gly Arg  
 225 230 235 240  
 Glu Ile Val Asn Lys Val Lys Asn Gly Tyr Ala Leu Thr Phe Glu Asn  
 245 250 255  
 Glu Ala Ile Leu Thr Gln Glu Gln Glu Asp Ser Asp Asp Lys Asp Lys  
 260 265 270  
 Pro Glu Lys Ser Ser Phe Asp Ile Val Leu Ser Lys Ala Asn Glu Lys  
 275 280 285  
 Pro Glu Thr Ile Ser Val Ser Ser Lys Asn His Lys Leu Val Arg Asp  
 290 295 300  
 Leu His Lys Val Leu Thr Asp Val Lys Asp Gly Lys Glu Leu Lys Val  
 305 310 315 320  
 Glu Val Leu Ser Gly Asp Ser Arg Phe Thr Thr Ala Val Glu Val Ser  
 325 330 335

Lys Glu Arg Phe Lys Asp Gly Glu Ala Glu Ala Ile Ile Leu Val Gly  
 340 345 350  
 Glu Asp Ala Ile Val Asp Gly Leu Ala Ser Ala Pro Leu Ala Ser Gln  
 355 360 365  
 Lys Asn Ala Pro Ile Leu Leu Ser Lys Lys Asp Ser Leu Pro Ser Glu  
 370 375 380  
 Ile Glu Ala Glu Ile Leu Arg Val Leu Gly Ser Asn Leu Ser Ser Lys  
 385 390 395 400  
 Lys Ile Tyr Ile Val Gly Gly Glu Ser Lys Val Ser Lys Glu Thr Glu  
 405 410 415  
 Glu Lys Leu Ser Lys Leu Gly Val Ser Lys Val Glu Arg Val Ser Gly  
 420 425 430  
 Glu Asp Arg Phe Glu Thr Ser Leu Glu Ile Ala Lys Gln Leu Lys Asp  
 435 440 445  
 Thr Phe Lys Thr Ala Phe Val Val Gly Gly Asn Gly Glu Ala Asp Ala  
 450 455 460  
 Met Ser Ile Ser Ala Arg Ala Ala Gln Phe Gly Ala Pro Ile Ile Val  
 465 470 475 480  
 Thr Gly Asn Glu Leu Asp Ala Asn Ala Glu Lys Leu Leu Lys Gly Lys  
 485 490 495  
 Glu Leu Glu Ile Val Gly Gly Glu Asn Ser Val Ser Lys Glu Val Glu  
 500 505 510  
 Asp Lys Leu Val Asp Ile Asp Leu Asn Asn Lys Val Glu Arg Leu Ala  
 515 520 525  
 Gly Glu Asn Arg Lys Asp Thr Asn Ala Lys Val Ile Asn Lys Tyr Tyr  
 530 535 540  
 Ala Gly Ala Thr Lys Ala Tyr Val Ala Lys Asp Gly Tyr Val Gly Gly  
 545 550 555 560  
 Asn Gly Gln Leu Val Asp Ala Leu Thr Ala Ala Pro Leu Ala Ala Ser  
 565 570 575  
 Ser Lys Ala Pro Ile Val Leu Thr Thr Glu Glu Leu Ser Lys Ser Gln  
 580 585 590  
 Glu Glu Val Val Glu Leu Arg Leu Lys Asn Ala Thr Lys Leu Val Gln  
 595 600 605  
 Ile Gly Glu Gly Ile Ala Lys Asn Ala Ile Glu Lys Ile Ala Glu Lys  
 610 615 620  
 Ile Asn Leu Phe Thr Lys Asn  
 625 630

&lt;210&gt; 30

&lt;211&gt; 477

&lt;212&gt; PRT



&lt;213&gt; Clostridium difficile

&lt;400&gt; 30

```

Met Lys Ser Thr Leu Gly Val Glu Asn Asn Met Lys Asn Ser Lys Lys
 1           5           10           15

Ile Leu Ala Ile Gly Leu Thr Leu Phe Leu Val Met Val Asn Thr Pro
 20           25           30

Met Val Ser Ala Leu Thr Ser Val Glu Gln Ile Lys Gly Asn Asp Arg
 35           40           45

Tyr Glu Thr Ala Ala Lys Ile Ala Asp Lys Gln Asn Tyr Asn Thr Ala
 50           55           60

Ile Leu Ile Asn Ser Asp Asn Ser Leu Ala Asp Gly Leu Ser Ala Ser
 65           70           75           80

Gly Leu Ala Gly Ala Leu Asn Ala Pro Ile Leu Met Thr Lys Gln Asn
 85           90           95

Gln Ile Pro Asn Thr Thr Met Glu Arg Leu Asn Lys Ala Lys Thr Val
100           105           110

Tyr Ile Ile Gly Ser Glu Ser Thr Ile Ser Lys Asn Val Glu Asn Gln
115           120           125

Leu Leu Ser Lys Lys Lys Val Val Gln Arg Ile Phe Gly Glu Asn Arg
130           135           140

Phe Asp Thr Ser Ile Lys Ile Ala Glu Lys Ile Lys Glu Ile Lys Pro
145           150           155           160

Ile Asp Lys Val Ile Ile Ala Asn Gly Phe Thr Gly Glu Ala Asp Ala
165           170           175

Ile Ser Ala Ser Pro Val Ala Ala Arg Asp Gly Val Pro Ile Ile Leu
180           185           190

Thr Asp Gly Asn Ser Val Gly Phe Asp Thr Thr Gly Leu Lys Ser Tyr
195           200           205

Ala Leu Gly Ser Ser Glu Ile Ile Ser Asp Glu Leu Val Lys Ser Thr
210           215           220

Asn Ser Ile Arg Leu Gly Gly Thr Asp Arg Phe Glu Thr Asn Lys Ile
225           230           235           240

Val Ile Gln Glu Phe Tyr Lys Asn Ser Lys Glu Phe Tyr Leu Ser Lys
245           250           255

Gly Leu Gln Leu Thr Asp Ala Leu Ala Ala Ser Thr Ile Ala Lys Asn
260           265           270

Ala Pro Val Val Leu Val Glu Asn Gly Ser Asn Lys Ser Ile Leu Ser
275           280           285

Gly Ala Asp Lys Leu Thr Val Leu Gly Gly Ile Asn Gln Asn Val Ile
290           295           300

```

Lys Gln Cys Ile Asn Gln Ala Ser Pro Asn Gln Gln Gly Leu Tyr Tyr  
 305 310 315 320  
 Asn Pro Asn Asp Arg Ala Phe Lys Glu Arg Ile Lys Gly Lys Val Tyr  
 325 330 335  
 Ala Leu Thr Lys Gln Tyr Arg Lys Glu Asn Gly Val Arg Ala Leu Ser  
 340 345 350  
 Val Ala Ser Arg Leu Glu Gly Leu Ala Asn Asp Trp Ser Asn Leu Met  
 355 360 365  
 Ala Asn Lys Lys Thr Leu Ser His Thr Ile Asn Gly Lys Asn Ser Tyr  
 370 375 380  
 Ser Thr Phe Leu Lys Tyr Leu Asp Trp Ser Glu Ile Lys Pro Gly Tyr  
 385 390 395 400  
 Ile Ala Val Gln Gly Glu Asn Ile Ile Lys Tyr Lys Ile Pro Asp Lys  
 405 410 415  
 Pro Val Tyr Thr Asn Arg Asp Ala Asp Asp Ile Gly Asn Phe Ile Phe  
 420 425 430  
 Asn Glu Trp Lys Thr Asn Pro Glu Glu Gly Thr Asn Met Leu His Lys  
 435 440 445  
 Gly Tyr Glu Ile Met Gly Phe Gly Ile Ala Ile Thr Gly Asp Lys Asn  
 450 455 460  
 Leu Tyr Ala Thr His Glu Phe Tyr Gly Arg Tyr Lys Glu  
 465 470 475

<210> 31  
 <211> 626  
 <212> PRT  
 <213> Clostridium difficile

<400> 31  
 Met Asn Lys Arg Lys Ser Phe Ile Arg Thr Ile Ala Val Ser Thr Met  
 1 5 10 15  
 Ala Val Ala Val Thr Gly Ser Ala Thr Cys Ala Tyr Ala Ala Pro Val  
 20 25 30  
 Leu Gln Gly Thr Lys Thr Tyr Glu Lys Val Asn Thr Ile Asp Ile Ser  
 35 40 45  
 Val Asp Ser Val Glu Asn Ile Val Tyr Ser Phe Gln Ala Ser Ile Lys  
 50 55 60  
 Val Gln Gly Glu Val Glu Val Val Asp Asn Glu Gln Lys Glu Lys Ile  
 65 70 75 80  
 Thr Trp Ser Asp Asn Ile Lys Ser Gln Ile Lys Ser Gly Asn Ala Asn  
 85 90 95  
 Ala Thr Cys Arg Ala Glu Tyr Asn Lys Ser Ser Asn Thr Thr Thr Leu  
 100 105 110

Asp Ile Tyr Val Thr Ser Asn Glu Asp Leu Leu Asp Gly Asn Arg Leu  
 115 120 125  
 Asn Ile Gly Arg Ile Ser Val Lys Lys Ser Gly Ser Asn Ser Asn Ala  
 130 135 140  
 Asp Tyr Lys Val Leu Gly Lys Gly Thr Ser Asp Lys Pro Ala Leu Lys  
 145 150 155 160  
 Ile Val Thr Tyr Asn Asn Lys Thr Val Asp Tyr Glu Asn Ile Ser Ser  
 165 170 175  
 Asp Glu Gly Leu Ile Phe Thr Leu Ile Asn Glu Ser Glu Val Lys Pro  
 180 185 190  
 Ile Gly Gly Thr Gly Ser Ser Lys Asn Asp Pro Glu Lys Tyr Lys Val  
 195 200 205  
 Glu Lys Ser Glu Ala Leu Glu Tyr Leu Leu Asn Asn Ile Arg Ile Asn  
 210 215 220  
 Tyr Ser Ile Val Ser Lys Glu Thr Gln Glu Ser Gly Ser Asn Val Ile  
 225 230 235 240  
 Leu Lys Leu Gly Leu Ala Gln Lys Thr Thr Lys Gly Arg Lys Ala Thr  
 245 250 255  
 Ile Asn Lys Tyr Val Glu Val Thr Leu Pro Lys Ser Leu Glu Tyr Ile  
 260 265 270  
 Val Glu Asn Glu Leu Ser Lys Pro Asp Glu Leu Pro Pro Asp Asn Gly  
 275 280 285  
 Ser Gly Gly Asn Asn Gly Gly Gly Ser Asn Ser Gly Gly Ser Ser Ser  
 290 295 300  
 Gly Gly Ser Ser Gly Gly Gly Asn Ser Ser Asp Ser Thr Ser Asn Val  
 305 310 315 320  
 Thr Val Lys Lys Leu Lys Gly Ala Asp Arg Phe Glu Thr Ala Ile Lys  
 325 330 335  
 Ile Ser Gln Ser Gly Trp Thr Lys Ser Asp Thr Val Val Ile Val Asn  
 340 345 350  
 Gly Glu Asp Lys Ser Met Val Asp Gly Leu Thr Ala Thr Pro Leu Ala  
 355 360 365  
 Ser Val Lys Asn Ser Pro Ile Leu Leu Ser Ser Asn Glu Lys Leu Pro  
 370 375 380  
 Gln Lys Thr Val Glu Glu Leu Lys Arg Leu Asn Pro Ser Lys Val Val  
 385 390 395 400  
 Val Ile Gly Gly Asn Asn Ser Met Pro Asn Ser Val Val Glu Ala Ile  
 405 410 415  
 Lys Ala Val Asn Ser Lys Ile Ser Val Gln Arg Ile Gly Gly Asp Thr  
 420 425 430

Arg Tyr Gln Thr Ser Ile Asn Ile Ala Lys Glu Ile Asp Arg Thr Asn  
 435 440 445  
 Asn Val Ser Lys Leu Tyr Ile Gly Ala Gly Asn Gly Glu Ala Asp Ser  
 450 455 460  
 Leu Ser Ile Ala Ser Leu Ala Gly Lys Glu Lys Thr Pro Ile Val Leu  
 465 470 475 480  
 Thr Gln Lys Asp Gly Val Asp Asn Glu Ala Glu Gln Phe Ile Lys Ser  
 485 490 495  
 Asn Lys Val Ser Asn Ile Tyr Phe Ile Gly Gly Val Glu Lys Ile Ser  
 500 505 510  
 Asn Lys Ala Ile Glu Gln Val Gly Lys Ile Ala Asn Lys Asp Ile Ser  
 515 520 525  
 Asn Asn Arg Val Ala Gly Gln Thr Arg Gln Glu Thr Asn Ala Lys Val  
 530 535 540  
 Ile Asp Lys Phe Tyr Ser Gln Ser Lys Leu Asp Gly Val Val Val Ala  
 545 550 555 560  
 Asn Gln Asp Lys Leu Ile Asp Ala Leu Ala Val Gly Pro Leu Ala Ala  
 565 570 575  
 Lys Asn Asn Ser Pro Val Ile Leu Ala Thr Asn Thr Leu Asp Lys Ser  
 580 585 590  
 Gln Glu Ser Ser Leu Lys Gly Lys Asn Ser Ser Lys Leu Phe Glu Val  
 595 600 605  
 Gly Gly Gly Ile Ala Ser Ser Val Ile Asp Lys Ile Lys Ser Leu Ile  
 610 615 620  
 Glu Lys  
 625

&lt;210&gt; 32

&lt;211&gt; 550

&lt;212&gt; PRT

&lt;213&gt; Clostridium difficile

&lt;400&gt; 32

Met Glu Asn Asn His Asn Ile Asn Ile Lys Tyr Lys Asn His Gln Gly  
 1 5 10 15  
 Asp Met Lys Met Asn Lys Lys Ile Leu Ser Leu Gly Leu Ala Val Ser  
 20 25 30  
 Leu Ile Leu Val Asn Phe Lys Ser Val Asn Ala Ser Ser Val Val Glu  
 35 40 45  
 Lys Ile Tyr Gly Lys Asp Arg Tyr Glu Thr Ala Ala Lys Ile Ala Asp  
 50 55 60  
 Lys Gln Thr Tyr Glu Thr Val Ile Leu Val Asn Thr Glu Lys Ser Leu  
 65 70 75 80



```
<210> 33
<211> 528
<212> PRT
<213> Clostridium difficile
```

<400> 33

```

Met Lys Val Asn Lys Arg Val Leu Ser Ile Gly Leu Ala Ile Ser Leu
  1             5             10             15

Ile Met Ala Gly Ala Pro Asn Ile Asn Ala Leu Ser Ser Ile Glu Lys
             20             25             30

Ile Gln Gly Lys Asp Arg Tyr Glu Thr Ala Ala Lys Ile Ala Gln Lys
             35             40             45

Gln Thr Tyr Glu Asn Val Val Leu Val Asn Thr Asp Asn Thr Leu Ala
  50             55             60

Asp Gly Leu Ser Ala Ser Gly Leu Ala Gly Thr Val Lys Ala Pro Ile
  65             70             75             80

Leu Leu Ser Gln Arg Asn Ser Ile Pro Ser Asp Thr Glu Lys Met Leu
             85             90             95

Lys Asp Val Lys Lys Val Tyr Ile Ile Gly Thr Glu Asp Ser Ile Gly
             100             105             110

Lys Ser Val Glu Asn Glu Leu Lys Gln Lys Gly Ile Glu Val Lys Arg
             115             120             125

```

Ile Gly Gly Asn Asp Arg Ile Glu Thr Ser Tyr Leu Ile Ala Lys Glu  
 130 135 140  
 Ile Ala Ser Ile Lys Pro Ile Asp Lys Val Phe Ile Thr Asn Gly Tyr  
 145 150 155 160  
 Thr Gly Glu Ala Asp Ala Met Ser Ala Ser Ser Val Ala Ser Arg Asp  
 165 170 175  
 Gly Ala Pro Ile Ile Leu Thr Asn Gly Lys Asn Val Pro Phe Glu Lys  
 180 185 190  
 Lys Glu Gly Val Gln Cys Tyr Ala Leu Gly Ser Glu Glu Ile Ile Ser  
 195 200 205  
 Asn Asp Leu Val Lys Lys Thr Asn Ser Val Arg Leu Ala Gly Glu Asp  
 210 215 220  
 Arg Phe Glu Thr Asn Lys Lys Val Ile Lys His Phe Tyr Ser Ser Ala  
 225 230 235 240  
 Lys Glu Phe Tyr Leu Ser Lys Gly Tyr Gln Leu Val Asp Ala Val Ala  
 245 250 255  
 Gly Ser Ser Ile Ala Lys Asn Ala Pro Ile Val Leu Val Asp Gly Asn  
 260 265 270  
 Ser Asp Lys Ser Val Leu Arg Ser Ala Asp Lys Ile Thr Ala Leu Gly  
 275 280 285  
 Gly Ile Asp Glu Lys Thr Leu Glu Gln Cys Leu Ser Ala Ser Ser Leu  
 290 295 300  
 Asp Ala Ser Ala Pro Thr Ile Thr Val Gly Asn Leu Asn Ile Tyr Gln  
 305 310 315 320  
 Gly Asp Lys Phe Asp Ile Ser Lys Leu Asn Ile Val Ala Lys Asp Ser  
 325 330 335  
 Asn Gly Asn Asp Leu Thr Pro Glu Leu Ile Gly Asn Ile Asn Thr Asp  
 340 345 350  
 Lys Val Gly Lys Tyr Lys Val Thr Ile Lys Ala Thr Asp Ile Gly Gly  
 355 360 365  
 Lys Thr Thr Ser Ile Ser Val Glu Val Asn Val Leu Glu Tyr Lys Thr  
 370 375 380  
 Asn Asp Met Asn Ser Ser Glu Phe Lys Arg Met Val Ser Ser Glu Met  
 385 390 395 400  
 Tyr Ser Leu Val Asn Ser Tyr Arg Lys Glu Lys Gly Lys Glu Pro Leu  
 405 410 415  
 Gln Val Ser Glu Asn Leu Gln Gly Met Ala Asn Tyr Trp Ser Lys Tyr  
 420 425 430  
 Met Ala Asp Lys Gly Glu Phe Ala His Val Ile Asn Gly Lys Asn Ala  
 435 440 445

Ala Glu Val Phe Ser Gly Gly Ile Arg Ser Glu Glu Asn Ile Ala Phe  
 450 455 460

Val Pro Leu Thr Thr Lys Ser Thr Tyr Thr Thr Lys Asp Ala Arg Glu  
 465 470 475 480

Val Ala Asn Val Ile Phe Thr Val Trp Lys Lys Ser Asp Lys Tyr Asn  
 485 490 495

Glu Asn Met Leu Ser Ser Lys Phe Ala Tyr Thr Gly Phe Gly Leu Tyr  
 500 505 510

Ile Leu Pro Asp Gly Gln Val Tyr Ala Thr Gln Glu Phe Leu Asn Lys  
 515 520 525

<210> 34  
 <211> 781  
 <212> PRT  
 <213> Clostridium difficile

<400> 34  
 Met Ser Val Ile Asp Ser Ile Leu Asp Lys Ala Asp Glu Gln Glu Ile  
 1 5 10 15

Lys Lys Leu Asn Val Ile Val Asp Lys Ile Asp Ala Leu Glu Asp Ser  
 20 25 30

Met Lys Asn Leu Ser Tyr Glu Glu Leu Lys Asp Met Thr Ala Ile Phe  
 35 40 45

Lys Asn Arg Leu Lys Lys Gly Glu Thr Leu Asp Asp Ile Leu Pro Glu  
 50 55 60

Ala Phe Ala Val Val Arg Glu Val Ser Lys Arg Lys Leu Gly Met Arg  
 65 70 75 80

Gln Tyr Arg Val Gln Leu Ile Gly Gly Ile Val Ile His Gln Gly Lys  
 85 90 95

Ile Ala Glu Met Lys Thr Gly Glu Gly Lys Thr Leu Val Glu Val Ala  
 100 105 110

Pro Val Tyr Leu Asn Ala Leu Thr Gly Lys Gly Val His Val Ile Thr  
 115 120 125

Val Asn Asp Tyr Leu Ala Glu Arg Asp Lys Glu Leu Met Ser Pro Val  
 130 135 140

Tyr Glu Ser Leu Gly Met Thr Val Gly Val Ile Ile Ser Asn Gln Asp  
 145 150 155 160

Pro Asn Ile Arg Lys Gln Gln Tyr Lys Cys Asp Ile Thr Tyr Gly Thr  
 165 170 175

Asn Ser Glu Phe Gly Phe Asp Tyr Leu Arg Asp Asn Met Val Pro Asp  
 180 185 190

Leu Ser His Lys Val Gln Arg Glu Leu Asn Phe Ala Ile Val Asp Glu  
 195 200 205



Val Asp Ser Ile Leu Ile Asp Glu Ala Arg Thr Pro Leu Ile Ile Ala  
 210 215 220  
 Gly Asp Gly Asp Glu Asp Leu Lys Leu Tyr Glu Leu Ala Asn Ser Phe  
 225 230 235 240  
 Val Lys Thr Val Lys Glu Glu Asp Phe Glu Leu Asp Arg Lys Asp Lys  
 245 250 255  
 Thr Ile Ala Leu Thr Ala Ser Gly Ile Ser Lys Ala Glu Ser Phe Phe  
 260 265 270  
 Gly Ile Thr Asn Leu Thr Asp Ile Lys Asn Ile Glu Leu Tyr His His  
 275 280 285  
 Ile Asn Gln Ala Leu Arg Gly His Lys Leu Met Glu Lys Asp Val Asp  
 290 295 300  
 Tyr Val Ile Ser Asn Gly Glu Val Met Ile Val Asp Glu Phe Thr Gly  
 305 310 315 320  
 Arg Val Met Asp Gly Arg Arg Tyr Thr Asp Gly Leu His Gln Ala Ile  
 325 330 335  
 Glu Ala Lys Glu Gly Val Glu Ile Lys Asn Glu Ser Lys Thr Met Ala  
 340 345 350  
 Thr Val Thr Tyr Gln Asn Phe Phe Arg Leu Tyr Glu Lys Leu Ser Gly  
 355 360 365  
 Met Thr Gly Thr Ala Lys Thr Glu Glu Gly Glu Phe Glu Ser Ile Tyr  
 370 375 380  
 Lys Leu Asn Val Val Gln Ile Pro Thr Asn Arg Pro Val Ile Arg Ala  
 385 390 395 400  
 Asp Leu His Asp Lys Val Phe Lys Thr Glu Glu Glu Lys Tyr Ser Ala  
 405 410 415  
 Val Val Glu Glu Ile Ile Arg Ile His Lys Thr Arg Gln Pro Ile Leu  
 420 425 430  
 Val Gly Thr Val Ser Val Glu Lys Ser Glu Lys Leu Ser Lys Met Leu  
 435 440 445  
 Lys Lys Gln Gly Ile Lys His Gln Val Leu Asn Ala Lys Gln His Asp  
 450 455 460  
 Lys Glu Ala Glu Ile Ile Ser Lys Ala Gly Lys Leu Asp Ala Ile Thr  
 465 470 475 480  
 Ile Ala Thr Asn Met Ala Gly Arg Gly Thr Asp Ile Ser Leu Gly Ala  
 485 490 495  
 Gly Asp Lys Glu Glu Glu Gln Glu Val Lys Asp Leu Gly Gly Leu Tyr  
 500 505 510  
 Val Ile Gly Thr Glu Arg His Glu Ser Arg Arg Ile Asp Asn Gln Leu  
 515 520 525

Arg Gly Arg Ser Gly Arg Gln Gly Asp Pro Gly Thr Ser Arg Phe Phe  
530 535 540

Val Ser Leu Glu Asp Asp Val Ile Lys Leu Tyr Gly Gly Lys Thr Ile  
545 550 555 560

Glu Lys Leu Met Lys Arg Thr Ser Ser Asn Glu Asn Thr Ala Ile Glu  
565 570 575

Ser Lys Ala Leu Thr Arg Ala Ile Glu Arg Ala Gln Lys Gly Val Glu  
580 585 590

Gly Lys Asn Phe Glu Ile Arg Lys Asn Val Leu Lys Tyr Asp Asp Thr  
595 600 605

Ile Asn Glu Gln Arg Lys Val Ile Tyr Asn Glu Arg Asn Lys Val Leu  
610 615 620

Asn Asp Glu Asp Ile Gln Glu Asp Ile Gln Lys Met Val Lys Asp Ile  
625 630 635 640

Ile Gln Glu Ala Gly Glu Thr Tyr Leu Ile Gly Arg Lys Arg Asp Tyr  
645 650 655

Tyr Gly Tyr Phe Lys His Leu Tyr Ser Thr Phe Met Pro Ala Asp Thr  
660 665 670

Leu Leu Ile Pro Gly Val Asp Lys Lys Ser Val Gln Glu Ile Ile Asp  
675 680 685

Ser Thr Tyr Glu Ile Ser Lys Arg Val Tyr Asp Leu Lys Lys Met Met  
690 695 700

Leu Gly Ile Asp Lys Val Ala Glu Leu Glu Lys Thr Val Leu Leu Lys  
705 710 715 720

Val Val Asp Gln Tyr Trp Ile Asp His Ile Asp Ala Met Glu Gln Leu  
725 730 735

Lys Gln Tyr Ile Gly Leu Lys Ser Tyr Ala Gln Lys Asp Pro Phe Lys  
740 745 750

Glu Tyr Ala Leu Glu Gly Tyr Asp Met Phe Glu Ala Leu Asn Lys Asn  
755 760 765

Ile Arg Glu Ala Thr Val Gln Tyr Leu Tyr Lys Phe Asn  
770 775 780

<210> 35

<211> 48551

<212> DNA

<213> Clostridium difficile

<400> 35

aggatatgatt tatgtttttaa taggagttat agctatagca ggactatttg ctataaaaaa 60  
gttttttagta acaaaagaca agagtgtctt taatataatg agtattcatg atgaagaagc 120  
aattaaggaa caacagaaca aagaagacga agataaatct atttagacaa taaattatat 180  
gaatagctta ttacaaatat atcttaagaa ataatgagaa ataaatatga ttaaggagaa 240  
ctttatgtac gcagtaagat acgaattgat aaagacttgt aagcagagtg gtgcaagggt 300  
aggaagatta cacactccac atgggattat tgaaacacca atattcatgc cagttggaac 360

tcaagcaact	gtaaaatcta	tgacaccaga	agaattaa	gagattgggt	cacaaataat	420
acttagtaac	acataccatt	tatatatgag	accaggtcat	gaacttataa	aaagagctgg	480
tggtcttcat	aagtttatga	attgggataa	accaatactt	acagatagtg	gaggggtcca	540
ggtatttagt	ctagggcctt	taagaaagat	aaaagaagaa	ggtgtagaat	ttaggtcaca	600
cttagatggt	tcaaaacatt	ttctaactcc	agaaaaagct	atggaaatac	aaaatgcatt	660
gggttcagac	ataatgatgg	catttgatga	gtgtgcacca	tacccatcag	atagagaata	720
tgtaaaaaat	tcttttagaaa	gaacaacaag	atgggttaaaa	agatgtaaag	atgctcataa	780
taataccgac	aagcaagctt	tatttggtat	aatacaaggt	gggatgtata	aagacttaag	840
ggaacaatca	gcaaaagaaa	taactaatat	tgatttacca	ggatatgcta	ttggtggact	900
tagtggttga	gaaccaaacc	cacttatgta	tgacgtgtta	gagcatacaa	ctccacttat	960
gccaaaggat	aaaccaagat	acctaattggg	agttggtagt	ccagatgact	tagtagaagg	1020
tggtataaga	ggagtagaca	tggttgatgt	tggtctacct	actcgtatcg	ctagaaaagg	1080
aactgcaatg	actagccaag	gtaaagtagt	gtttagaaat	gcaacttatg	ctgaagactt	1140
tactccactt	gacctgaat	gtgattgtta	tgcttgtaaa	aactattcaa	gagcttataat	1200
aagacacctt	ataaaagcaa	atgaaatact	gggtgcaaga	tttaataacta	ctcataatct	1260
tcacttttta	ttaaatttaa	tgaaacaaat	aagacaagcg	ataatggaag	atagattact	1320
tgatttttaga	aatgaatttt	ttgcaaaata	tgatatgaa	atatagggtt	tgactatta	1380
attataatca	ttttaatata	taatatctaa	tagtactcaa	agatagagaa	agtaaatgt	1440
tatataaaca	tttattatga	tttacaanaag	agtataat	acaaaggagt	gaaaaattaa	1500
tgccacagca	acaaataatt	atgagtatag	gtctatgggt	agtagtaatc	gcaatctttt	1560
atttccta	gatttagacct	caaaagaaaa	aagacaaaca	attaaaagaa	atgagaagta	1620
gcctaagtgt	aggagacaag	gtcattacaa	taggtggaat	agtagcta	gtagctaag	1680
ttgaagatga	tggtgtaata	ttagaatttag	gaccaaatag	aactaaagtt	ccatttgaaa	1740
aatgggctat	aggaactggt	aaatctaaaa	aagaagaaat	agaagaagac	taaattagta	1800
atttaaaagc	cttttctcga	atatatatta	atttgagagg	agggtttttt	atgggaaaaat	1860
ctaattatat	ttttaaagg	ctaggatatg	catatataat	aactttggct	gttttattag	1920
tgtataactt	atttttgaca	tttacagata	ttggtggaga	taatataact	atggtctcat	1980
catttataac	tactatttct	gcagctatag	gaggggtttt	cacttcaaaa	cacatgaaag	2040
aaaaaggact	tatgtatggg	cttttagtag	gacttttgta	catagtatgt	atttttttga	2100
ctgtgttttt	agcacaagaa	aagtttgat	ttgaagtagg	gatgatttat	aagttattgc	2160
taatatcagc	agcaggaggc	attggaggcg	tattgggtgt	caacttttaa	tagagtttat	2220
attaaagtata	ttttatgata	taatcaatag	gaaacaatta	gaaatggagg	cgttgaaaa	2280
ggaaaatata	cagttaaaaa	ctttatctca	agctacttta	aagcaaaagt	ctgcaaaagg	2340
tggtatgtga	gagtgtaaaa	cttctgttca	atcagcttgt	aaaacttctt	gtacagttgc	2400
aaatcaagag	tgtgaaagat	agaaaattat	tgatttagtt	gtttgtaaga	gggtatctca	2460
aagtgattct	tttatcatta	tgaggtatcc	tcttttattt	tcaattatta	tataaaaaata	2520
tattctattt	atgtaaaatt	aaatttgatt	tcttatcggt	taattatgca	aaaatatcta	2580
atcatataag	agcattcatt	ttaaagtagc	tactcttata	ttttattttt	tatcaattac	2640
atgtattaga	ataattttta	attttaaact	ctaatttatg	atgattttta	taatatgtta	2700
tatattttaa	tatatggaga	tagatattaa	tttagtgata	tatatttaag	taataagagt	2760
aaagcaaaaa	ttagtgcata	gtagtatata	aaaatggagt	ttgaataact	ttgttagtat	2820
atgattgttt	agtagaaaat	cactataaaa	taggatttct	atctttgagc	ataattatag	2880
agagctaagt	agggggaagg	gggtcaagaa	gatattatta	gtgtttgtgt	tagtaaat	2940
tgtaaacaca	taaaaatata	tgaanaataa	tattatatat	aatattattt	atgtgtggga	3000
aatacgaata	taaaaaataa	aaaataatat	tcggaggaca	taaatatgtt	aagcaacaaa	3060
aagagaagta	tggcaatagt	aatggcaggt	gctacagtta	tgagtgcagc	agcaccaata	3120
ttcgcagaca	atactgtgac	agaaaatgtt	gataaaaaat	acacagtaag	tgcgaaagat	3180
tctgctaacc	taatcgaaga	ggtagaaaa	gcattagaag	ttagtttga	agatacaaaa	3240
gcaggagcaa	atgtaaatga	tagagtttac	gatataaaag	tagacaatgt	aaacttaact	3300
aatgcaactc	aattacaaaa	taaaataaac	tctttaacag	aagggcaaa	tttaaaagt	3360
actattcaag	ataaagggca	tcaagtgtta	gggtggaaa	tagttgacta	taaaattgaa	3420
aactacaaaa	ctgctcaaga	aatagtagat	gcagttaatg	cttacaatgc	aactttagca	3480
gaagattctg	ataacaagtt	aacagcaact	ataaaatcta	caaatacagt	tgaagtaaaa	3540
agagcaaaa	attcagcaaa	tgttattaca	ttaaatgtag	gagaccaaca	tttagatttc	3600
tcaaaagtga	taacatcaga	agaaggtact	tttgaaggtt	atgaaaagcg	ttatagtgc	3660
atagattcaa	aagaattaca	tacggtaaca	gttaaaaaac	cagatttaca	agacatatca	3720
gcagaagaat	tatttgatgg	aattagatta	actacattag	gaagagaaat	agttaataaa	3780
gttaaaaaat	gatatgcttt	aacatttgaa	aatgaagcaa	tccttactca	agaacaagaa	3840
gatagtgatg	ataaagataa	accagaaaaa	tcaagctttg	atatagtttt	aagcaaggca	3900
aatgaaaagc	cagaaacaat	aagtgtttca	agtaaaaacc	ataagttagt	tagagactta	3960
cacaaagtat	taactgatgt	aaaagatgga	aaagaattaa	aagtagaagt	tttatctggt	4020

gattcaagat	ttacaacagc	agtagaagtt	agtaaagaga	gattcaaaga	tggagaagca	4080
gaagctataa	tttttagttg	agaagatgct	atagttgatg	gatttagcatc	agcaccactt	4140
gcctctcaaa	aaaatgcacc	aattattatta	tctaaaaaag	attcactacc	atcagaaata	4200
gaagctgaaa	tattaagagt	acttggaagt	aattctatctt	ctaagaaaat	atatatagta	4260
ggtggagaat	ctaaagtatc	aaaagaaaact	gaagaaaaac	tttctaaatt	aggtgtaagt	4320
aaagttgaga	gagtttctgg	agaagataga	tttgaaaactt	cttttagaaat	agcaaaaaca	4380
ttaaaagata	catttaagac	tgcttttgta	gtaggtggaa	atggagaagc	tgatgctatg	4440
agtatatcag	ctagagctgc	tcaatttggg	gctccaataa	tagttacagg	taacgaatta	4500
gatgcaaatg	ctgaaaaatt	attaaaagga	aaagaattag	aaatagtggg	tggagaaaat	4560
tctgtatcaa	aagaagttga	agacaaaatta	gttgatatag	atttaaataa	taaagttgaa	4620
agattagctg	gagaaaatag	aaaagatact	aatgctaaag	taatcaataa	atactatgca	4680
ggtgcaacta	aagcatatgt	agcaaaaagat	ggttatgtag	gtggaaatgg	acaattagtt	4740
gatgcactta	cagcagcacc	acttgcagct	agttcaaaaag	ctccaatagt	attaactaca	4800
gaagaacttt	ctaaatcaca	agaagaagta	gttgagttaa	gacttaagaa	tgctactaag	4860
ttagtacaaa	taggtgaagg	aatagctaaa	aacgctattg	aaaaaatagc	agaaaaaata	4920
aacttatttta	ctaagaacta	ataagatata	ggaattaata	ttaaaagt	atttagataa	4980
tatatgatta	tacaatgcgt	taatatgtaa	gacatagtgt	agatattata	aaatatacaa	5040
ataacaagat	aaaataataa	aaataaggtg	tctcattttag	aattagattg	agatacctta	5100
ttttatttttt	tgattataat	gttgagaaat	tctatttgca	aaaattatta	taaaaagata	5160
cttctttataa	tgattaaaca	tattactaac	ataaattatt	ttgaacttat	aacttaatta	5220
aataaaataa	attataatgt	tttgaattta	tagagaataa	aaactttagg	tataattgtt	5280
gagttgtatg	gataactatg	gtatcatata	taaaatatgg	taacaaaatt	gtaacttttg	5340
attaaaaata	agtatgagta	ctaataaaaa	atatgaaaag	tacattgggg	gtagaaaata	5400
atatgaagaa	tagtaaaaag	atattagcta	taggacttac	actattttta	gtaatggtaa	5460
atactcctat	ggtaaagtgcg	ttgacatcag	ttgagcaaat	aaaagggaat	gacagatatg	5520
agacagcagc	aaaaatagca	gataagcaaa	actacaatac	agcaatacta	atcaactcag	5580
ataatagctt	agcagatggg	ctaagtgcga	gtggtttagc	tggagcctta	aatgcgccta	5640
ttttgatgac	aaaacaaaaat	cagattccaa	acacaactat	ggaaagatta	aacaaagcaa	5700
agactgtgta	tataataggc	tcagaatcaa	caataagtaa	aatggttag	aatcagttac	5760
tatccaaaaa	gaaggttgta	caaagaatct	ttggtgaaaa	tagatttgat	acaagtataa	5820
aaatagctga	aaaaattaaa	gaaattaagc	caatagataa	agtaattata	gccaatggat	5880
ttacaggaga	agcagatgcg	ataagtgcac	caccagtagc	tgctagagat	ggagtaccta	5940
taatacttac	agatggaaat	agtgtgggat	ttgacacgac	aggcttaaag	agttatgcgc	6000
ttgggtctag	tgaaataata	agcgacgaat	tagttaagag	cacaaattct	attagattgg	6060
gtggcacaga	tagatttgaa	actaataaga	tagttataca	agagttttat	aaaaatagta	6120
aagaatttta	tctaagtaaa	ggcttacaat	taacagacgc	tttggtgcc	tctacaatag	6180
caaaaaatgc	tcctgtagtt	ttagttgaaa	atggcagtaa	taagagtatt	ttaagtgggtg	6240
cagacaaatt	gactgtacta	ggaggcataa	atcagaatgt	tatcaagcaa	tgtataaatc	6300
aagcttcacc	aaatcaacaa	ggtttatat	acaattccaa	tgatagagca	tttaaagaga	6360
gaataaaggg	taaagtatat	gctcttaca	agcagtatag	aaaagagaat	ggagtaagag	6420
cgttatctgt	agcaagcaga	ttagagggtc	tagcaaatga	ttggtcta	ttaatggcaa	6480
ataagaagac	tttatctcat	acgataaatg	gaaaaaattc	atattctact	tttttgaaat	6540
acttagattg	gagttagatt	aaacctggat	atatagcagt	tcaaggtgaa	aatataatta	6600
aatataaaaat	tcagataaaa	cctgtatata	caaatagaga	tcagatgat	ataggaaact	6660
ttatTTTTTaa	tgagtggaa	acaaatccag	aagaaggaac	taatattgtg	cataaagggt	6720
atgaaataat	gggattttgg	atagctataa	caggagataa	gaatctatat	gctacacatg	6780
aatttttatgg	aagatatataa	gaataagaaa	atatacacaa	aattatgctt	gtaagggaagc	6840
tatatataa	attaaattat	aatggaaaaat	agcattttata	ttataaaatt	aatataaatg	6900
ttatTTTTTtt	ataattta	gtaaaaaaat	tgtatttaaa	agtttaaaata	tagtataaaa	6960
tgggtaagaa	agtagtgaaa	attaaatggc	aaaattaggg	aaatctaata	acgcaaaact	7020
atagggatta	aagcttttta	agccatatca	gccagttgct	aaaaagagta	aagtcctttt	7080
attttataat	taagattctc	tttgagtgtc	tttttttatt	gtctgaaact	tatagtattt	7140
aaattcatga	aaaattagat	atttattaag	ggaggattgt	tatttcatga	aaaaggcaat	7200
atcttgtgta	ctagcagtat	ctatgtgtag	tgccaccata	aacgtttttg	cagaacctat	7260
attagagggg	aagctaaggg	cagttgaaaa	atcagttaca	gaaaagttaa	gggggagctc	7320
tgaagtagat	ttaaattttt	cactgccaat	aaaaaatagc	gagtcagaaa	gtatgactaa	7380
tatgcagtta	cgattaaaag	atgatagtaa	caatagtggg	ataataaagc	ttggagaaaa	7440
agattcagga	acagtaaatg	ttgggacaga	taatatagaa	tatactataa	aaaaattatc	7500
tgcaagtaga	agtgaataaa	aaaataaaga	tgaaaatgta	tcatattata	atattgtatt	7560
taataatttta	cctgttgga	aatacaatgt	agaagtctct	ggagctggg	ttaagaacaa	7620
agaaattaaa	gacattgata	tatcaagtta	ttcacaagga	gtcttattaa	gcaatagagc	7680

aagtgtagat	aataaaaaata	aaccttttaaa	ttatgaacta	ttcttaaatgg	gtgatgtaaa	7740
tggagacggg	aaagtagata	agagtgatta	taataaggtt	ttagagaata	tagattcaaa	7800
taaaagagaa	tttgatttaa	atagagatgg	aaaggtagat	attgttgact	tagattatgt	7860
acagaaaaat	ttaggacaaa	gtgaagatgc	aaaaagccta	gaaagtatag	tatctacca	7920
tcctatagta	gatacaagta	aggttgaatt	aaaaggaaat	agtgatgtag	atattaatgg	7980
taatgtagaa	aattttattca	gtggagaaaa	taatggagtt	acaatatctt	caaaaaatga	8040
tgaatatatct	gaggaaaaac	cagcagtaat	ggatatagct	ttatcagaag	cacttaatat	8100
ggaagtcac	aatataaaaag	ctccagaagg	tactgctcca	gcaaaaggaa	tagttgaggt	8160
tactgacaaa	gatggtaata	ttcaacagat	tccttatgaa	aacaaagaat	atatgaaaaa	8220
cttaaaagat	gaggagcaac	aagaggaagt	gaataacgaa	caagaagaag	taaatcaaga	8280
agaagtaaat	aataaagatg	aaaatataaa	ccagactgat	aataactcaa	ataataaaga	8340
tgaaaataat	gataaacaac	aggaaaatag	cgataaacia	gaagcaaact	ctaataccaa	8400
tcaagataaa	aataatgata	aagaaaattc	atcaatggat	aaacctaatt	cagatgaaga	8460
tactccaaaa	gaaggggata	atatcagtaa	atgatataac	tttacaacat	ctacatatag	8520
taggtcagca	ttagaacat	atgcagatgt	tcaaaatagt	gatatagtaa	taaatcttgg	8580
taaaacaagta	gcagtaaaaa	agataactat	aaaaataaca	gctacaacag	caaatgatag	8640
aaatcttgca	gaaatatcaa	aagttgaatt	tttaacaat	gtatataaag	aaataccaaa	8700
gccagagatg	aacataccta	aaataaataa	aatattgact	agcactgctg	ttggaaatga	8760
aaaagtatct	ttttattgga	ataatgaagt	aaatgtatca	gcatacgaac	ttatactaga	8820
caaaattgat	agtaagggaa	aagtattatc	tactaaaaaa	ttacaaacca	gtaaaaatac	8880
aatagaagtt	agagatatag	aagcatatgg	actatataga	gtagcatac	aatcttttaa	8940
tggagattgg	tctagtgggt	ataaagatgc	aacaccagaa	aacttagacg	gagtaccaga	9000
aaatgtagat	tcaaactata	aacctataca	atttagccca	gatagtatat	tagaatttca	9060
agttgttcca	gacacaaaac	cagaaccacc	agaaggtata	aatgttaaag	gtaaatttaa	9120
atctcttgat	gtaagctgga	aaaatcataa	acaagctaaa	gattttgact	tgtactataa	9180
agaaaaaggg	tctaattgggt	cttgataaaa	atacaatgat	aagcttattc	gtgggacaag	9240
ttcaaagata	gaaggtctta	aggataatac	aacatatgaa	gttagaatga	cagctactaa	9300
ccacttggga	actagtggaa	tgtctaaaac	atatataggt	acaactgaaa	gcatggatgc	9360
accaacaaca	cctaactata	gattaattaa	tactcctaaa	gagaatgggtg	gatttggaa	9420
accaacaaat	cacattgtag	atgttactta	tccagcagga	tatgacaaaa	gtgaatatag	9480
tgataaaaaat	ccatttgata	aattcaatgt	agtagatggg	gactttacaa	cacattggac	9540
attoccaaca	tataatgcag	ctgagggaa	gaatagagga	ccaataataa	cttttgataa	9600
atcttatact	atggatactt	ttatggctac	acctcgtttt	gatgtttctg	ataagacaat	9660
aaatgacttt	gaggtacgat	attgggatga	aaatgggtcaa	ttacacaatc	tgggggatgt	9720
tccttttgaa	acaaagacag	accctgtaaa	taataataag	tatgtaatgg	ttagatttaa	9780
agaaccaata	actgctacca	agatacaagt	aagtgtttgt	tcctctatt	catataagtc	9840
taatatatca	gaacttaagt	tttatgagta	tgacagtata	gaagatgatg	taagaaatct	9900
atztatggat	gacctacaag	ttgaattaaa	agaagatgta	actcaagaaa	aaataactga	9960
attaaaagat	agattaaata	ctccagataa	agctagtggg	gaatatcatc	catataagac	10020
agttattgag	agagaattaa	aattagctca	agacttatat	aatgatagag	ccacacttac	10080
tgatgtaact	actgtaaatc	aagaaataaa	taaattagga	aaatataaag	ataaaaacgg	10140
acagtatact	gtaaattcca	ataatttagg	tatgcaaaa	gactggcagg	cattaggagt	10200
agctgctaga	gcaggagatg	aaatcacagt	atatgttggg	agtaaaaagt	gaaaaactcc	10260
taagttgata	tatacacagt	actatggaga	atctgggtgca	tataagagtg	gagaaataaa	10320
tcctttgact	ggaaaaaatg	taattcaatt	aagcaaaact	catagttag	atatagaatg	10380
tggaggaagt	ttatacataa	gatacgtgta	agatactcca	agtggaggag	atataaaaagt	10440
acgtgtagca	ggagctacta	agataccaca	ccttaatcta	aatggaacaa	taaatgataa	10500
gagtcctgog	ggagtctctg	aatctaagaa	aaagataaaa	gcatatatag	agaatttagc	10560
tagatataaa	gatgctgtaa	aaggagaacc	atattatcca	ggaacaaatg	ttggtaattc	10620
atatggatat	tcagaaaaaa	ctgggtgtgt	aaacactaca	gatattgaaa	gtgataaagt	10680
aacattaaat	gttctctgcaa	cagctgtata	tcaagggata	tcaaaaggaa	atgctgattt	10740
aaatacacaa	gtagatagat	tatataatag	tatgcttgca	tgggaacaaa	taatggattt	10800
agtttactca	gaaagaggag	tattttaaacc	acaagattta	gatggtaatg	gaactgtaga	10860
tgataatgaa	tttaacttaa	ctaaaaatga	tatggctcca	aaatctcgta	tgaacataaa	10920
ataaccaaga	atgtttattg	gagcatttat	gatgcttctc	ggattacatg	taggagtaga	10980
atatggttca	gttccagggc	ttatgaaatg	agtacctttc	caaatagatg	atagtggaaa	11040
agctacagga	ggaaacttat	ttggatgggg	aataggtcat	gagataggac	atgttacaga	11100
tattggaaaag	atgacatata	gcgaaactag	taacaatgta	cttgctttac	ttgctcagac	11160
ctttgatgat	aagacacatt	caagacttga	aggttcaaca	atggataaga	tatatgagca	11220
tgtaacttca	aatagcttag	gtataccaag	caatgtattt	gaaaggcttg	gaatgttatg	11280
gcaattacat	ctagcttatg	atgatgattt	tacgggttca	atgttaaaga	ataactcaga	11340

tgctgattta	agtaatgata	cattctatgc	aaagattttca	agaaaataca	gagcactttc	11400
aagtagtgac	ccaattaata	gtttgccaaa	agaccaaaatg	ctagtagcaa	tggtcatcaag	11460
tggtgttgaa	aaagattttaa	gagattactt	taaggcatgg	ggagtagaaa	ttacaccaga	11520
attaaacagt	ataatggata	gtaagaacta	cgaaaaagaa	tcaagaaaga	tacaatatgt	11580
aaatgacgaa	gctagaagaa	agatttttaa	tgaaaatata	gctagtatgg	caaaagatac	11640
taaaagtaagt	gctagttttt	ctgatggaa	tcaaagtgg	tctctagtta	aatctaataa	11700
aatatctata	gacctagatg	tagataagga	taaagataag	atactaggat	acgaaataat	11760
aagaagtgat	ggaaattatt	tagatgggtg	aaatggaaca	caggtaaaat	acagaacagt	11820
tggatttgtg	gatgcaaaa	gaaatagtaa	gactaaattt	actgataata	tatctcctct	11880
taataacaga	gcattttacat	ataaggtagt	tgcatatgat	tatcacctaa	atccaacaga	11940
agaatttgaa	gttggtactg	taaaattatc	agatgaaggg	aaaataaata	aatcagcatg	12000
gtcattcata	acaaatactg	tttctgacgg	tgatgttaga	actgaaaatg	actcacatgg	12060
accattacaa	aatcctgaaa	ttgataatat	aaaagataat	gatgcaagta	ctacatatata	12120
aggaaagata	ataagtaaa	cagagtggaa	caaagaccct	caaaaagacc	ctgatataaa	12180
tgtagaagaa	aatccatata	taatagtaga	catgaaagaa	acattaccta	tagtaggact	12240
aaaataacata	aaaccagagg	cagaagctag	gaagtgttca	ttaaaaggat	tatttcaact	12300
tagaaagaat	actagcacaa	catacaatcc	tttaactaat	tacaagatac	aagtaagtaa	12360
tgataaaaa	aactgggaag	atgtaagtca	aggaaactttt	gaatatggac	aagatttagt	12420
tggtggagga	aaagaccaag	ataatgaagc	tagagttaca	tttaataaa	atggtaaaact	12480
ttggacatac	caagctagat	atgtaaaact	aatctcaa	aataagtcta	atatagaagt	12540
tgacagaatta	gatataatag	gacctccagg	tgataatata	gaaattggca	caatgaataa	12600
tgacagacaa	actcgtacta	atggaattgg	aaaattagca	gaagattatg	tatatcaaa	12660
tgatgactct	agcacaagtg	aagatgaaac	aggtaagatt	ccagcaggtt	caatagttat	12720
tacaggtgaa	tacagtggaa	atccagcatt	taattttacca	ctacttatag	ataaaaaaca	12780
taagactata	tctggagaag	cactttttatt	tgacagaagt	ccagaaaaag	gtgaactagg	12840
tgaaatcagt	aaaggtacat	gggttttacta	tatgccaaaa	gaaaactttg	attcctttaag	12900
tgataaaggt	aaggctgaac	tatatagata	caatgacctt	caaggagata	ctccagtagg	12960
tcaaaagattt	gtaagtgata	caatgtatgt	acctgttagg	gctaagaatt	atgatgagtt	13020
aaagacaata	agtcttactg	atagtaatat	taaagctaga	aaagctacaa	gaaatgctat	13080
tgatatcagt	aataagaagg	ttataactgt	aagcaataaa	gtaaagggta	atattgttag	13140
agataaatag	taggtggaac	tctaaaaacg	taaaaatact	aacactaaca	ttgtatatct	13200
aataagatta	ggtactagaa	tagtttttaa	atagtagata	ttaaaagtaa	agtatatata	13260
tgtaactata	agtttaaggt	aattttataaa	ttaaatataa	aagctaggtg	gataagtaaa	13320
tttatccact	tagttaatga	tagataagga	cgggtgttaa	tgaataaaa	aaaatccttt	13380
ataagaacta	tagcagtatc	aactatggca	gtagctgtaa	ctggtagtgc	tacatgtgca	13440
tatgcagctc	cagttttaca	aggaacaaaa	acgtatgaga	aagtaaatac	tatagatata	13500
tctgtggact	cagtagaaaa	catagtatat	tcatttcaag	ccagtataaa	agtacaaggt	13560
gaggttgaa	tagtagataa	tgagcagaaa	gaaaagatta	cttggctctga	taatataaaa	13620
tctcagataa	aaagtggaaa	tgcaaatgct	acctgtagag	cagaatacaa	caaatcaagt	13680
aatacaacta	cattagatat	atacgttaca	tcaaatgaag	atttactaga	cggaaataga	13740
ttgaatatag	gaagaatatc	tgtaaaaaaa	tctggcagta	actcaaatgc	tgactataaa	13800
gttttaggaa	aaggaacaag	tgataaacca	gcttttaaaa	ttgtaacata	taacaacaaa	13860
acagtggatt	atgaaaatat	aagctctgat	gaaggattaa	tatttaccct	tattaatgag	13920
agtgaagtaa	aaccaattgg	tggaactgg	tctagtaaaa	atgaccctga	aaagtataag	13980
gtagagaaaa	gtgaagcact	tgaatattta	ttaaacaata	ttaggataaa	ctattctatt	14040
gttagtaaa	aaacacaaga	aagtggttca	aatgtaatat	taaaacttgg	attagcacia	14100
aaaacaacta	aggggaagaa	agctacaata	aacaaatatg	ttgaagttac	acttccaaaa	14160
agtttaga	acatagttga	aaatgaatta	tcaaaaccgg	atgaattacc	accagataat	14220
ggtagtgg	gaaataatgg	tggaaggaag	aattcaggcg	gctcatctag	tggtggaagc	14280
tctggaggag	gaaattcaag	tgattctaca	tctaattgtca	cagtcaaaaa	attaaaagg	14340
gctgatagat	ttgagacagc	tatcaaaata	tctcaaagt	ggttgacaaa	atcagatata	14400
gttgtaatat	taaatggaga	agataaaa	atgggtggatg	ggcttacagc	tacaccactt	14460
gctagtgtaa	aaaactcacc	gattctttta	tcatcaaatg	aaaagcttcc	tcaaaaaact	14520
gtagaagaat	tgaaaaggct	aaatccatca	aaagtagttg	tgataggtgg	aaacaactca	14580
atgccaaatt	cagttgttga	agctatcaaa	gctgtaaatt	caaaaatatc	agtacaaaaga	14640
ataggtggag	atacaagata	tcaaaactct	attaatatag	caaaagaaat	agataggacg	14700
aataatgtg	gcaagttgta	tataggagca	ggaaatggag	aagctgattc	attatcaata	14760
gcatcactgg	ctggtaaa	aaaaactcca	atagtactta	cacaaaaaga	tggtgtagat	14820
aatgaagctg	aacaatttat	taaatcaaat	aaagtgctca	atatatat	tattggtgga	14880
gtagaaaaaa	tatctaacaa	ggctattgaa	caagttggaa	aaatagcaaa	caaagatata	14940
tcaaaacaata	gggtggcagg	tcaaaactaga	caagagacga	atgccaaagt	aatagataaa	15000

ttctactctc	aatctaagct	agatggagtt	gtagttgcaa	atcaagacaa	acttatagac	15060
gcttttagcag	ttggaccatt	agcagctaaa	aacaattcac	cagttatact	tgctacaaat	15120
acgttagata	aatcacaaga	atcgtcatta	aaaggtaaaa	attcatcaaa	gttgtttgaa	15180
gtaggtggag	gaatagcctc	atctgtaata	gacaagatta	agagtttaat	tgaaaaataa	15240
atattaagta	aatagaaaat	aagaatgtat	ttagatataa	agttatatat	ttaaatgcat	15300
tcttatttta	ttttcattat	cttaacaatc	taataatatt	gaggttaaaa	tgtataataa	15360
tttgagctat	attaagttta	attttaaata	aagcagatat	cttttatata	tataaatata	15420
aattcaattt	tgaatatggt	taaaaagaag	ttaaaaataa	ttttttaaat	ttaataaaat	15480
aaaacactaa	tatatagacg	agatatattc	atatcatata	tactctataa	tatatggaaa	15540
ataaagagaa	aataaatgga	aaataatcac	aatataaata	taaaatataa	aaatcatcag	15600
ggggatatga	aaatgaacaa	aaaaatatta	tcattaggtc	tagcagtatc	attaattctta	15660
gtaaacttca	aaagtgtaaa	tgcatcatca	gtagtagaaa	aaatatacgg	taaagataga	15720
tatgaaacag	cagcaagatg	agctgataaa	cagacttatg	aaacagtaat	tttagttaa	15780
acagaaaaat	cacttgacga	tggattaaagt	gtaagtggat	tatcaggagc	tacaaaagct	15840
ccaatactat	ttacacaaca	aaataagata	ccagcagaca	caaatagatg	tctaaaaaat	15900
atcaaaaaag	catatataat	aggtacagaa	gatactataa	gtaaatcagt	agaaaaagaa	15960
ctagattcta	aaaatataga	agtaaaaaaga	attgggtggag	aagatagact	aaaaacaagc	16020
tatttaatag	ctaaagaaat	agcaactata	aaaaaagtgg	ataaagtact	attaactaat	16080
gcatatagt	gagaagcaga	tgcaatgagt	gtatcatcag	tagctactag	agatggagct	16140
ccaatcatac	ttacagatgg	aaagagtgtg	ccttttgatg	taaaaaatat	tcaatcatat	16200
tgtatagggt	cagaggagat	aatgagtaat	cctctagtta	aaaatacaaa	ctcagtaaga	16260
atagaaggaa	ctgaccgatt	tgaaactaac	aaaaacgtaa	tagactactt	ttttaatagt	16320
gcagatggat	tctatgtatc	agatggatac	caattagttag	atgcaatagc	agctgcacca	16380
cttactaaaa	actctccaat	ggtagtagta	aatgatggaa	gcgataaaac	tgtattagaa	16440
ggagctaaga	atataacctc	tgtagggtgaa	ataaatgaga	aggtaataca	acagtgtata	16500
aatgcttcaa	agtcaaatgg	acaacccccca	acaattacag	ttggaagtac	agaggatat	16560
aagggtgaaa	agtttgacac	tggcaagtta	aatatagtag	ctaaagataa	cacagggaa	16620
gtattaccaa	tagaagttga	tgggtttata	gatacaaaata	gagtaggtac	atatatattg	16680
acattaaaag	ctacagacga	atggggaaaa	agtacaggaa	aaagagtaga	aataaaagt	16740
ttggatgaca	aatctcatga	ttacaacagc	ccagaattta	aaaaaatggg	atctactgaa	16800
atgtataact	taatcaattc	ctatagaaaa	gaaaaaggca	aagagccttt	ggtagtgtct	16860
agtaggctag	aagggtatggc	aaatgcatgg	tctaagtata	tgatggataa	aaaagtattt	16920
gcacattata	tagatggaaa	aaatgctcca	caagtatttt	ctgagtttgg	aatgagaagt	16980
gaggaaaaata	tagcatatat	ttatatagat	tctaagaacg	ttcaaactac	acaagatgca	17040
aaagacttgg	caaaggctat	atttgaggtg	tggagaanaat	ctccagagta	caatgcaaac	17100
atgcttagtg	atgagtttta	tagtactgga	tttgggcttt	atatattatc	tgatgggtcaa	17160
gtgcatgcta	ctcaagagtt	tttaaattggc	aatgaaggta	gcttgtaaaa	aattccataa	17220
atcttgtttt	tttggttaatt	tactgatatt	ataaaatgaa	aatcacatct	aaagtacttt	17280
ttgtaatctg	ttttgttatt	gctttacagg	ttatgggtatt	ttaggtgttt	tttggttttag	17340
tagtttataa	attttgaggg	aattactact	gaggatgtaa	tttatgatta	gctgaattat	17400
ataaaaaaat	tagtaggagg	agtttgatga	aggtaataaa	gagagtatta	tcaataggat	17460
tggcaatatc	attaataatg	gcaggagcac	caaatatcaa	tgcactttca	agtagagaa	17520
agatacaagg	gaaagataga	tatgagacag	ctgccaaaaa	tgcacagaaa	caaacatatg	17580
aaaatgtaag	attagtaaat	acagataata	cactagctga	tggattaagt	gcaagtggat	17640
tagcaggaac	agttaaagca	cctatattat	tatcacaaaag	aaacagtata	ccatcagaca	17700
cagaaaaaat	gttaaaagat	gtaaagaagg	tttatataat	aggaactgaa	gattcaatag	17760
gaaaatcagt	tgaaaatgaa	ttaaagcaaa	aagggtataga	agtcaaaaga	atagggtggt	17820
atgatagaat	tgaaacaagt	tatcttatag	caaaagaaat	agcttcaata	aaacctatag	17880
ataaggtttt	tataactaat	ggttatacag	gtgaagcaga	tgctatgagt	gcgtcatcag	17940
tagcatcaag	agatggagca	cctataattt	taactaatgg	taaaaatggt	ccatttgaga	18000
aaaaagaagg	tgtgcaatgt	tatgctcttg	gttcagagga	gataataagt	aatgatttag	18060
ttaaaaaaac	taattcagtt	agatttagctg	gagaagatag	atttgagact	aataaaaaag	18120
ttataaaaaca	tttttatagt	tcagcaaaaag	aatttttatct	ttctaaagggt	tatcaattag	18180
tagatgcagt	tgctgggtct	tctatagcta	aaaatgcacc	tattgtattg	gttgatggaa	18240
atagtgaata	gtctgtactt	agaagtgcgt	ataagattac	agcactagga	ggaatagatg	18300
aaaagacgct	agaacaatgt	ttaagtgcac	catcactgga	tgcaagtgca	ccaacaataa	18360
ctgttggaag	tttaaatatt	tatcaagggtg	ataagtttga	tataagtaaa	ctgaatatcg	18420
tagcaaagga	tagcaatggt	aatgatttga	caccagagct	tataggaaat	attaatactg	18480
ataaggttgg	taaatacaaa	gttactataa	aagctactga	tattgggtgga	aagactacct	18540
caataagtgt	agaagtaaat	gtacttgaat	ataagactaa	tgatatgaat	agtagtgagt	18600
ttaaaagaat	gggttctagt	gaaatgtata	gtttgggtaaa	ttcatataga	aaagaaaaag	18660



ggaaagagcc	attacaagtg	tctgaaaatt	tacaaggaat	ggcaaattat	tgggtctaagt	18720
atatggctga	taaagggtgaa	tttgacatg	taataaatgg	taaaaatgct	gctgaagtat	18780
tttctgggtg	tattagaagt	gaagaaaata	tagcctttgt	accacttact	acaaagagta	18840
cttatacaac	aaaagatgct	agagaagtag	caaatgtcat	atttacagtt	tggaagaagt	18900
ctgataagta	taatgaaaaa	atggttgagtt	ctaagtttgc	atatactgga	tttgggttat	18960
atatattacc	agatggacaa	gtgtatgcta	ctcaggagtt	tttaaataag	taaaatttaa	19020
tttttttagtt	tattacatttt	taaaattttag	ggtataaaaa	cttgtaaact	tggagaaaaat	19080
aataattttaa	aaaaatagct	tgcaaaaaga	ataaaaaatgg	attattatag	agatgtgaga	19140
aatattagga	atatatggat	gattatttcta	tgtacataat	aaagagatgt	aatttttaata	19200
taatgttggg	aggaatttaa	gaaatgaata	agaaaaatat	agcaatagct	atgtcagggt	19260
taacagtttt	agcttcggct	gctcctgttt	ttgctgcaac	tactggaaca	caagggtata	19320
ctgtagttaa	aaacgactgg	aaaaaagcag	taaaacaatt	acaagatgga	ctaaaagata	19380
atagtatagg	aaagataaact	gtatctttta	atgatggggt	tgtgggtgaa	gtagctccta	19440
aaagtgttaa	taagaaaagcg	gacagagatg	ctgcagctga	gaagttatat	aatcttggtta	19500
acactcaatt	agataaaatta	ggtgatggag	attatggtga	tttttctgta	gattataaatt	19560
tagaaaacaa	aataataaact	aatcaagcag	atgcagaagc	aattggttaca	aagttaaatt	19620
cacttaatga	gaaaactcctt	attgatatag	caactaaaga	tacttttggga	atgggttagta	19680
aaacacaaga	tagtgaaggt	aaaaatgttg	ctgcaacaaa	ggcacttaaa	gttaaagatg	19740
ttgctacatt	tgggttgaag	tctgggtggaa	gcgaagatac	tggatatgtt	ggtgaaatga	19800
aagcaggagc	tgtagaggat	aagtatggta	aagttggaga	tagtacggca	ggtattgcaa	19860
taaatcttcc	tagtactgga	cttgaatatg	caggtaaagg	aacaacaatt	gatttttaata	19920
aaacttttaa	agttgatgta	acaggtgggt	caacacctag	tgctgtagct	gtaagtgggt	19980
ttgtaactaa	agatgatact	gatttagcaa	aatcagggtac	tataaatgta	agagttataa	20040
atgcaaaaaga	agaatcaatt	gatatagatg	caagctcata	tacatcagct	gaaaatttag	20100
ctaaaagata	tgtattttgat	ccagatgaaa	tttctgaagc	atataaggca	atagtagcat	20160
tacaaaatga	tggatatagag	tctaatttag	ttcagttagt	taatggaaaa	tatcaagtga	20220
ttttttatcc	agaagggtaaa	agattagaaa	ctaaatcagc	aaatgataca	atagctagtgc	20280
aagatacacc	agctaaaagta	gttataaaaag	ctaataaatt	aaaagatttta	aaagattatg	20340
tagatgattt	aaaaacatat	aataataactt	attcaaatgt	tgtaacagta	gcaggagaag	20400
atagaataga	aactgctata	gaattaagta	gtaaatatta	taattctgat	gataaaaaatg	20460
caataactga	taaagcagtt	aatgatatag	tattagttgg	atctacatct	atagttgatg	20520
gtcttggtgc	atcaccatta	gcttcagaaa	aaacagctcc	attattatta	acttcaaaag	20580
ataaattaga	ttcatcagta	aaatctgaaa	taaagagagt	tatgaactta	aagagtgaca	20640
ctggtataaaa	tacttctaaa	aaagtttatt	tagctgggtg	agtttaattct	atatctaaag	20700
atgtagaaaa	tgaattgaaa	aacatgggtc	ttaaagttac	tagattatca	ggagaagaca	20760
gatacgaaac	ttcttttagca	atagctgatg	aaataggtct	tgataatgat	aaagcatttg	20820
tagttgggtg	tactggatta	gcagatgcta	tgagtatagc	tccagttgct	tctcaactta	20880
aagatggaga	tgctactcca	atagtagttg	tagatggaaa	agcaaaaagaa	ataagtgatg	20940
atgctaagag	tttcttagga	acttctgatg	ttgatataat	aggtggaaaa	aatagcgtat	21000
ctaaagagat	tgaagagtca	atagatagtg	caactggaaa	aactccagat	agaataagtg	21060
gagatgatag	acaagcaact	aatgctgaag	ttttaaaaga	agatgattat	ttcacagatg	21120
gtgaagttgt	gaattacttt	gttgcaaaaag	atggttctac	taaagaagat	caattagtag	21180
atgccttagc	agcagcacca	atagcaggta	gatttaagga	gtctccagct	ccaatcatat	21240
tagctactga	tacttttatct	tctgaccaa	atgtagctgt	aagtaaagca	gttcctaaag	21300
atggtggaac	taacttagtt	caagtaggta	aaggtatagc	ttcttcagtt	ataaacaaaa	21360
tgaaagattt	attagatatg	taatataagt	tttaataaaa	ctttaaatag	aaaaaggctt	21420
ctctcatagg	aagtcttttt	tatttaaaat	aaatataaaa	taaaatagag	gctataaata	21480
gcctctatgt	tatgtgagaa	atccctaaat	aaaaagatgt	ttaattttatt	taagtataaat	21540
cattatacta	taaatcaaaa	taattgtcta	ctattatgga	tttttaattgt	ataatagtaa	21600
ttggacaata	gaaaaggagg	tacttatatg	tcagttatag	attcgatact	tgacaaagca	21660
gatgagcaag	aaataaagaa	attgaatgtt	atagtagata	aaatagatgc	attagaagat	21720
agtatgaaaa	atctatcata	tgaagaacta	aaagacatga	cagctatatt	taaaaataga	21780
cttaaaaaaag	gtgaaacctt	agatgatata	cttcagagaag	catttgctgt	agtaagagaa	21840
gtatcaaaaa	gaaaattggg	aatgcgtcaa	tatagagttc	aattaattgg	tggaaatagta	21900
atacatcaag	gaaagatagc	agagatgaaa	actgggtgaag	gtaaaaactct	agttgaagta	21960
gcaccagtat	atttaaattgc	ccttacaggt	aagggtgtac	atgtaatcac	agtaaattgat	22020
tacctggcag	aacgtgataa	ggagcttatg	agcccagttt	atgaatctct	tggaatgaca	22080
gtaggagtaa	ttatatctaa	ccaagacccta	aatataagaa	aacaacaata	taaatgtgat	22140
ataacttatg	gtacaaaatag	tgaatttggga	tttgattatt	taagagataa	tatggtgccca	22200
gatttatctc	ataaagtaca	aagggaaacta	aatttttgcta	tagtagatga	agttgactca	22260
atattaatag	atgaagccag	aactccactt	attattgcag	gagatgggtga	tgaagattta	22320



aaacttttatg	agtttagcaaa	tagcttttgta	aaaactgttta	aagaagaaga	ctttgagctg	22380
gacagaaaag	ataaaactat	agcattaaca	gcaagtggta	taagtaaagc	tgagtcattt	22440
tttgggtataa	caaacctttac	tgatataaag	aacatagaat	tatatcatca	tataaatcaa	22500
gcttttaagag	gtcataagct	tatggaaaaa	gatgttgact	atgttatttc	aaatggagaa	22560
gtaatgatag	ttgacgaatt	tacaggaaga	gtaatggatg	gtagaagata	tacagatgga	22620
cttcaccaag	ctatagaagc	aaaagaaggt	gttgagataa	agaatgaatc	taaaactatg	22680
gctactgtga	cttatcaaaa	tttcttcaga	ctatatgaaa	aactttctgg	tatgactggg	22740
actgcaaaga	cagaagaagg	ggaatttgag	tcaatctata	aattaaatgt	tgtccaaata	22800
ccaactaata	gaccagttat	tagagctgat	ttacatgata	aggtatttaa	aacagaagaa	22860
gaaaagtata	gtgctgttgt	agaagaaata	ataaggatac	ataagactag	acagccaata	22920
cttgtgggaa	cagtttctgt	tgaaaaatct	gagaagctgt	ctaaaatgct	taaaaaaca	22980
ggtattaagc	atcaagtctt	aaatgctaaa	caacatgata	aagaggcaga	gataatttct	23040
aaagccggta	aattagatgc	tataacgatt	gctacaaata	tggcaggtag	aggaacagat	23100
atttctctag	gtgcaggaga	taaagaagaa	gaacaagaag	ttaaagattt	aggtggactt	23160
tatgtttatag	gaacagaaag	acacgaatca	agaagaattg	ataatcagct	tagagggcgt	23220
tctggtcgtc	aaggagaccc	aggtacatca	agattccttg	taagtcttga	agatgatgta	23280
ataaagcttt	atggtggaaa	aactatagag	aaacttatga	agagaacaag	ttcaaacgaa	23340
aatacggcta	ttgaaagtaa	agcacttaca	agagctatag	aaagagcgca	aaaaggtgta	23400
gaaggtaaaa	attttgaaat	aagaaaaaat	gttcttaaat	atgatgatac	tattaatgag	23460
caaagaaaag	ttatatataa	tgaaagaaat	aaagtgttaa	atgatgaaga	tatacaagaa	23520
gatattcaaa	aaatggttaa	agacatcata	caagaagcag	gagaaactta	tttaattgga	23580
agaaaaagag	attattatgg	atatttttaa	catttatata	gtacatttat	gccagcagat	23640
acattactaa	tacctgggtg	agataaaaaa	agtgtccaag	agataattga	tagcacatat	23700
gaaatttcaa	aaagagttaa	tgacttgaaa	aagatgatgc	ttggtattga	taaggttgca	23760
gagtttagaaa	aaacagtact	tttaaaagtg	gttgatcaat	actggataga	tcatatagat	23820
gctatggaac	agttaaaaca	gtatataggt	cttaaatctt	atgctcaaaa	agaccctttt	23880
aaggaatatg	cttttagaagg	atatgacatg	tttgaagctt	taaaataaaa	tataagggaa	23940
gcaacagtgc	aataacttata	taaattttaac	taaaaaagtt	catgtaattt	ttattaaatg	24000
aaaaataaag	tatttttgatt	cttttatattg	taattacaat	taaataatta	aggaatacaa	24060
atgaaaaaga	agctgaaaaat	cagcttctttt	ttctgtgtga	gaaatacggg	aaaaagagat	24120
gcttgaaaatt	tctttcttcc	atttaatatatt	atactataaa	gaagatgaaa	agtctattag	24180
ttataaaactg	gattatttaa	aaaaattaca	attttgttac	ttttttgtat	acataaatat	24240
taaaaaaata	aaatataaga	aaaagtttat	atcttttggg	taattattac	aataagcttc	24300
attttattgaa	ataatatcaa	atatatatta	taattggtaa	ataaggaaaa	aataataaaa	24360
atttgaatttt	tttaggggga	aaataccatg	ataaaaaaaa	atctttctgt	aattatggct	24420
gctgcaatga	taagtacatc	agtagctcca	gttttttgctg	cagaaactac	acaggtaaaa	24480
aaagaaacaa	taactaagaa	agaagctaca	gaactagttt	cgaaagttag	agattttaatg	24540
tctcaaaagt	atacaggagg	ttctcaagtt	gggcaaccaa	tatatgaaat	aaaagttggg	24600
gagactttat	caaaattaaa	aataataact	aatatagatg	aattagagaa	attagtaaatt	24660
gctttggggag	aaaataaaga	acttatttgta	actataacag	ataaagggca	tataacaaat	24720
agtgcaaatg	aagtagttgc	agaagcaact	gaaaaaatatg	aaaattcagc	agacctttcc	24780
gctgaagcta	actctataac	agaaaaagct	aaaactgaaa	ctaattggaat	ttataaagtt	24840
gcagatgtaa	aagcttcata	tgatagtgtc	aaagataagt	tagttataac	tttaagagat	24900
aaaacagaaa	cagtaacttc	taaaactata	tatgtaggta	ttggtgatga	aaaagttgat	24960
ttaacagcaa	atccagttga	ttcaactgga	acaaattttag	acccttctgc	agaaggattt	25020
agagtaaata	aaatcgataa	actaggtgta	gcaggagcta	aaaatatattga	tgatgtccaa	25080
ttggctgaaa	taactataaa	aaatagtgac	ctaaatacag	tttcaccaca	agattttatat	25140
gatggatata	gattaaactgt	taaaggtaat	atggtagcaa	atggtacatc	aaagtcaatt	25200
agtgatattt	cagcaaaaaga	ttcagaaaca	ggaaaatata	aattttactat	taagtatact	25260
gatgcatctg	gaaaagcaac	agagcttact	gtagagagta	ctaatagaaa	agatttaaaa	25320
gatgccaaaag	ctgcattaga	aggttaattca	aagggttaaat	tgatagctgg	agatgataga	25380
tatgcaactg	cagtggctat	agcaaaaaca	acaaaatata	ctgacaacat	agttatagtt	25440
aattcaaata	aactagttga	tggattagca	gctacaccac	ttgctcaatc	taaaaaagca	25500
cctatattat	tagcatctga	taatgaaata	ccaaaagtaa	cttttagatta	tataaaagat	25560
ataattaaga	aaagcccatac	agctaaaaata	tatatagtag	gtggagaatc	agcagtatca	25620
aatacagcta	aaaagcaatt	agaatcagta	actaagaata	ttgaaagact	agctggagat	25680
gatagacata	cgacttctgt	agcagtagca	aaagctatgg	gttcttttta	agatgcattt	25740
gtagtaggtg	cgaaagggga	ggctgatgct	atgagtatag	ctgctaaagc	tgctgaactt	25800
aaggctccta	taatagtaaa	tggttggaat	gatctttcag	cagacgctat	caaattgatg	25860
gatggaaaag	agattgggat	agttgggtgg	tctaacaatg	tatctagcca	aattgaaaat	25920
caacttgccg	atattgataa	agatagaaaa	gttcaaagag	ttgaaggaga	aacaagacac	25980

gataactaatg	ctaaagttat	agaaacatat	tatggcaaat	tagataaact	atatatagca	26040
aaagatggat	atggaaataa	tggtatgcta	gtagatgcat	tagcagcagg	acctctagca	26100
gcaggtaaag	gtccaatact	tctagctaaa	actgatataa	cagactcaca	aaagaatgca	26160
cttagtaaaa	aattaaatct	tggtgcagaa	gtaactcaaa	taggtaatgg	agttgaattg	26220
acagtaatac	aaaagatagc	taaaatacta	ggttggtaat	aacttatact	gtggagaaga	26280
ttccttaatt	aatacaggaa	tcttttttcta	taattatgct	at ttatttga	aattataaca	26340
ttttataaag	taggaggatg	tagtagtggt	agatgcaact	ttattttttta	caccacatca	26400
agatgatgaa	accttaagta	tgggaagtgc	tataattgaa	catggtgaaa	aaagtgtatc	26460
acatgtcata	ttatgcacag	atggtagtaa	atctataata	agaaaagttt	tggatgatgg	26520
agggtcatgt	tcttatcata	ttaaagatgt	tcataaatac	tctttatcag	aaagtgaatt	26580
ttcaaaagat	agagatgaag	aattttaagga	tagctgtgaa	gcaatgggtg	taaaagaaag	26640
taatatacat	atagaagata	atagagcaca	tgatgggtgag	ttaagtaaag	aaaaggcaag	26700
ggaaataata	cttaagtact	tagaagaata	tccagatgca	aaagtaaaaa	ctgttactcc	26760
at ttaagct	agtggaaatac	atgaagacca	tagagcatta	ggagaagctg	catttagagtt	26820
atatagagaa	ggcaaaataa	aagattttaag	at tttatgta	gaaccatacg	actataagga	26880
ttttaaaaaa	gttaatccaa	atgtagaagt	atgggaagga	ttacctagtc	aagaggagaa	26940
gttattaagt	gcaatgaatg	catataaaaa	atggaaatcct	gaaagtggac	attatgctat	27000
aggatatcat	tcagtaaaat	ctcattttga	tgagtttagct	acaaataaaa	tacagtatgt	27060
acatgctcca	taaaatatta	cgaaagaatt	aggaggttaag	aattaatgaa	aatatcaaaa	27120
aagatagtgt	ctttgttaac	tatgacattt	ttaactgtta	catttatatgg	aaatacatct	27180
aatgcatcta	caaaagatac	attaacgggt	tctggaagat	gggaaacagc	aataaaaaata	27240
agtcaagctg	ggtggacaaa	gtctgaaagt	gctgtcttag	taaatgacaa	ttccatagca	27300
gatgctttat	cagctactcc	at ttgctaaa	gcaaaagatg	cgccaatatt	attaactcaa	27360
agtaataaat	tggatagtag	aacaaaagca	gaattaaaaa	gacttggtgt	gaaaaatgta	27420
tattttaatag	gtggttcaat	tgcatttaagt	tcagagattg	aaaagcaatt	aaatgcagaa	27480
aatataaatt	ttgaaagaat	atccggaaat	agtagatatg	atacttcttt	aaaattagct	27540
gaaaagctag	atagggaaaa	gtctatatct	aaaatagtag	tagtaaatgg	agaaaagggt	27600
cttgctgatg	cagtgaagtgt	tggagctata	gctgctcaag	aaaacatgcc	aataatactt	27660
tctgattcag	agaatggaac	tgaagtagct	gataatttta	tagatagtaa	agatatagca	27720
aaatcgtatg	taatatgttg	tacatatctt	at tttctaatt	ctgtagaaag	aagtttacca	27780
aatgcaacaa	gaatagcagg	tagtagtaga	agtgaacaa	atgcaaaaaat	tatagaagaa	27840
ttttataaag	atactgatata	aaaaaatatt	tatgtaacaa	aagatgggtac	aaagaataag	27900
aatgat ttaa	tagattcttt	agcagtgggt	gtattagcag	ctaaaaatag	ttctccaata	27960
gtattagcag	gaaataagct	tgatactact	caaaaagatg	ttttaaatac	taagattata	28020
gataaagtta	ctcaaat tgg	tggcttaggt	aatgaaaaatg	ttgtagaaga	tatactagat	28080
atacaagaag	agactaagta	tactgttgaa	actattgatg	aacttaatgc	tgctataaaa	28140
agagcagatg	caaatagat	tataaagttt	aaaccagaga	aagaaaaaac	tataaataat	28200
tcttttagta	ttgaaactaa	aaaaactggt	actattgaat	tagatggaag	atatagacaa	28260
acaataactt	tggatatacc	taatggaaaa	tttaataact	atgcagaaat	agaaggtgga	28320
gtaaaagctaa	aaaacataaa	aaatgaatca	ttagttaata	aaggaagtat	acaggatttta	28380
gatataatg	atgaaaaatgg	ttgtaaaaata	gagaatgaaa	gttctggaga	aattttggttt	28440
gttactatag	ttgaagaagc	taatgatgta	tatatagtta	atagtgggtga	tataacaaaa	28500
atatcaaata	attcttctag	tactataata	agaaattctg	gaaatattga	tacagttaca	28560
ggaaaaaaag	aacctgcaat	aagtggaaat	aaacctaaag	tcaatgatac	agaaaaagaa	28620
actaaagctg	ctagaggctt	aaatcctagg	gtagaagcat	gttctgttcc	taaaaaagat	28680
tatgttatga	taacaattcc	taattcacct	aaagattcaa	gatataaaaat	ttactataga	28740
gtagtttata	ataaacctta	tgcaatggat	gttgggtgata	aaattaatat	tggagaatgg	28800
actgttgcac	caacagatga	agagccattt	cttgaaaaag	ctaaaaatgg	ttgttatgta	28860
gaagctgttg	aggtaaatat	ttcaacgaaa	gaggtttcta	gatgggggaag	aactaacgct	28920
acagatgatg	gatttttaact	tttaaaaact	tttaaaaaca	aaaggtagac	aatttttttga	28980
ttgtctactt	tttatttttta	aaaattatca	ttttactaaa	caaataattt	aaaactataa	29040
caattatatt	agtaataact	tttacaataa	catcattttac	aaaaagcaaa	tcaatcatta	29100
catacataat	taacattttca	acgacaccag	aaagtgcct	aaaactaaca	aaagacaaaa	29160
attccttaaa	taagaacttt	at tttcaattc	ttttactttc	aaactcaaaa	aacttattag	29220
taacataagc	aaataagaca	gcaagtatcc	atgcctaaagc	at tttgcaacc	ataaaaattaa	29280
ataaaaataac	tctagtaaaa	tataaatac	aaactatatt	tacaagagta	gtaaatgctc	29340
caaagaataa	atataaatatt	gtctctttat	gttttttttaa	aatcaaatat	aatcctccta	29400
agctaattta	atataaatat	acttaagat	actaacaac	attagtatag	cacatctatt	29460
aataaatcaa	tatatcttaa	aatattataa	aaataaaact	ataatattct	ataattaata	29520
tgtactcata	atccatattt	tatcaaaaata	aaatatgacc	tttaatttaga	aatctgatat	29580
aataaaaaatt	aaatacatat	aaggggggtaa	acatgagaaa	gtataaatca	aaaaaattgt	29640

caaagctact	agcattgtcg	acagtttgc	ttttaatagt	gtcaacaata	cctgtctcag	29700
cagaaaacca	taaaactcta	gatggagtag	aaactgcaga	gtattctgaa	agctatcttc	29760
aatacctaga	agatgtcaaa	aatggagaca	cagcaaaaata	taatggagta	ataccttttc	29820
cacatgaaat	ggaaggtaca	acacttcgaa	ataagggtag	aagtagtctt	ccatcagcat	29880
ataaatcaag	tgtagcatat	aacccaatgg	atcttgggtc	tactacacca	gcaaaaaaat	29940
aaggaagtct	taatacatgc	tgggtctttt	caggtatgtc	aacttttagaa	gcatacttta	30000
aattaaaagg	atatggaacg	tatgacttat	cagaagaaca	tttaagatgg	tgggcaactg	30060
gtggaaaata	tggatggaac	ttagatgata	tgtcagggag	ttcaaagtga	acagccatag	30120
gatattctgac	agcatgggca	ggtcctaagt	tagaaaagga	cataccatat	aatctttaa	30180
ctgaggcaca	aggtgcaact	aaaccttcaa	atatgggata	tgcacctact	caatttaagt	30240
taacagatgt	tgttcgtctt	aataaagata	aggaaactgt	aaaaaatgct	ataatgcaat	30300
atggttcagt	gacatcaggt	tatgcacatt	attcaacata	ttttaataaa	gatgaaacag	30360
catataactg	tacaaataag	agggctccat	taaatcacgc	tgtagcgata	gtaggatggg	30420
atgataatta	ttcaaaaagat	aactttgcat	ctgatgtaaa	accagaatca	aatggagcat	30480
ggtttagtaaa	aagtagttgg	ggagagttta	attctatgaa	aggattcttc	tggatttctt	30540
atgaagataa	aactctacta	acagatacag	ataactatgc	aatgaaatca	gtatcaaaac	30600
cagatagtga	tacaaaaaatg	taccaacttg	aatatgctgg	tcttagcaag	ataatgtcaa	30660
ataaagtaac	agcagcaaat	gtatttgatt	ttagcagaga	ttctgaaaaa	cttgactctg	30720
ttatgtttga	aacagattct	gtaggagcaa	aatatgaagt	atattatgca	ccagtagtaa	30780
atggagttcc	tcaaaacaat	tcaatgacaa	aacttgcaag	tggaaacagta	tcatattctg	30840
gatacataaa	tgtacctact	aattcttaca	gcttaccaaa	aggtaaagga	gcaatagtag	30900
tagttataga	caacacagca	aatcctaata	gagaaaaatc	aacttttagca	tatgaaacta	30960
atatagatgc	atattatttta	tatgaggcta	aagcaaacctt	aggtgaaagc	tacatacttc	31020
aaaacaataa	gtttgaagac	ataaatacat	atagtgaatt	ttctccttgt	aactttgtta	31080
taaaagctat	aacaaaaaca	tcttctggac	aagctacttc	aggagaatct	ttaactggag	31140
cagatagata	tgaacacagca	gttaaagtta	gtcaaaaagg	atggacttct	tcacaaaatg	31200
cagtatttagt	aaatggagac	gcaatagtgg	atgctttaac	agctacacca	tttacagcag	31260
caatcgactc	tccaattctt	ttgacaggaa	aagataaactt	agattcaaaa	actaaggcag	31320
agttacaaag	atttaggaact	aaaaaagttt	atctaatagg	tggagaaaaat	tctttaagca	31380
agaatgtaca	aactcaactt	agtaatatgg	gtatatcagt	agaaaagaatt	tcaggtagtg	31440
atagatacaa	gaccagtata	tctctagctc	aaaagttaaa	tagtataaaa	tctgtttcac	31500
aagttgcagt	ggcaaatggg	gtaaatggac	ttgcagatgc	aataagtgtt	ggtgcagcag	31560
ctgctgataa	caatatgccca	ataatactta	ctaataaaaa	gagtgagttg	caaggtgctg	31620
atgagttttt	aaattcatct	aaaataacta	agtcttatat	aattgggtgg	acagctactt	31680
tatcatcaaa	tttagaaaagt	aagctttcaa	atccaacaag	acttgacagga	agtaatagaa	31740
atgaaactaa	tgctaagata	atagataaat	tctaccctag	ctcagatttg	aaatatgctt	31800
tgttggttaa	agatgggttca	aagagtcagg	gagacttgat	agatggatta	gcagtaggag	31860
cttaggtgac	taaaactgat	tcaccagtag	tcttagttgg	aaataagtta	gatgaaagtc	31920
aaaaaaatgt	acttaagtct	aagaaaatag	aaactcctat	tagagttggg	ggaaatggaa	31980
atgaaagtgc	ttttaatgaa	ctaaatactc	ttttaggaaa	atagtgaatg	gaaagatagt	32040
aaggatatac	atttttattat	taaaaaatgt	tcaatagaag	tttataaata	taaaaataca	32100
aaaaggggtg	tctcaaaatg	aacttttttag	ttcatgaggc	actctttttt	gttgttttat	32160
atttttatca	tttaataaat	gaaaaatagg	cttgtctata	tttttagata	gttttcttat	32220
tattgatgaa	aaatgtaata	attttaaaact	gtaaaatagt	gtaaagtatt	tttataatat	32280
aaatatagta	agatttttaa	caaaaataagg	gggaataata	atgaaagcac	caaaaactat	32340
tttaacaata	ctgacaatag	cgcttacttt	aagtagcatt	tctataatac	catcatatgc	32400
acttacagag	gaaaaattaa	taggtaatgg	tagatatgaa	actgctgtaa	agataagtca	32460
aaaggcatac	agttctagtc	ctaagttagt	gctagttaat	gacaattcat	tagctgatgc	32520
actttcagct	acaccatttg	caaaggctaa	aggagcacca	atacttttga	cagaaagtga	32580
taagctagat	gataggactg	aaaaagagat	aaaacgtcta	ggagctaaag	atattttatt	32640
gataggagggt	actgctgtac	ttaataaaga	tatagagaac	aagttaaaag	gtaatggact	32700
taatgtagaa	agaattaatg	gtaagaatag	atatgaaact	tcactgattt	tggctaataa	32760
gcttaaggat	ataaaaagata	ttaaggaagt	tgtgttagta	aatggtgaaa	aaggattaag	32820
tgatgctgtt	agtgttgagg	caccagcagc	tcaaaaataag	atgcctataa	ttttgtctaa	32880
ccctaaggac	ggagtagaag	catttgataa	atttataagg	gatgagaaaag	ttataaaaagc	32940
atatgtaatt	ggaggtaaaa	attctgtttc	aagagctgta	gaaaagagtc	ttcctaattgc	33000
agaaagagtg	agtgggaaag	atagaaatga	gactaatgct	aaagttatag	aaaagtttta	33060
cactgatata	aatttaagta	atttatatgt	aactaaggat	ggtagtaaaa	atgagaatca	33120
actaatagac	tcgttagctg	ttggagtatt	ggctgcaaaa	aatgagtctc	ctattgtttt	33180
ggttggcaat	aaaattaaca	caaaacaaag	agatatactt	agtactaaaa	aattaaatac	33240
tataacacag	gtaggtggaa	atggaaatga	agaagctttt	gatgaaatta	aaagtctaca	33300

```

agaaaaaact gtgtttgaag cgaaaaactgt tgaagaatta actgatatga taaatatagc 33360
aagtcctaata gatattataa acttttaaacc aaaggagaat actgtaaatg aagcttttag 33420
aatggtaaca aataaaccaa taactgtgaa tataaagggg gattgttcaa agactcttac 33480
agtagatatg cctaattggtg aagtgaataa ctatgctact ttggtaaatg ttatagttag 33540
aaatatgtga gaaggtggat ttaataatca tgatacaatc actatatgtt cagttcgtga 33600
caaaaatggg agagttatag aaaatactag aaattcagat atagatacgt tgatgatatt 33660
agcttctgct aatgatataa aattgattaa tgatggatat ataggaaaaac ttatagataa 33720
ttcatctaata agtgatataa ctaataatgg tactatagat aaaaaagtta atcagggtga 33780
agatttagag gctaaagtag attcaattga aaaagctata gattctataa gtcaaaaagt 33840
caataaaaata caagatatat tagataagct tggatttcta aagaaatttc ttagctaaga 33900
atatatatat ttaagtgaag tgtgtgtgtg aattgttcta aagttatgta gttaaacagg 33960
tttatataca aatagattta tagataaata tgtagtttag tatatatatt aataagcttt 34020
ggggagggtaa catgggggatt ggaattgaat gtaacaaaaa aataaataaa agtattgaca 34080
aaactagaat aaagaaaaat aatctaaata gcaatagcaa tagcaatagc aatagcaata 34140
gcaatagcaa tagcaatagc aatagcaata gcaatagcaa tagcaatagc aatagcaata 34200
gcaataaaaat ttcaataaat gctgtagatt ctaaacttat gaatagtagg attaaatcaa 34260
tagatataat aagagggcta tcaatagccc tcatgatagt ctgcaataat cctgggactt 34320
ggatgagaat gtatcctcaa ttaagacatg ctgtatggca tgggtgaaca cttgcagatt 34380
ttgcatttcc tttttttgtt atactcttag gtgttacaat tccaatatca ataaatagta 34440
aactaaaaaa taataaatcc acactgagta taactcttag tatattcaag agaagtatct 34500
tattgatatt atttggaatt ttcttaaaat acctaggaaa tcttgattta gatactgtaa 34560
gaatacttgg agtacttcaa cgtatgggat tagtctatct tgtgactagt ttagtatatt 34620
tactactaaa aaaattaaat gttggaagta ctgctacaat aattactttt ctatgtatat 34680
ctacattttat aatagttggg tactacatat tagcaaaaacc ttatggattt gagctagaag 34740
gttctcttgc acagtttagt gatttacatt tcttcaaggg acatttgtac aagccagaat 34800
ttgaaccaga tggtttttta actagtatag tagcaatttc ttcaggtagt ttgggatgca 34860
ctatgggatg tgtactttta aaagaagaca taggagaata taaaaaattc tttaaaatac 34920
tgggttatgag cataatcctg ctaatagggtg cttttatctt taatcagtat tttccattta 34980
ataaaagatt atggtctagc tcatttgtac ttttaatggc tggcagttat ggtatattac 35040
tgtcaatatt ttattttata tgcgacatca aaaataaaag taaaatattt accccaataa 35100
tagcacttgg aagtagccct atttttacat atatgtgcct tgaaatatta agtcatgttt 35160
tttggaaatg tccaaaactt acgaacaaag tctgattatcc aactaccttg gtagagtggg 35220
ctacatatga acttataaca ccttgggcag gtacaacttg ggattctctt atattttctt 35280
tattgtatgt cctattctgg attatagtaa tgtctataat gtacaagaaa aaaatattta 35340
ttaaatttta gtgcaatatt ttttagatgat aaagtctttg atttgttcaa atatacggaa 35400
actatcaaata ataataaaaa tttctcctcg ataaaattta aaaaccatgc cataaaagat 35460
tacaaaaagt ttacaatttg taattttttg tgctataatg aaaaaagatt tacataatat 35520
tgaatttatg taatataaaa caagcttaag atgtattaaa aaagatttaa agagttaaag 35580
aaccaaaaaa tgagggggga tatgatgaaa aaacaacaa aattattggc aacaggaatg 35640
ttgtctgtag caatgggtgg accaaacgta gcttttagcg ctgaaaatac tactgcaaag 35700
acagaatcta attcagatat taatataaat ttacaaagaa aaagtgtagt gttaggaagt 35760
aaatctaatt caagtgtaaa atttaaagaa aaattaaatg ctgattcaat aacattaaat 35820
tttatgtgtt atgatatgcc acttgaagct actttaaatt acaatgaaaa gacagattca 35880
tatgaagggtg taataaacta taacaaagac ccagaatact taaatgtatg ggaacttcaa 35940
agcataaaaa taaatggtaa agatgaacag aaagttttta acaaagaaga cttagagtct 36000
atgggatttaa atcttaagga ttatgatgtt acacaagagt ttataataag tgatgcaaat 36060
tcaacaaaag cagtaaatga atatatgaga aaaacttctg caccagtaaa aaaacttgct 36120
ggagctacta gatttgaaac tgctgttgag ataagtaaac aggggttgaa agatgggtct 36180
agtaagggtt taatagttaa tggagaatta gcagcagatg gaattacagc tactccactt 36240
gcatctacat atgacgcacc aatattactt gcaataaaag atgatatacc agaaagtaca 36300
aaagcagagt taaaacgtct aaatccaagt gatgtaataa ttataggtga tgatggttca 36360
gtttcacaaa aggcagtgtc tcaaatttaa tctgtgttaa atgtaaatgt gactcgtatt 36420
ggtggagttg atagacatga aacatcttta cttatagcaa aagaaataga taagtatcat 36480
gagttaataa aaatatacat agcaaatgga ttgtcgtgtg aatatgatgc tcttaatat 36540
tcatcaaaaag ctggtgaaga ccaacaacct ataatttag ctaataaaga ttctgtacca 36600
caaggtagat ataattgggt atcttcacaa ggttttagaag aagcttatta cataggtggg 36660
agtcaaagtt tatcatcaaa aattatagac caaatatcta aaatagcaaa aatgggtact 36720
tcaaaaaata gagtttctgg agcagataga catgaaacaa atgcaaatgt aattaagact 36780
ttctatccag ataaagagtt gagtgcaatg ttagttgcaa aatctgacat aatagttgat 36840
tcaataacag caggaccttt agcagctaaa ttaaaagctc caatacttat aactccaaaa 36900
acttatgttt ctgcatacca tagtacaacaa ttatcagaaa aaacagcaga aacagtatat 36960

```

caaattggtg	atggaatgaa	agatagtgtt	ataaattcta	tagcatctag	tttatcaaaa	37020
cacaatgcac	caacagaacc	agataattca	ggttcagcag	caggaaaaac	tgtagtgata	37080
gacccaggtc	atggggggttc	tgattcaggg	gcaacaagtg	gattaaatgg	tgagcacaa	37140
gaaaagaaat	atactcttaa	tacagcatta	gcaactactg	aatatttacg	ttcaaaaggt	37200
ataaatgttg	ttatgactag	agatacagat	aaaacaatgg	ctttaggaga	aagaacagca	37260
ttatcaaata	ctataaagcc	agatttattt	acaagtatac	attacaatgc	gtctaattgga	37320
tctggtaatg	gtgtggaaat	ttattacaaa	gtaaaagata	aaaatggtgg	aacaactaag	37380
actgcagcat	caaatatctt	aaaaagaata	ctagaaaaat	ttaacatgaa	aaatagagga	37440
atcaagacaa	gaacacttga	taatggaaaa	gattacctat	atgtgttaag	aaataacaat	37500
tatccagcaa	tacttggtga	atgtgcattt	attgacaata	agagcgatat	ggataagtta	37560
aatactgcag	aaaaagtaaa	aacaatggga	actcaaattg	gaataggaat	agaagataca	37620
gtaaaataac	atagataata	ttaaagcaaa	acatatcatc	tcctgttata	atgtaaaata	37680
taacaagaga	tgatatatctt	ttgtaaagaa	tcaaattggt	tacaatgatg	tttcttgtat	37740
tgataatctg	ctcattattg	agtataatat	agaaataagt	caaaaataag	atagaaataa	37800
gttgaaatta	agaattaaaa	aaatttttat	tccaatatgg	aggtaactat	gtctggatat	37860
actaatgatg	agtgtgaaat	acctaaaatt	aaaagctatc	caggggcaga	taaagaaatt	37920
gcaagtgaga	ttgactatag	catagtaaaa	ggaacagttt	tattcgactt	gtaccaaaaga	37980
attatggatt	tagttttatc	tataatagga	cttgtgatag	gtttaccttt	aatagcaata	38040
tttggaattc	ttattaaaaat	agaggacaaa	ggaccaataa	cttataagca	agaaagacta	38100
ggaaaatgcg	ggagaagatt	ctatatatat	aaattgagat	ctatgagaac	agacgcagaa	38160
aaatttggtg	cacaatgggc	tgaaaaagat	gatcctagaa	taactaaggt	tggaaggttt	38220
attaggaaaa	ctagaattga	tgagattcca	cagctgttta	atatattaaa	gggggacatg	38280
ggcttaatat	gtcctagacc	agagagaccg	aactttactg	ttcaatttaa	tgaggaaata	38340
cctggattta	tcaacagact	cgctataaaa	cctgggctta	caggttgggc	acaagtaaat	38400
ggtggatagc	aaatcactcc	tgaagaaaaa	cttaaagaag	atatttacta	tataaaaaat	38460
aggagtatat	tacttgactt	taagatactc	tttaaaactg	ttaaagtagt	attgacagga	38520
gacggagcta	gataatttta	aaatattaga	ttactaatca	atgacaaagc	tctctaaaaa	38580
gagctttttt	attgtcaaaa	aatacaaaaa	agaaaaggaa	tgagaactta	tgattaaaaa	38640
aatctctact	atattatcgt	tagttttggt	aatctctatt	tctagtacga	taggagtatt	38700
tgagatgca	aatcccaaaa	gagagttgat	agaaggaggt	ataccagaga	tttcgacaga	38760
attaaataaa	agagcattta	aggattctaa	agaagtaatt	cttgtaaatg	aagagtcaat	38820
tgtagatagt	ataagtgcaa	caccttttagc	atattcaaaa	aatgtctcaa	tagttgtaac	38880
taagagtaaa	aatcttggtg	gagtaacaag	aaattatcta	aaagaactag	gtccagagaa	38940
ggttacgata	gtaggtggtt	taaaggcagt	atctaaagat	gctgaacgta	atatagaaaa	39000
aatgggaatg	aaagtagaaa	gaataagagg	gaaagacaga	tatgatacat	ctttaaagat	39060
agctagagaa	atgtatagaa	cagtaggatt	tgatgaagca	tttttactta	gctcaacaac	39120
aggattagag	aacgccatat	ctgtatatct	ttatgcccgt	aaaagtggaa	tgccaataat	39180
atgggctaaa	gatgaagggt	ttgaggaaca	aattgatctt	ttaaaaggca	agaattctaa	39240
aaaaatatat	gcattagggt	attcaaaaga	atttattgca	gagattgatt	ctaacttaaa	39300
aaatatagaa	ggaataaagc	aaataaataa	gtcaagtaca	aatgttgatt	taataaagaa	39360
gttttatgat	gaaaaagata	taaagaaaaat	ctatactgct	agatttagact	ttggtagttag	39420
atctgatgta	aacgaatata	tttctttggg	agtagtatca	gcaaaagaaa	atatgccaat	39480
actgatttgt	agtataact	taagtcgtgc	acaagataaa	ttcttgaaag	atagcaatat	39540
aaatgatgta	gttgaagtgt	ggtatacagt	agggtgattat	agtctattta	agagcatatt	39600
taatctaaca	ttcttatctt	gtatagtatt	aatactatta	ttactattga	ttacatttag	39660
ggcacttaga	tatgagtcga	aatagttttt	gattttataag	gaagcttttc	atagggttaa	39720
tttacttaat	tagaataatg	tttataagca	gttttaaaag	ttctactttt	agagtttaaa	39780
ttagataagg	agaaagaaga	tgtcaaaaac	agctaaagca	gcattgtgga	ttatggcagc	39840
cactatgttt	tcaaaagtat	taggtttcct	tagagaactt	gtacttgcaa	atttctatgg	39900
gacagggtat	tatgagacg	tatttgtatt	aacattaaat	atacctggtt	taatcatagc	39960
agtcataagg	tctgcagttg	ctacaactta	tatacctatg	tactttgaga	caaagaaaag	40020
actcggagat	gaaggtgcct	taaaattttac	taataatggt	ttaaatatat	gctatataat	40080
ggctatagta	atagctatta	ttggtctttt	atttacagag	caatttggtta	cagtatttgc	40140
agcaggattt	agaaacgacc	ctgctaagtt	ccaagcagca	atattattta	ctaaaataat	40200
gatttcagga	gttttattcc	tgagtggaaag	taagatatatt	agttcgtttt	tacaggtgaa	40260
tgacagtttt	gtaattcctg	ggcttatagg	aatcccatat	aatattataa	taatagcagc	40320
aatagcacta	agtgcaggta	aaaatgtatg	gatactgcca	gcaggagcat	tattggctat	40380
ggcaggtcag	ctattgtttc	aactgccatt	tgcgtttaag	aagtcctaca	agtataagcc	40440
atatataaat	ttaaaggatg	agtctataaa	agagtttagtc	aacctagtat	tgcctagtgt	40500
ggtaggagta	gctgttggac	agttaaatat	atttgttgac	cgattattgg	caaccacttt	40560
gggtgatgga	aagttgtctg	ctcttaacta	tgcaaatagg	ctaaatgagt	ttgttatggc	40620

attatattgtt	acatctataa	taacagttat	atatcctaaa	ctagcaaaaa	tgtctggtaa	40680
agataataaaa	gaaggattta	taagcactat	tgtaaaatct	tcaaattgta	ttatattggt	40740
ggtgcttcct	atatctatag	gagctattat	attagcagag	cctcttgta	gaatattatt	40800
ccaaaggggt	aaatttgatg	cgttatcaac	agaccttaca	tctatagcgc	ttagattata	40860
ttcactaggt	ttattggctt	gtggagtaag	agatgtctta	tatagggcat	tttactctct	40920
ttcagataca	aaaacgcaa	tgataaatgg	aagtatagct	ttaataataa	atatagttct	40980
taatttaata	ttgataagac	cattgggtca	tgcagggtatt	gctatatcaa	ctagtacatc	41040
aaacataatt	acagttatat	tattattttat	atctcttaaa	aagaaaaatg	gatatttttg	41100
aggggataaa	attataaaga	ctgggtctaaa	aagtttggtg	gcatctgggtg	tgatggcagt	41160
agctacttta	ttgatataata	ataatctata	tgcattttatg	ggttcaggaa	caattaaaga	41220
aattatatca	gttggagctg	gagtattagg	tggagcttct	gtatatactg	tattgatagt	41280
tttattttaag	gtagaagaga	tggattttagc	ttttgaatttt	ttaagaaaaag	gaaaacaaaa	41340
actacttaga	agatagataa	ttaccttgaa	atttttaaatt	agagtcagaa	aggatgaaaa	41400
tatggattat	aagaataatt	atgagatgtg	gttaaattct	ccttattttg	atgaacaaac	41460
gaaaaatgag	ctattaagta	taaaagatga	tgaaaaagaa	atacaagata	gattttataa	41520
aaatttagag	tttggaaacag	gtggtttaag	aggtataata	ggagcgggaa	ctaatagaat	41580
caatatattat	actgtagac	gtgcgacact	gggcgtttta	aattatatca	tgaaaactca	41640
aggagaagaa	ggtaagcaaa	agggaaatagt	tatagcacat	gatagtagat	acatgtctag	41700
ggagttttgt	atagaagtag	caaagacttt	aagtgccttat	ggtgtaaagg	cttatatttt	41760
tgaagaatta	aagcctacac	cagaactttc	ttttgctggt	agatatttaa	agtgtgctat	41820
gggaatagta	ataacagcaa	gtcataatcc	taagaataac	aatggatata	aagtatatga	41880
ctcagatggt	ggtcaaatct	gtatagatat	ggcaaatgat	ataatagcag	aagttaataa	41940
gattgatgat	tacagtacta	ttaaaagtat	agattttgaa	gaagctttat	ctaaaaattt	42000
aataactata	ttagataatg	aagtagatga	tgagtttata	aaagctgtta	agaaacaagt	42060
tttaagacaa	aatataatag	atgaatacgg	gaaaaaatta	aaaataatat	atacacctat	42120
tcatggtaca	ggaaataagc	cagtaagaaa	agtccttaaat	gaatgtgggt	ttgaaaatgt	42180
aatggtagtt	aaggagcaag	aattgccaga	ttctaatttt	tcaacagtaa	agtatcctaa	42240
tcttgaagaa	aagtcagttg	ttaatatagc	tattgaaatg	gcaaagaata	atggcactga	42300
tttgataata	ggaacagacc	ctgattgtga	tagagtaggt	atcgttgtaa	aagattcaag	42360
tggtgaatat	gtagtcctaa	atggaaatca	agtcggttca	cttttagtta	gatataatat	42420
agagacttta	gttgaagaaa	ataaacttcc	taaaaaacaat	cctacaataa	taaaaaactat	42480
agttacatca	gaaactaggtg	caaaaatagc	aaaggcttat	aatgtagatt	gtttaaatac	42540
tttaacagga	tttaaaattta	taggagaaaa	aataaaagcg	tttgaagaaa	gcaatgatag	42600
aagttttata	atgggatatg	aggaaagcta	tggttactta	attggaactc	atgctagaga	42660
taaggatggt	gttgatctct	cacttatgat	atgtgaaatg	gcagcatact	attcttcaaa	42720
aggtatgaac	ttgtatgaag	ctttgataga	tacatataat	aaatttggtat	actacaaaga	42780
ggacttaaaag	tcagtaactt	taaaaggcat	agatggaata	aaaaaaataa	aagagatgat	42840
gctttacttt	agaagtgtta	aaatagataa	tgtggctgat	gtaaaagtag	ataaaatttt	42900
agactataaa	gatgggtgtg	atgattttacc	taagtcagat	gtactttaaat	ttttacttga	42960
agatggttct	tggatagcta	taagaccttc	aggtacagaa	cctaaaatta	aattttactt	43020
tggagcaaat	tcagacaatc	aagaagatgt	agaatttaag	ttgaacaatt	taatatctta	43080
tatatctaat	atgtgtgatt	ctatataaaa	agctaagggg	agataaaacaa	tgaaagtata	43140
caatgtcata	gttgctggag	gaggaggaac	tcgtttctgg	ccacttagta	gacaagaagt	43200
acctaaacag	cttataaatt	taagtggaga	agatgcttta	ataaacgaaa	ctataaacag	43260
aatagattct	ctagcaaaaa	aagatgattt	atttatagta	accaatgaga	agcaattgga	43320
agcttttaaag	gatatagtaa	aagataaatg	tttggtatag	aatatattgc	cagagccatg	43380
tgccagaaat	acagcagcag	ctatagggtt	tgcagcattt	aacataatga	aaaaatatgg	43440
tgatgggtgt	atgtgtgtat	atccagctga	ccattatata	aaagatgaaa	aagaatttaa	43500
atctattttg	gaaaaagcta	tctatatagc	agaaaaataa	gataaaactag	taactattgg	43560
gataactcct	acattttccat	caacaggata	tggatacata	aatttttaata	gagaaaatac	43620
tatagaagat	gttgcatatg	aggtagtggg	gtttgtagaa	aaaccaaact	atgaaattgc	43680
taaagaatat	gtaaaactcta	aaaagtattg	atgggaatagt	ggaatgttcg	tatggaaggt	43740
atctaaaata	ctagaagatt	ttaagagata	cttgccaaaa	gtatatgaaa	aattagaaga	43800
tataagtaaa	tacttaggaa	caaaagaaga	aatggaaaaa	ataaaaagaaa	tatacccaac	43860
tatacaatca	atctctatag	attatgggtat	tatggaaaga	agtaatgatg	ttatagtagt	43920
tcttgggtgat	tttggatgga	atgatgttgg	ttcttgggat	tctctaggtg	caatatatcc	43980
aactgatgat	gaaggtaata	tcaaaagagg	tgaaaaatatt	actatagata	ctaaaaattc	44040
aattatataat	tcagatgaca	aattaatatc	gacaatagggt	attagtgatt	tgatagtagt	44100
atctacaaat	gatgcagtta	tggttttag	aaaagataaa	gcctcaagat	gtcataaaaa	44160
tagtagagca	attaaaagaa	gaagatagac	aagaatatat	gtaggagggtg	ttcattttgat	44220
tcttaaaaaa	atacattatg	tatggtttgg	tgggtccaaaa	ggaaatatag	aaaatatatg	44280



tataaattct	tggaaagaaa	agcttccaga	atatgagata	gtagagtgga	atgaaaaaaa	44340
ctttgatata	gagaaagaaa	taaaaggaaa	taagtttttg	gaagaatgtt	ataaaagaaa	44400
actttgggca	ttcatatctg	attatacaag	aataaaagta	ctctatgaac	aaggtggagt	44460
ttatatggat	acagatatgc	aaatattaaa	ggatataaact	cctcttttag	aaaataatag	44520
gctaattctgt	gggtatgaag	atgatagaga	gtatataaat	ggtgccataa	taggtgtaga	44580
aaaaggacat	ccttttttaa	aagacttatt	ggagtattat	gaaaaagaag	tacttacttc	44640
ttcattgttt	acaataccaa	agataatgac	tcatctttatg	gaaaaagaatt	ataagaagat	44700
tgacccaaat	aattatgaag	aaggaatacag	tgtttacgac	aaagaatatt	tttatccttt	44760
tggattttaa	gaagacttta	caccagaatg	tataacagaa	aacacttttg	ggatacattg	44820
gtggggaaaa	agttgggcta	aaaagagaaa	ttatttctta	gaaagcaaac	acttaacagg	44880
tgtaaataag	atatggaagt	gttgtaaaat	atgtgcaagt	aatactttac	gaagctaaaa	44940
gggagaaagt	aaaatgaaga	ctttaaatga	tgatagaaat	attatgatta	gaaaaatatct	45000
ttatatattt	gttttggtta	tcatattact	tagaaagtat	atgatatcat	ttattgaccc	45060
aaatatggat	atagggatga	ttaaatcagc	gttattttat	tcttcaatag	gaataactaat	45120
gctacttttt	ttatttgata	agagaaaaag	tataacagaa	atggtactag	ttggagtgtg	45180
tgtcttactt	tatttattaa	atagggaagg	agctatatta	ttaatcgttt	tattatattt	45240
atcagctaaag	caaatagatg	ataaatttat	agttaaaaat	tattttaataa	tatcagcatg	45300
ttttttaaag	gtagctatat	tattatttaa	tctatttcct	tcattgatat	ttaatcagga	45360
agtcactat	agatatatag	aaaaaataga	tatgcttggt	acaagaatgg	attttggact	45420
tggtaatcca	aatagtgtat	attatcatat	ggttactata	tatgcagcat	atatattctt	45480
aagatataaa	gattataata	aatgggatag	aataatttta	tttggttctg	cattttttgt	45540
ttatcaaacc	acatatagta	gaactggatt	tttactata	ttagcaggat	taatatttgt	45600
agaaataata	agatggatag	atataaaaaa	aataaaagga	ctaccaatgc	tcctaaagac	45660
attaccgata	atacttactt	tatttagtgt	aattattggt	acagtatttg	ataaaagcac	45720
tttattaaat	agattattag	caagtagacc	taagttctgg	catgtatact	tggctgaaaa	45780
ggggaacttt	ctaaaacctt	ttggaaattc	atatagtcct	gctataaaaag	caactaatcc	45840
attagatagt	tcttatgtat	atataattag	tgtactagga	gtagttgctt	gtatattatt	45900
tatgtatttg	atgtataaag	gaatagaaag	ctttatagaa	aaagataaaa	aagcatattt	45960
agtagcttta	tttatatttt	tactatactc	ttttgcagaa	aatttattat	tagaagcttc	46020
atthagtttt	gcagtagtat	tactaataaa	agaagtgata	ttaaatgata	aaagggaaat	46080
agatttgtgg	aagatgaaaa	gtaggaggta	gtaatgttaa	tatcattaat	aatgccaaac	46140
ttaaatagat	atgatgatat	atacttactt	atggacagct	tagaaaaatca	aacttacaaa	46200
aactttgaac	ttatagtagt	agacccaaat	gataatagca	aagttaaaga	aatagttgat	46260
aaatacattg	ataagttaga	tataaagtat	ataaaaagct	ctaaaaaagg	attgagttat	46320
aacagaaatg	taggcataga	caatgctgta	gggcaaatta	taggtttccc	tgatgatgat	46380
tgtgtctacg	aaaatgatac	attagaaaaa	gttataaaat	tctttaataa	aaataaagat	46440
tataaaatat	acagttgcaa	aactatggat	tcaaataaag	ttgatgcatt	taagaaaaatg	46500
tatgatggaa	cttgtgatat	cacaagttcc	aacgtttttg	atactataac	atccataact	46560
ttttttattg	attttgaagg	taaagactat	acaagatttg	atgagaaatt	gggtgttggt	46620
ggagagtttg	gtgctgggta	agagatagat	tatgttttaa	acttattgag	tttaggtttt	46680
aaaggaaaaat	actttggaaa	tgatataatt	taccatcctg	caaaaaaaca	ttctaaatct	46740
aaagaaaagt	atcaaaaaga	ttataactat	ggaagaggct	ttggagcgct	ttgcaaaaag	46800
gaaatagtct	atagaaaaaa	ttataagttt	gcaaaagtta	tggtatcaaa	acttgttaaga	46860
aatataggag	gtcttatatt	gagttccaat	agagattatc	atagtgccac	tataaaagga	46920
agaattaatg	gatttagaca	gtacaagtta	taatgaggat	ggtgttttatg	gacaagggtta	46980
aatttttaat	aaatatgctt	ttggcttttt	ttatatatcc	atttaataag	cataagttta	47040
atggtagaga	tatctggctt	ataggtggac	attcagggga	tataataaat	gataattcaa	47100
agttttttta	tgagtatatg	ctaaaagaac	ataatgatgt	agaaacttat	tgggtagttaa	47160
ataaagacag	taaagtattt	gataaaaattc	caggcaaaaa	attaataaga	ggaagtgtag	47220
aaaactatct	gtattattac	aattcaaaag	ctatagtatt	ttcacatgca	ccatcagcag	47280
atatagcacc	atacaatttt	gctgtacctg	ttcttaatta	ttttcataaa	aaaacgataa	47340
aagtgttttt	aaatcatggg	actataagtt	ttaaaaaaag	aaaacctatg	aacaaaaagt	47400
tcaaaaatat	tatagataac	ctatataaaa	gctataatat	agttacagca	agttctgaat	47460
ttgaaagaaa	tgttatggta	aatgattggg	gaatgtttaga	tgactcagtt	tatataattg	47520
gaaatgcaag	atatgacaat	ttaccaacaa	atgaagttgc	acaaactcga	gatataattat	47580
atactcctac	atggagagat	tggataaaaat	ttagttcagg	aaaattttacg	gatacagatt	47640
attttaaaaa	tataatgaat	tttttaaatg	atgacaaaat	aaataaaaata	ttagatgaga	47700
aagatataaa	tgtaaaaatt	tatatgcac	atcttatgca	tgagtttata	gatgatataa	47760
aagaaaaat	aacaggaaaa	cgaatttgtgt	ttttagataa	aggggtaact	ctagcaaatg	47820
aatcagaaa	atcagctgca	aatataactg	attattcaag	tgtagccata	gactttttgt	47880
atatgaatag	acctattttg	ttttaccaat	tcgattttgga	tgaatatatg	gaaaaagttg	47940

```
attcatatat agattttaaaa agtgaaatgt ttgggtcttt ggcttataat aatgatgaag 48000
ccgtgaataa acttattgat attattgaaa ataattttga ggttatggat aaccaaaaaa 48060
atgaaagaaa taagtttttc agatataatg ataataagaa ctgcaagagg atatatgatt 48120
gtgtattaag taaaattaaa taaatatattt ataaatgatg aggaaggtaa tatgaagaag 48180
aatttagtat ctataattac tcccatgtat aattctgaaa aatttattga agcaaccata 48240
aaatcagtat taaaccaaac ttatcaagaa tgggaaatgt taattattga tgattgctca 48300
acagataata gtcctaatat agtcaaactt tatatgcaac aggatagtag aataaaatgt 48360
ataaagactg agactaataa ggggtgtctt aatgctagaa atttagcact aagtaaggca 48420
acaggacaat ttatagcttt tttangtagt gatgaccaat ggaatagtag taagttagaa 48480
aaacaagtaa attttatgtt agaaaatgac tatgtaattt catttacttc atatgaactg 48540
atggatgaaa a                                     48551
```



## INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 01/01280

## A. CLASSIFICATION OF SUBJECT MATTER

IPC7: C12N 15/63, A61K 39/08

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: A61K, C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

BIOSIS, WPI DATA, PAJ, EPO-INTERNAL

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relévant to claim No.
Y	WO 9519371 A2 (SOLVAY), 20 July 1995 (20.07.95), page 12, line 1 - line 9, claim 12, abstract --	1-33
Y	Microbial Pathogenesis, Volume 28, 2000, Marina Cerquetti et al, "Characterization of surface layer proteins from different Clostridium difficile clinical isolates" page 363 - page 372 --	1-33
P,Y	Molecular Microbiology, Volume 40, No 5, 2001, Emanuela Calabi et al, "Molecular characterization of the surface layer proteins from Clostridium difficile" page 1187 - page 1199 --	1-33

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

## \* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier application or patent but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search

28 Sept 2001

Date of mailing of the international search report

02 -10- 2001

Name and mailing address of the ISA/

Swedish Patent Office  
Box 5055, S-102 42 STOCKHOLM  
Facsimile No. +46 8 666 02 86

Authorized officer

Carolina Gómez Lagerlöf/BS  
Telephone No. +46 8 782 25 00

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 01/01280

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,Y	Infection and Immunity, Volume 69, No 5, May 2001, Tuomo Karjalainen et al, "Molecular and Genomic Analysis of Genes Encoding Surface Anchored Proteins from Clostridium difficile" page 3442 - page 3446  -- -----	1-33

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/SE01/01280**Box I** Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 30-32  
because they relate to subject matter not required to be searched by this Authority, namely:  
see next sheet
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

**Box II** Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/SE01/01280

Claims 30-32 relate to methods of treatment of the human or animal body by surgery or by therapy (PCT Rule 39.1(iv)). Nevertheless, a search has been executed for these claims. The search has been based on the gene expression cassette according to claims 1-10.

**INTERNATIONAL SEARCH REPORT**  
Information on patent family members

International application No.  
**PCT/SE 01/01280**

Patent document cited in search report				Publication date		Patent family member(s)		Publication date	
WO	9519371	A2	20/07/95	AU	6380194	A		24/10/94	
				EP	0691845	A		17/01/96	
				EP	0738278	A		23/10/96	
				GB	2291594	A,B		31/01/96	
				GB	9400650	D		00/00/00	
				GB	9519866	D		00/00/00	
				JP	8508474	T		10/09/96	
				JP	9508012	T		19/08/97	
				US	5874267	A		23/02/99	
				US	6028098	A		22/02/00	
-----									



**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☒ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**